Strategies to control community-associated antimicrobial resistance among enteric bacteria and methicillin-resistant *Staphylococcus aureus* in Canada – executive summary

Jeff Wilson DVM DVSc PhD¹, John Conly MD², Thomas Wong MD MPH FRCPC³,
Gayatri Jayaraman MPH PhD^{3,4}, Jan Sargeant DVM MSc PhD⁵, Andrew Papadopoulos MBA⁵,
Virginia Young MSc¹, Melanie Quist-Moyer MSc¹, Sharon Bauer MSc¹

Resistance to antimicrobial drugs is a concern that exists worldwide and has a significant impact on human and animal health. Knowledge and practice gaps exist around the control of antimicrobial-resistant infections in Canada, particularly in the community setting. Although much research exists on the control of hospital-acquired resistant infections, currently no comprehensive synthesis or review of the literature exists on the control of antimicrobial-resistant organism infections within the community. In particular, there is little synthesis of information on those infections that represent a large component of community-level impact, namely resistant enteric bacteria and community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA); these infections pose a significant health burden to Canadians.

In addition to representing a significant human health impact within the Canadian community, enteric bacteria and MRSA have similar spread and control mechanisms (eg, hygiene and handwashing, sanitation, housing density and crowding, person-to-person spread and animal exposure), thus representing areas for common policy, intervention and other control activities.

The objective of the present study was to conduct a formal, comprehensive review of control strategies and interventions available to reduce both the development of antimicrobial-resistant enteric bacteria, specifically *Campylobacter* species, *Salmonella* species, verocytotoxinogenic *Escherichia coli*, *Shigella* species and CA-MRSA, and the spread of such infections within Canadian communities.

A comprehensive review was undertaken of the relevant peer-reviewed and grey English language literature from 1970 to the present using a protocol based on a systematic review methodology. A total of 1467 references were identified; of these, 563 met the abstract relevancy screening criteria and of these, 203 were reviewed in detail. Consequently, 66 references were reviewed, from which data were abstracted specifically regarding CA-MRSA, and 57 references were reviewed, from which data were abstracted specifically regarding enteric pathogens.

In general, there was reasonable scientific evidence regarding risk groups and risk factors for CA-MRSA. This information

provided some insight into potential approaches to control these infections. Identified potential risk groups and risk factors included the following:

- Children;
- Specific ethnic groups;
- Athletes;
- Drug use;
- Men who have sex with men;
- Heterosexual high-risk activities;
- Military;
- Veterinarians and animal handlers;
- HIV infection;
- Tattoo recipients;
- Living with a carrier or case of CA-MRSA;
- Emergency departments and hospitalized patients; and
- Antibiotic use

There was a paucity of information on risk groups and risk factors for the community-acquired, antimicrobial-resistant enteric bacterial infections studied; some information exists on risk settings and risk factors for community-acquired enteric bacterial infections (without reference to resistance), but the extent to which this can be extrapolated to resistant infections is unknown; much of this information takes the form of outbreak reports. Potential risk groups and risk factors for community-acquired, antimicrobial-resistant enteric infections included the following:

- Daycare centres;
- Schools;
- Households;
- Nursing homes;
- Immunocompromised individuals;
- Specific sociodemographic factors (ie, rural residents, specific ethnic groups, income, education and access to health services);
- Population density; and
- Season

There was a paucity of scientifically based information (randomized controlled trials or observational studies) on

This is the executive summary from a comprehensive review with the same title. To access the full review, please visit <http://nccid.ca/en/amr-reviews>

¹Novometrix Research Inc, Moffat, Ontario; ²Department of Medicine, University of Calgary, Calgary, Alberta; ³Centre for Communicable Diseases and Infection Control, Public Health Agency of Canada; ⁴Department of Epidemiology and Community Medicine, University of Ottawa, Ottawa; ⁵Ontario Veterinary College, University of Guelph, Guelph, Ontario

Correspondence: Dr Jeff Wilson, Novometrix Research Inc, RR #1 Moffat, Ontario L0P 1J0. Telephone 519-824-7771, fax 519-822-5362, e-mail jeff_wilson@novometrix.com

Reprints: National Collaborating Centre for Infectious Diseases, 413-445 Ellice Avenue, Winnipeg, Manitoba R3B 3P5.

Telephone 204-943-0051, fax 204-946-0927, e-mail nccid@icid.com

interventions for CA-MRSA or for the enteric bacterial infections studied (resistant or otherwise). A number of randomized controlled trials have shown the effectiveness of handwashing in the prevention of gastrointestinal illness.

A limited number of intervention studies of hospital-acquired MRSA exist; however, the extent to which the results of these studies can be extrapolated to CA-MRSA is unknown. Such extrapolation may be misleading.

There was extensive literature concerning recommendations, guidelines and suggested approaches to the control of both CA-MRSA and, to a lesser extent, for enteric bacterial infections in community settings. Although the efficacy of these approaches is plausible, it has, for the most part, not been formally evaluated. Proposed approaches to control these infections included the following:

- CA-MRSA
 - Hand and personal hygiene;
 - Prudent use of antibiotics;
 - Decolonization;
 - Early diagnosis and appropriate treatment;
 - Education programs (hygiene and antibiotic use);
 - Regular cleaning and laundering in households and facilities;
 - Equipment disinfection; and
 - Exclusion of those with active infection from certain high-risk settings.
- Community-acquired enteric bacterial infections
 - Hand, household and institutional hygiene;
 - Equipment disinfection in high-risk settings;
 - Public and public health education;

- Early diagnosis and appropriate treatment; and
- Exclusion of those with active infection from certain high-risk settings

We concluded that formal evaluations of the efficacy of strategies for the control of CA-MRSA and community-acquired enteric infections (resistant and otherwise) are warranted, and should form the basis for public health guidelines and policy. Until such time as evaluations can be undertaken, recommendations for the control of these infections must be largely dependent on historical practice, conventional wisdom, extrapolation from other contexts, consensus and conjecture.

Potential interventions that would warrant formal evaluation in various settings and groups include the following:

- Hand and personal hygiene;
- Prudent use of antibiotics;
- Decolonization;
- Early diagnosis and appropriate treatment;
- Public education programs (hygiene and antibiotic use);
- Regular cleaning and laundering in households and facilities;
- Equipment disinfection; and
- Exclusion of those with active infection from certain high-risk settings

Ongoing collection and evaluation of information (including surveillance and epidemiological studies) on the occurrence, settings, risk factors and risk groups for CA-MRSA and resistant enteric infections are warranted. Such information will be useful in determining disease trends, identifying risk groups, settings and risk factors, and in identifying and evaluating potential interventions.