Emerging sequence type-269 serogroup B *Neisseria meningitidis* has the potential to cause outbreaks. *Can J Infect Dis Med Microbiol* 2014;25(1):8.

To the Editor:

Invasive meningococcal disease (IMD) is a serious life-threatening infection with an average case fatality rate of approximately 10%. Conjugated capsular polysaccharide vaccines against serogroups A and C have successfully reduced the incidence of disease and carriage due to these two common serogroups of meningococci in Africa and Western countries (Europe, North America and Australia), respectively (1,2). Conjugated vaccines against serogroups Y and W135 have also been developed and licensed as part of a tetravalent A,C,Y,W135 conjugated vaccine (3). Because control of IMD due to serogroups A, C, Y and W135 meningococci has become feasible, serogroup B *Neisseria meningitidis* (MenB) is now the leading cause of IMD in many industrialized countries (4,5). Hypervirulent clones associated with MenB diseases have been described as electrophoretic type-5 (sequence type [ST]-32), cluster A4 (ST-8), electrophoretic type-37 (ST-11) and lineage 3 (ST-41/44) (6). In the past decade, MenB strains of ST-269 or ST-269 clonal complex (cc) have emerged as common causes of MenB disease. For example, in England and Wales, ST-269 cc was responsible for approximately one-third of MenB cases between 2006 and 2011 (7). However, control of MenB disease has proven to be difficult because the serogroup B capsular polysaccharide is nonimmunogenic due to its antigenic similarities to host components (8).

The ST-269 cc is a large group of *N meningitidis* strains that are grouped together according to the definition of isolates sharing four or more of their seven housekeeping gene alleles when genotyped using multilocus sequence genotyping (MLST). Studies that analyzed data from the *N meningitidis* MLST website (<http://neisseria.org/nm/typing/mlstdb/>) reported that most isolates of the ST-269 cc were MenB and were more often involved in invasive disease than in normal carriage (9,10). In addition, the ST-269 cc found in the database comprised a total of 319 isolates with 271 unique STs when data were assessed on January 22, 2008 (9), and 415 isolates with 312 unique STs when data were assessed on February 6, 2009 (10). A search of the *N meningitidis* MLST database performed on June 7, 2013 revealed a total of 439 unique STs. This suggests either rapid expansion of this cc, heightened awareness of the importance of this cc as a cause of invasive disease or increased surveillance of MenB disease isolates as a result of the relative shift in the prevalence of MenB over MenC subsequent to the introduction of MenC conjugate vaccines in different countries.

Since the first description of ST-269 cc in the early 1970s (11,12), there have been four reports of clusters of cases or outbreaks of MenB disease due to strains of this cc (12-15). In three instances, the strains involved belonged to ST-269; in the fourth, the exact ST was not described but was reported to be ST-269 cc. In the endemic cluster of ST-269 MenB cases in Belgium (13), 20 of 41 cases (49%) occurred among individuals 15 to 24 years of age, and another 10 (24%) patients were between five and 14 years of age. In the community outbreak in Landes, France (14), 36% and 18% of the patients were between 20 and 24 years of age, and 15 to 19 years of age, respectively. In Quebec, of the 120 laboratory-confirmed ST-269 MenB cases detected between 2003 and 2008, 30% occurred among individuals 15 to 19 years of age; but by 2009 to 2011, the clone had spread to cause disease in other age groups, including those younger than five years of age (who represented 33% of the cases) (16). In the most recent outbreak, at a university in Ohio, United States (15), all 13 cases of MenB ST-269 occurred

among individuals 18 to 23 years of age. The ST-269 MenB clone in Quebec has also been described as replacing other circulating cc's in the geographical regions where the clone was observed to persist (16). This suggests that ST-269 either has a higher virulence or a better propensity for spread. The stability of the ST-269 clone and/or prolonged nature of the outbreak reported in three of the four outbreaks also support hyperinvasiveness of ST-269 compared with other STs in this cc (12,13,15,16).

A new genome-derived and protein-based multicomponent MenB vaccine (Bexsero; Novartis, USA) has recently received approval for use by the European Medicines Agency (17). Whether Bexsero can prevent IMD due to ST-269 MenB, other endemic MenB or other serogroups of meningococci will be a much-anticipated outcome.

ACKNOWLEDGEMENT: RSWT thanks Dr Shelley Deeks, of Ontario Public Health and University of Toronto (Toronto, Ontario), for useful comments and discussion. This study made use of the *Neisseria* Multi Locus Sequence Typing website (<http://pubmlst.org/neisseria/>) developed by Keith Jolley and Man-Suen Chan and based at the University of Oxford (Oxford, United Kingdom). The development of this site has been funded by the Wellcome Trust and European Union.

DECLARATIONS: Raymond Tsang, Frances Jamieson and Brigitte Lefebvre have no conflicts of interest to declare. Rodica Gilca received a research grant for an unrelated study from Pfizer.

Raymond SW Tsang PhD
Vaccine Preventable Bacterial Diseases
National Microbiology Laboratory, Public Health Agency of Canada

Frances B Jamieson MD
Public Health Ontario Laboratory, Public Health Ontario
Faculty of Medicine, University of Toronto
Toronto, Ontario

Brigitte Lefebvre PhD
Laboratoire de santé publique du Québec,
Institut national de santé publique du Québec
Ste-Anne-de-Bellevue

Rodica Gilca MD
Institut national de santé publique du Québec
Centre de recherche du CHU de Québec
Laval University
Laval, Québec

REFERENCES

- Novak RT, Kambou JL, Diomandé FVK, et al. Serogroup A meningococcal conjugate-vaccination in Burkina Faso: Analysis of national surveillance data. *Lancet Infect Dis* 2012;12:757-64.
- Trotter CL, Andrews NJ, Kaczmarski EB, Miller E, Ramsay ME. Effectiveness of meningococcal serogroup C conjugate vaccine 4 years after introduction. *Lancet* 2004;364:365-7.
- National Advisory Committee on Immunization (NACI). An advisory committee statement (ACS) on conjugate meningococcal vaccine for serogroups A, C, Y and W135. *Canada Communicable Disease Report* 2007;33(ACS3):1-24.
- Harrison LH, Trotter CL, Ramsay ME. Global epidemiology of meningococcal disease. *Vaccine* 2009;27:851-63.
- Racloz VN, Luiz SJ. The elusive meningococcal meningitis serogroup: A systematic review of serogroup B epidemiology. *BMC Infect Dis* 2010;10:175.

Continued on page 38

Continued from page 8

REFERENCES – CONTINUED

6. Caugant DA. Population genetics and molecular epidemiology of *Neisseria meningitidis*. APMIS 1998;106:505-25.
7. Ladhani SN, Flood JS, Ramsay ME, et al. Invasive meningococcal disease in England and Wales: Implications for the introduction of new vaccines. Vaccine 2012;30:3710-6.
8. Finne J, Leinonen M, Makela PH. Antigenic similarities between brain components and bacteria causing meningitis. Implications for vaccine development and pathogenesis. Lancet 1983;322:155-7.
9. Caugant DA. Genetics and evolution of *Neisseria meningitidis*: Importance for the epidemiology of meningococcal disease. Infect Gen Evolu 2008;8:558-65.
10. Caugant DA, Maiden MCJ. Meningococcal carriage and disease – population biology and evolution. Vaccine 2009;27S:B64-B70.
11. Law DSK, Lorange M, Ringuette L, et al. Invasive meningococcal disease in Quebec, Canada, due to an emerging clone of ST-269 serogroup B meningococci with serotype antigen 17 and serosubtype antigen P1.19 (B:17:P1.19). J Clin Microbiol 2006;44:2473-9.
12. Zhou J, Lefebvre B, Deng S, et al. Invasive serogroup B *Neisseria meningitidis* in Québec, Canada, 2003-2010: Persistence of the ST-269 clone since it first emerged in 2003. J Clin Microbiol 50:1545-51.
13. Bertrand S, Van Meerveme E, De Baere T, et al. Detection of a geographical and endemic cluster of hyper-invasive meningococcal strains. Microbes Infect 2011;13:684-90.
14. Delisle E, Larrieu S, Simoes J, et al. Community outbreak of group B meningococcal disease in southwest France – December 2008 to September 2009. Euro Surveill 2010;15:19655.
15. Mandal S, Wu HM, MacNeil JR, et al. Prolonged university outbreak of meningococcal disease associated with a serogroup B strain rarely seen in the U.S. Clin Infect Dis 2013;57:344-8.
16. Gilca R, Deceuninck G, Lefebvre B, et al. The changing epidemiology of meningococcal disease in Quebec, Canada, 1991-2011: Potential implications of emergence of new strains. PLoS One 2012;7:e50659.
17. Iacobucci G. First meningitis B vaccine could be available in UK next year. Br Med J 2012;345:e7867.