Research Article

Compatibility and Conjugacy on Partial Arrays

S. Vijayachitra¹ and K. Sasikala²

¹Department of Science and Humanity, Sathyabama University, Chennai, India

²Department of Mathematics, St. Joseph's College of Engineering, Chennai, India

Correspondence should be addressed to S. Vijayachitra; vijayachitra@live.com

Received 27 March 2016; Revised 28 July 2016; Accepted 3 August 2016

Academic Editor: Qi Dai

Copyright © 2016 S. Vijayachitra and K. Sasikala. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Research in combinatorics on words goes back a century. Berstel and Boasson introduced the partial words in the context of gene comparison. Alignment of two genes can be viewed as a construction of two partial words that are said to be compatible. In this paper, we examine to which extent the fundamental properties of partial words such as compatibility and conjugacy remain true for partial arrays. This paper studies a relaxation of the compatibility relation called k -compatibility. It also studies k -conjugacy of partial arrays.

1. Introduction

The genetic information in almost all organisms is carried by molecules of DNA. A DNA molecule is a quite long but finite string of nucleotides of 4 possible types: a (for adenine), c (for cytosine), g (for guanine), and t (for thymine). The stimulus for recent works on combinatorics is the study of biological sequences such as DNA and protein that play an important role in molecular biology [1–3]. Sequence comparison is one of the primitive operations in molecular biology. Alignment of two sequences is to place one sequence above the other [2, 4] in order to make clear correspondence between similar letters or substrings of the sequences. Partial words appear in comparing genes. Indeed, alignment of two strings can be viewed as a construction of two partial words that are compatible. The compatibility relation [5] considers two arrays with only few isolated insertions (or deletions). In some cases, it allows insertion of letters which relate to errors or mismatches. A problem appears when the same gene is sequenced by two different labs that want to differentiate the gene expression. Also, when the same long sequence is typed twice into the computer, errors appear in typing.

Partial array A of size (m, n) over Σ , a finite alphabet, is partial function $A : Z_+^2 \rightarrow \Sigma$, where Z_+ is the set of all positive integers. In this paper, we extend the combinatorial properties of partial words to partial arrays. Also, this paper

studies a relation called k -compatibility where a number of insertions and deletions are allowed as well as k -mismatches. The conjugacy result [6] which was proved for partial words is extended to partial arrays. k -Conjugacy of partial arrays is discussed.

2. Preliminaries on Partial Words

In this section, we give a brief overview of partial words [7].

Definition 1. Partial word u of length n over A , a nonempty finite alphabet, is partial map $u : \{1, 2, \dots, n\} \rightarrow A$. If $1 \leq i \leq n$, then i belongs to the domain of u (denoted by $\text{Domain}(u)$) in the case where $u(i)$ is defined, and i belongs to the set of holes of u (denoted by $\text{Hole}(u)$), otherwise.

A word [8–10] is a partial word over A with an empty set of holes.

Definition 2. Let u be a partial word of length n over A . The companion of u (denoted by $u_\diamond$) is map $u_\diamond : \{1, 2, \dots, n\} \rightarrow A \cup \{\diamond\}$ defined by

$$u_\diamond(i) = \begin{cases} u(i) & \text{if } i \in \text{Domain}(u) \\ \diamond & \text{otherwise.} \end{cases} \quad (1)$$

The symbol $\diamond$ is viewed as a “do not know” symbol. Word $u_\diamond = ba\diamond ab\diamond$ is the companion of the partial word. The length of the partial word is 6. $D(u) = \{1, 2, 4, 5\}$. $H(u) = \{3, 6\}$.

Let u and v be two partial words of length n . Partial word u is said to be contained in partial word v (denoted by $u \subset v$), if $\text{Domain}(u) \subset \text{Domain}(v)$ and $u(i) = v(i)$ for all $i \in \text{Domain}(u)$. Partial words u and v are called compatible (denoted by $u \uparrow v$), if there exists partial word w such that $u \subset w$ and $v \subset w$ (in which case we define $u \vee v$ by $u \subset u \vee v$ and $v \subset u \vee v$ and $\text{Domain}(u \vee v) = \text{Domain}(u) \cup \text{Domain}(v)$). As an example, $u_\diamond = aba\diamond\diamond a$ and $v_\diamond = abab\diamond a$.

The following rules are useful for computing with partial words:

- (i) *Multiplication*: If $u \uparrow v$ and $x \uparrow y$, then $ux \uparrow vy$.
- (ii) *Simplification*: If $ux \uparrow vy$ and $|u| = |v|$, then $u \uparrow v$ and $x \uparrow y$.
- (iii) *Weakening*: If $u \uparrow v$ and $w \subset u$, then $w \uparrow v$.

Lemma 3. Let u, v, x, y be partial words such that $ux \uparrow vy$.

- (i) If $|u| \geq |v|$, then there exist partial words w, z such that $u = wz$, $v \uparrow w$, and $y \uparrow zx$.
- (ii) If $|u| \leq |v|$, then there exist partial words w, z such that $b = wz$, $v \uparrow w$, and $x \uparrow zy$.

Definition 4. Two partial words u and v are called conjugate, if there exist partial words x and y such that $u \subset xy$ and $v \subset yx$.

Definition 5. Two partial words u and v are called k -conjugate, if there exist nonnegative integers k_1, k_2 whose sum is k and partial words x and y such that $u \subset_{k_1} xy$ and $v \subset_{k_2} yx$.

3. Preliminaries on Partial Arrays

This section is devoted to review the basic concepts on partial arrays [11].

Definition 6. Partial array A of size (m, n) over Σ , a nonempty set or an alphabet, is partial function $A : Z_+^2 \rightarrow \Sigma$, where Z_+ is the set of all positive integers. For $1 \leq i \leq m$, $1 \leq j \leq n$, and if $A(i, j)$ is defined, then we say that (i, j) belongs to the domain of A (denoted by $(i, j) \in D(A)$). Otherwise, we say that (i, j) belongs to the set of holes of A (denoted by $(i, j) \in H(A)$).

An array [5] over Σ is a partial array over Σ with an empty set of holes.

Definition 7. If A is a partial array of size (m, n) over Σ , then the companion of A (denoted by $A_\diamond$) is total function $A_\diamond : Z_+^2 \rightarrow \Sigma \cup \{\diamond\}$ defined by

$$A_\diamond(i, j) = \begin{cases} A(i, j) & \text{if } (i, j) \in D(A) \\ \diamond & \text{otherwise,} \end{cases} \quad (2)$$

where $\diamond \notin \Sigma$.

The bijectivity of map $A \rightarrow A_\diamond$ allows defining the catenation of two partial arrays in a trivial way.

Example 8. Partial array $A = \begin{pmatrix} b & a & b \\ \diamond & a & b \\ b & \diamond & b \end{pmatrix}$ is the companion of partial array A of size $(3, 3)$, where $D(A) = \{(1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (3, 1), (3, 3)\}$, $H(A) = \{(2, 1), (3, 2)\}$.

Let

$$X = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix}, \quad (3)$$

$$Y = \begin{pmatrix} b_{11} & \cdots & b_{1n'} \\ \vdots & & \vdots \\ b_{m'1} & \cdots & b_{m'n'} \end{pmatrix}.$$

By column catenation, we mean

$$X \oplus Y = \begin{pmatrix} a_{11} & \cdots & a_{1n} & b_{11} & \cdots & b_{1n'} \\ \vdots & & \vdots & \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} & b_{m'1} & \cdots & b_{m'n'} \end{pmatrix}. \quad (4)$$

By row catenation, we mean

$$X \ominus Y = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \\ b_{11} & \cdots & b_{1n'} \\ \vdots & & \vdots \\ b_{m'1} & \cdots & b_{m'n'} \end{pmatrix}. \quad (5)$$

If A and B are two partial arrays of equal size, then A is contained in B denoted by $A \subset B$ if $D(A) \subseteq D(B)$ and

$$A(i, j) = B(i, j) \quad \forall (i, j) \in D(A). \quad (6)$$

Definition 9. Partial arrays A and B are said to be compatible denoted by $A \uparrow B$, if there exists partial array C such that $A \subset C$ and $B \subset C$.

4. Compatibility and k -Compatibilty of Partial Arrays

4.1. Compatibility. The rules mentioned for partial words are also true for partial arrays.

Let A, B, X, Y be partial arrays.

- (i) *Multiplication*: If $A \uparrow B$ and $X \uparrow Y$, then $AX \uparrow BY$ either by column catenation or by row catenation.
- (ii) *Simplification*: If $AX \uparrow BY$ either by column catenation or by row catenation with A and B being of same size, then $A \uparrow B$ and $X \uparrow Y$.

(iii) *Weakening*: If $A \uparrow B$ and $C \subset A$, then $C \uparrow B$.

Lemma 3's version for partial arrays can be stated as follows.

Lemma 10. Let A, B, X, Y be partial arrays such that $AX \uparrow BY$, either by column catenation or by row catenation.

- (i) If order of $A \geq$ order of B , then there exist partial arrays C, Z such that $A = CZ$, $B \uparrow C$, and $Y \uparrow ZX$.
- (ii) If order of $A \leq$ order of B , then there exist partial arrays C, Z such that $B = CZ$, $A \uparrow C$, and $X \uparrow ZY$.

4.2. k -Compatibility

Definition 11. If A and B are two partial arrays of same size and k is nonnegative integer, then A is said to be k -contained in B denoted by $A \subset_k B$ if $D(A) \subset D(B)$ and there exists subset E of $D(A)$ of cardinality k called the error set such that

$$\begin{aligned} A(i, j) &= B(i, j) \quad \forall (i, j) \in D(A) \setminus E, \\ A(i, j) &\neq B(i, j) \quad \forall (i, j) \in E. \end{aligned} \quad (7)$$

Definition 12. If A and B are two partial arrays of same order and k is a nonnegative integer, then A and B are called k -compatible denoted by $A \uparrow_k B$ if there exist partial array Z and nonnegative integers k_1, k_2 such that

- (i) $A \subset_{k_1} Z$ with error set E_1 ;
- (ii) $B \subset_{k_2} Z$ with error set E_2 ;
- (iii) $E_1 \cap E_2 = \emptyset$;
- (iv) $k_1 + k_2 = k$.

Example 13. $A = \begin{pmatrix} b & a & b \\ a & \diamond & c \end{pmatrix}$, $B = \begin{pmatrix} a & b & a \\ a & \diamond & c \end{pmatrix}$, then there exists partial array $Z = \begin{pmatrix} a & b & b \\ a & \diamond & c \end{pmatrix}$ with $E_1 = \{(1, 1), (1, 2)\}$, $E_2 = \{(1, 3)\}$ and $k_1 = 2, k_2 = 1 \Rightarrow k = 3$; that is, $A \uparrow_3 B$.

Equivalently, A and B are k -compatible, if there exists subset E of $D(A) \cap D(B)$ of cardinality k called the error set such that

- (i) $A(i, j) = B(i, j) \forall (i, j) \in D(A) \cap D(B) \setminus E$;
- (ii) $A(i, j) \neq B(i, j) \forall (i, j) \in E$.

If A and B are arrays, then $A \uparrow_k B$ means $A = B$. We sometimes use notation $A \uparrow_{\leq k} B$, if set E has cardinality $\leq k$.

Multiplication. If $A \uparrow_{k_1} B$ and $X \uparrow_{k_2} Y$, then $AX \uparrow_{k_1+k_2} BY$ where A, B, X , and Y are partial arrays and k_1, k_2 are nonnegative integers, using column catenation.

Example 14. $A = \begin{pmatrix} \diamond & a & a \\ b & b & \diamond \\ a & b & a \end{pmatrix}$, $B = \begin{pmatrix} b & b & \diamond \\ a & a & \diamond \\ a & a & b \end{pmatrix}$, $X = \begin{pmatrix} b & \diamond & a \\ a & b & a \\ a & \diamond & b \end{pmatrix}$, $Y = \begin{pmatrix} a & b & b \\ b & a & b \\ \diamond & \diamond & a \end{pmatrix}$.
 $AX \uparrow_{6+7} BY$.

Simplification. If $AX \uparrow_k BY$ and order of A is equal to order of B , then $A \uparrow_{k_1} B$ and $X \uparrow_{k_2} Y$ for some k_1, k_2 , satisfying $k_1 + k_2 = k$.

Example 15. $A = \begin{pmatrix} \diamond & a & a \\ b & b & \diamond \\ a & b & a \end{pmatrix}$, $B = \begin{pmatrix} b & b & \diamond \\ a & \diamond & a \\ b & a & b \end{pmatrix}$, $X = \begin{pmatrix} b & \diamond \\ a & b \\ a & \diamond \end{pmatrix}$, $Y = \begin{pmatrix} a \\ b \\ b \end{pmatrix}$.
 $AX \uparrow_8 BY \Rightarrow A \uparrow_5 B$ and $X \uparrow_3 Y$ with $5 + 3 = 8$.

Weakening. If $A \uparrow_k B$ and $Z \subset A$, then $Z \uparrow_{\leq k} B$.

Example 16. $A = \begin{pmatrix} \diamond & a & a \\ b & b & \diamond \\ a & b & a \end{pmatrix}$, $B = \begin{pmatrix} b & b & \diamond \\ a & \diamond & a \\ b & a & b \end{pmatrix}$, $Z = \begin{pmatrix} a & a \\ b & \diamond \\ b & a \end{pmatrix}$.
 $Z \uparrow_{\leq 7} B$ with $k = 7$.

Theorem 17. Let A and B be partial arrays of orders $a \times b$ and $a \times c$, respectively. If there exist array Z of order $a \times d$ and integers k_1, k_2, m , and n such that $A \subset_{k_1} Z^m$ with error set E_1 and $B \subset_{k_2} Z^n$ with error set E_2 , then there exist integers p and q such that $A \uparrow_{\leq k} B^q$ with

$$\begin{aligned} k &= \|(D(A)(a, |b|, p) \cap E_2(a, |c|, q)) \\ &\cup (D(B)(a, |c|, q) \cap E_1(a, |b|, p))\|. \end{aligned} \quad (8)$$

Moreover, if $E_1(a, |b|, n) \cap E_2(a, |c|, m) = \emptyset$, then $A \uparrow_k B^q$.

Proof. Let A and B be partial arrays of $a \times b$ and $a \times c$, respectively. Let array z of order $a \times d$ exist such that, by using column catenation, $A \subset_{k_1} Z^m$ and $B \subset_{k_2} Z^n$ for some integers k_1, k_2, m , and n . Let E_1 be the error set of cardinality k_1 such that $A(i, j) = Z^m(i, j)$ for all $(i, j) \in D(A) \setminus E_1$ and $A(i, j) \neq Z^m(i, j)$ for all $(i, j) \in E_1$ and E_2 be the error set of cardinality k_2 such that $B(i, j) = Z^n(i, j)$ for all $(i, j) \in D(B) \setminus E_2$ and $B(i, j) \neq Z^n(i, j)$ for all $(i, j) \in E_2$. We have $A \subset_{nk_1} Z^{mn}$ with error set $E_1(a, |b|, n)$ of cardinality nk_1 and $B \subset_{mk_2} Z^{mn}$ with error set $E_2(a, |c|, m)$ of cardinality mk_2 .

Let $(1, 1) \leq (i, j) \leq (a, d^{mn})$ and $Z^{mn}(i, j) = a$ for some letter a . There are 4 possibilities.

Case 1. If $(i, j) \notin E_1(a, |b|, n)$ and $(i, j) \notin E_2(a, |c|, m)$, then $A^n(i, j) \in \{\diamond, a\}$ and $B^m(i, j) \in \{\diamond, a\}$. It does not give any error, when we align A^n with B^m .

Case 2. If $(i, j) \notin E_1(a, |b|, n)$ and $(i, j) \in E_2(a, |c|, m)$, then $A^n(i, j) \in \{\diamond, a\}$ and $B^m(i, j) = b$ for some $b \neq a$. It gives an error in the alignment of A^n with B^m only when $A^n(i, j) = a$ or when $(i, j) \in D(A)(a, |b|, n)$.

Case 3. If $(i, j) \in E_1(a, |b|, n)$ and $(i, j) \in E_2(a, |c|, m)$, then $B^m(i, j) \in \{\diamond, a\}$ and $A^n(i, j) = b$ for some $b \neq a$. It gives an error in the alignment of A^n with B^m only when $B^m(i, j) = a$ or when $(i, j) \in D(B)(a, |c|, m)$.

Case 4. If $(i, j) \in E_1(a, |b|, n)$ and $(i, j) \in E_2(a, |c|, m)$, then $A^n(i, j) = b$ for some $b \neq a$ and $B^m(i, j) = c$ for some $c \neq a$. It gives an error in the alignment of A^n with B^m only when $b \neq c$.

Therefore, if $E_1(a, |b|, n) \cap E_2(a, |c|, m) = \emptyset$ then $A^n \uparrow_k B^m$ with $k = \|(D(A)(a, |b|, n) \cap E_2(a, |c|, m)) \cup (D(B)(a, |c|, m) \cap E_1(a, |b|, n))\|$ and $E_1(a, |b|, n) \cap E_2(a, |c|, m) \neq \emptyset$ then $A^n \uparrow_{\leq k} B^m$. $\square$

Example 18. $A = \begin{pmatrix} a & b & \diamond \\ c & a & b \\ \diamond & b & a \end{pmatrix}$, $B = \begin{pmatrix} a & c \\ c & b \\ \diamond & b \end{pmatrix}$, $Z = \begin{pmatrix} a \\ c \\ b \end{pmatrix}$.

We have AC_4Z^3 with error set $E_1 = \{(1, 2), (2, 2), (2, 3), (3, 3)\}$, and BC_2Z^2 with error set $E_2 = \{(1, 2), (2, 2)\}$.

$k = 6$:

$$(i) D(A) = \{(1, 1), (1, 2), (2, 1), (2, 2), (2, 3), (3, 2), (3, 3)\}.$$

$$D(B) = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 2)\}$$

$$(ii) D(A)(a, |b|, 2) = \{(1, 1), (1, 2), (2, 1), (2, 2), (2, 3), (3, 2), (3, 3), (1, 4), (1, 5), (2, 4), (2, 5), (2, 6), (3, 5), (3, 6)\}.$$

$$D(B)(a, |c|, 3) = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4), (1, 5), (1, 6), (2, 5), (2, 6), (3, 6)\}$$

$$(iii) E_1(a, |b|, 2) = \{(1, 2), (2, 2), (2, 3), (3, 3), (1, 5), (2, 5), (2, 6), (3, 6)\}.$$

$$E_2(a, |c|, 3) = \{(1, 2), (2, 2), (1, 4), (2, 4), (1, 6), (2, 6)\}.$$

$$E_1(a, |b|, 2) \cap E_2(a, |c|, 3) \neq \phi.$$

$$k = \|(D(A)(a, |b|, 2) \cap E_2(a, |c|, 3)) \cup (D(B)(a, |c|, 3) \cap E_1(a, |b|, 2))\| = \|((1, 1), (1, 2), (2, 1), (2, 2), (2, 3), (3, 2), (3, 3), (1, 4), (1, 5), (2, 4), (2, 5), (2, 6), (3, 5), (3, 6)) \cap ((1, 2), (2, 2), (1, 4), (2, 4), (1, 6), (2, 6)) \cup ((1, 1), (1, 2), (2, 1), (2, 2), (3, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4), (1, 5), (1, 6), (2, 5), (2, 6), (3, 6)) \cap ((1, 2), (2, 2), (2, 3), (3, 3), (1, 5), (2, 5), (2, 6), (3, 6)))\| = \|(1, 2), (1, 4), (1, 5), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6), (3, 6))\|.$$

$k = 9$:

$$A^2 \uparrow_{\leq 9} B^3 (A^2 \uparrow_6 B^3).$$

5. Conjugacy and k -Conjugacy of Partial Arrays

5.1. Conjugacy

Definition 19. Two partial arrays A and B of same order are called conjugate if there exist partial arrays X and Y such that $A \subset XY$ and $B \subset YX$ using row catenation or column catenation.

0-conjugacy on partial arrays with same order is trivially reflexive and symmetric but not transitive.

$$\text{Example 20. } A = \begin{pmatrix} a & \diamond & b \\ b & c & a \\ a & a & \diamond \end{pmatrix}, B = \begin{pmatrix} b & c & \diamond \\ a & a & \diamond \\ a & c & b \end{pmatrix}, C = \begin{pmatrix} b & a & \diamond \\ b & a & \diamond \\ a & b & c \end{pmatrix}.$$

By taking $X = \begin{pmatrix} a & c & b \end{pmatrix}$ and $Y = \begin{pmatrix} b & c & a \\ a & a & \diamond \end{pmatrix}$, we get $A \subset XY$ and $B \subset YX$ showing that A and B are conjugate similarly and, by taking $X' = \begin{pmatrix} b & c & \diamond \end{pmatrix}$ and $Y' = \begin{pmatrix} a & c & b \end{pmatrix}$, we get $B \subset X'Y'$ and $C \subset Y'X'$ showing that B and C are conjugate. But A and C are not conjugate.

That is, conjugate relation is not transitive.

Proposition 21. Let A and B be nonempty partial arrays of same size. If A and B are conjugate, then there exists partial array C such that $AC \uparrow CB$, either by column catenation or by row catenation.

Proof. Let A and B be nonempty partial arrays of same order. Suppose A and B are conjugate and let X, Y be partial arrays such that $A \subset XY$ and $B \subset YX$ either by column catenation or by row catenation; then $AX \subset XYX$ and $XB \subset XYX$. So, for $C = X$, we have $AC \uparrow CB$. $\square$

5.2. k -Conjugacy

Definition 22. Two partial arrays A and B of same order are k -conjugate, if there exist nonnegative integers k_1, k_2 whose sum is k and partial arrays X and Y such that $AC_{k_1}XY$ and $BC_{k_2}YX$ with row or column catenation.

Theorem 23. Let A and B be nonempty partial arrays of same order. If A and B are k -conjugate, then there exists partial array Z such that $AZ \uparrow_{\leq k} ZB$.

Proof. Let A, B be two partial arrays of same order. Supposing that A and B are k -conjugate, then, by definition, there exist nonnegative integers k_1, k_2 whose sum is k and partial arrays X and Y such that $AC_{k_1}XY$ with error set E_1 and $BC_{k_2}YX$ with error set E_2 using row catenation or column catenation accordingly.

Then, $AXC_{k_1}XYX$ with error set E_1 and $XBC_{k_2}XYX$ with error set $E'_2 = \{(i + \text{number of rows of } X, j)/(i, j) \in E_2\}$ or $E'_2 = \{(i, j + \text{number of columns of } X)/(i, j) \in E_2\}$ according to row or column catenation and so, for $Z = X$, we have $AZ \uparrow_{\leq k} ZB$. $\square$

$$\text{Example 24. Given } A = \begin{pmatrix} a & \diamond & b \\ b & c & a \\ a & a & \diamond \end{pmatrix}, B = \begin{pmatrix} b & c & b \\ a & a & \diamond \\ a & a & b \end{pmatrix}.$$

There exist $X = \begin{pmatrix} a & \diamond & b \end{pmatrix}$ and $Y = \begin{pmatrix} c & a & b \\ a & a & \diamond \end{pmatrix}$ with AC_3XY and BC_2YX , $k = k_1 + k_2 = 5$.

There exist $Z = \begin{pmatrix} a & \diamond & b \end{pmatrix}$ such that $AZ \uparrow_{\leq 5} ZB$.

6. Conclusion

Motivated by compatibility and conjugacy properties of partial words, we define the conjugacy of partial array and derive the compatibility properties of partial arrays. By giving relaxation to the compatibility relation, we discuss k -compatibility and k -conjugacy of partial arrays. We prove that, given partial arrays A, B and integers p, q satisfying $|A|^p = |B|^q$, we find k such that $A^p \uparrow_k B^q$. Also, there exists partial array Z such that $AZ \uparrow_{\leq k} ZB$.

Disclosure

S. Vijayachitra is a Research Scholar at Department of Science and Humanity Sathyabama University, Chennai, India.

Competing Interests

The authors declare that they have no competing interests.

References

- [1] D. G. Arquès and Ch. J. Michel, “A possible code in the genetic code,” in *STACS 95*, vol. 900 of *Lecture Notes in Computer Science*, pp. 640–651, Springer, Berlin, Germany, 1995.
- [2] F. Blanchet-Sadri and R. A. Hegstrom, “Partial words and a theorem of Fine and Wilf revisited,” *Theoretical Computer Science*, vol. 270, no. 1-2, pp. 401–419, 2002.
- [3] A. Colosimo and A. de Luca, “Special factors in biological strings,” *Journal of Theoretical Biology*, vol. 204, no. 1, pp. 29–46, 2000.
- [4] F. Blanchet-Sadri, D. Dakota Blair, and R. V. Lewis, “Equations on partial words,” *RAIRO—Theoretical Informatics and Applications*, vol. 43, pp. 23–29, 2009.
- [5] F. Sweetey, D. G. Thomas, and T. Kalyani, “Collage of hexagonal arrays,” in *Advances in Visual Computing*, G. Bebis, R. Boyle, B. Parvin et al., Eds., vol. 5359 of *Lecture Notes in Computer Science*, pp. 1167–1175, Springer, Berlin, Germany, 2008.
- [6] F. Blanchet-Sadri and D. K. Luhmann, “Conjugacy on partial words,” *Theoretical Computer Science*, vol. 289, no. 1, pp. 297–312, 2002.
- [7] J. Berstel and L. Boasson, “Partial words and a theorem of Fine and Wilf,” *Theoretical Computer Science*, vol. 218, no. 1, pp. 135–141, 1999.
- [8] A. de Luca, “On the combinatorics of finite words,” *Theoretical Computer Science*, vol. 218, no. 1, pp. 13–39, 1999.
- [9] C. Choffrut and J. Karhumäki, “Combinatorics of words,” in *Handbook of Formal Languages*, G. Rozenberg and A. Salomaa, Eds., vol. 1, chapter 6, pp. 329–438, Springer, Berlin, Germany, 1999.
- [10] M. Lothaire, *Combinatorics on Words*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, UK, 1997.
- [11] F. Sweetey, D. G. Thomas, and V. R. Dare, “Subarray complexity of partial arrays,” in *Intelligent Optimization Modeling*, pp. 85–94, Allied, 2006.