

## Research Article

# On the Observability of Leader-Based Multiagent Systems with Fixed Topology

Bo Liu <sup>1,2</sup>, Ningsheng Xu,<sup>2</sup> Housheng Su <sup>3</sup>, Licheng Wu,<sup>1</sup> and Jiahui Bai<sup>2</sup>

<sup>1</sup>School of Information Engineering, Minzu University of China, Beijing 100081, China

<sup>2</sup>College of Science, North China University of Technology, Beijing 100144, China

<sup>3</sup>School of Artificial Intelligence and Automation, Huazhong University of Science and Technology, Wuhan 430074, China

Correspondence should be addressed to Housheng Su; [houshengsu@qq.com](mailto:houshengsu@qq.com)

Received 18 June 2019; Revised 7 September 2019; Accepted 16 September 2019; Published 5 November 2019

Academic Editor: Xianggui Guo

Copyright © 2019 Bo Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper investigates the observability of first-order, second-order, and high-order leader-based multiagent systems (MASs) with fixed topology, respectively. Some new algebraic and graphical characterizations of the observability for the first-order MASs are established based on agreement protocols. Moreover, under the same leader-following framework with the predefined topology and leader assignment, the observability conditions for systems of double-integrator and high-integrator agents are also obtained. Finally, the effectiveness of the theoretical results is verified by numerical examples and simulations.

## 1. Introduction

Self-organizing behavior of multiple intelligent dynamic agents is widespread in nature, for example, bird flocking flight [1], fish group patrol [2], microbial collective foraging [3], and ant colony cooperation [4]. This self-organizing behavior of such intelligent and dynamical networked system, through transferring and interaction of information between local agents, as a whole, presenting the ability of effective coordination and the level of advanced intelligence, has led to the wide concern and interest in many fields including applied mathematics, ecology, biology, sociology, physics, computer science, control, and engineering. At present, coordinated control of intelligent systems has become a hot research topic in the field of control and is widely used in smart grids, UAV formation, intelligent transportation systems, artificial satellite networks, and so on [5–20].

Observability is a significant problem and becomes the basis of the design of optimal control and optimal estimation in modern control theory, which was put forward by Kalman in 1960. Intuitively speaking, the so-called system's observability means that the internal states of the system can be correctly known by observing the external variables of the system. In general, for a multiagent network, the

observability is the key problem and means that the entire system's state can be completely reconstructed by only observing a small quantity of intelligent dynamic agents (i.e., leaders). Observability plays a fundamental role in state estimation and system control as well as has recently been widely applied in wireless sensor networks, satellite navigation systems, autonomous underwater vehicles, machine games, and so on.

The concepts of controllability and observability were first tried to apply to networked multiagent systems by Tanner in 2004 [21], where all agents are partitioned into leader agents and follower agents, in which the follower agents are only affected by the neighbor interaction protocol and the leader agents are influenced by the external control input. This structure, called leader-follower structure, can make the whole system as a standard linear time-invariant system so that the classical controllability and observability are all applied into networked multiagent systems. Observability is very important to study distributed estimation and intrusion detection problems in distributed sensor networks [22]. Zelazo and Mesbahi [23] investigated the observability of networked systems with homogeneous and heterogeneous dynamics, respectively, as well as provided an analysis for the observability of networked linear systems. In

[24], the study mainly focused on controllability and observability of multiagent networks under a leader-follower framework with the chain and cyclic topologies, respectively. In [25], the authors established a decentralized condition for controllability and observability of networked MASs using decentralized Laplacian spectrum estimation. The observability of a network system was investigated based on average consensus algorithm for path and cycle graphs in [26], graph Cartesian products in [27], regular graphs and distance-regular graphs [28, 29], and equitable partition [30], respectively. Furthermore, the observability of switched linear systems and switched multiagent systems was studied in [31, 32], respectively. Recently, the researchers [33–35] investigated the observability of MASs with time-varying and switching topologies, respectively. In addition, Liu et al. [36] analyzed the observability for complex networks.

It is worthwhile to note that most of the previous research conclusions mainly focused on controllability and observability of first-order networked MASs. However, it is well known that, in control engineering, almost all control systems are high-order systems, that is, higher-order differential/difference equations are used to describe those control systems studied in many areas, such as consensus, controllability, flocking, containment control, and stabilizability. In this paper, the observability of a class of MASs is considered, and a unified framework for the inertia of first-order, second-order, and high-order MASs is given. Most of the existing literatures on the observability of MASs focused only on a single dynamic equation, such as a first-order dynamics, a second-order dynamics, or a high-order dynamics, rather than a unified consideration. In general, it is difficult to bring the three kinds of networks into a unified framework, which is a hard point in modelling and analyzing. This paper aims at the observability of MASs with first-order/second-order/high-order agents based on agreement protocols, respectively. The main contributions of this paper are summarized as follows: (i) Some necessary and sufficient conditions for the observability of first-order MASs are built by Laplacian matrix, which relies only on the information flows within the agents and between the agents from the leaders. (ii) It is proved that the observability of the second-order/high-order MASs is equivalent to that of first-order MASs with the same information of topology structure and the same prescribed leaders. (iii) A unified framework for the observability inertia of first-order, second-order, and high-order MASs is established, which relies only on the information topology of such network.

## 2. Mathematical Preliminaries

For the convenience of discussing the observability of networked systems, the information communication links among agents can be described by a weighted undirected/directed graph denoted as  $\mathcal{G} = (\mathcal{V}, E, \mathcal{A})$ , in which  $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$  is a *vertex set* and  $E = \{(v_i, v_j) : v_i, v_j \in \mathcal{V}\}$  is an *edge set*, respectively. The weighted adjacency matrix of  $\mathcal{G}$  is marked as  $\mathcal{A} = [a_{ij}] \in \mathbb{R}^{n \times n}$  with  $a_{ij} \neq 0$  if and only if  $(v_i, v_j) \in \mathcal{E}$  and  $a_{ij} = 0$ , otherwise. Moreover,  $\mathcal{N}_i = \{v_j \in \mathcal{V} : (v_i, v_j) \in E\}$  represents the neighbors' set of agent  $v_i$ . The

Laplacian matrix is given as  $L \triangleq \Delta - \mathcal{A}$ , where  $\mathcal{A}$  is the weighted adjacency matrix and  $\Delta$  is the diagonal degree matrix.

For a network with leader-follower structure (see Figure 1), the Laplacian matrix  $L$  can be divided into the following block matrices:

$$L = \begin{bmatrix} L_f & L_{fl} \\ L_{lf} & L_l \end{bmatrix}, \quad (1)$$

where  $L_f$  and  $L_l$  stand for the communication information links within the followers and the leaders, respectively,  $L_{fl}$  stands for the communication information links from the leaders to the followers, and  $L_{lf}$  stands for the communication information links from the followers to the leaders, where  $L_{fl}^T = L_{lf}$  if graph  $\mathcal{G}$  is undirected.

## 3. Observability Analysis

**3.1. Observability of First-Order MASs.** A first-order MAS consists of  $(n + l)$  agents, in which the former  $n$  agents are regarded as followers and the remaining  $l$  agents are regarded as leaders, and its dynamic equation is described by

$$\begin{aligned} \dot{x}_i &= u_i, \\ i &\in \mathcal{L}_{n+l}, \end{aligned} \quad (2)$$

where  $x_i \in \mathbb{R}^n$  represents the state and  $u_i \in \mathbb{R}^q$  is the external input. Interactions among agents of system (2) in this paper are given by the agreement protocol as follows:

$$u_i = \sum_{j \in \mathcal{N}_i} a_{ij}(x_j - x_i), \quad i \in \mathcal{L}_{n+l}, \quad (3)$$

where  $\mathcal{N}_i$  is the neighbor set of agent  $i$  and  $a_{ij} \geq 0$  for all  $i, j$ .

Let  $x \triangleq [x_1, \dots, x_{n+l}]^T$ ,  $x_f \triangleq [x_1, \dots, x_n]^T$ , and  $x_l \triangleq [x_{n+1}, \dots, x_{n+l}]^T$ . From the partition of  $L$ , the dynamic equations (2) and (3) can be redescribed by

$$\dot{x} = \begin{bmatrix} \dot{x}_f \\ \dot{x}_l \end{bmatrix} = -Lx = -\begin{bmatrix} L_f & L_{fl} \\ L_{lf}^T & L_l \end{bmatrix} \begin{bmatrix} x_f \\ x_l \end{bmatrix}. \quad (4)$$

In the actual control process, because of the limitation of the measuring devices, the state of the system is difficult to be measured directly, but can be determined by observing the output of the system. The observability of MASs refers to the observation of the state of the whole MAS only from a few agents as much as possible. Note that it is congenial with reason that some leaders can measure their neighbors' states and exchange information flows between the other leaders, so we can let  $y$  be the output vector measured by the leaders. Supposed the leaders can directly exchange information with the other agents, there is a complete communication graph between the leaders. In other words, each leader can broadcast its measured output to all other leaders and has access to the entire output vector  $y$ . Therefore, each leader here can get all the output vectors. According to the partition of Laplace matrix,  $y = L_{fl}^T x_f$  is chosen as the output. Therefore, the dynamic equation of the networked MAS can be looked upon as a classical linear time-invariant (LTI) system as follows:




FIGURE 1: A leader-follower framework.

$$\begin{cases} \dot{x}_f = -L_f x_f - L_{fl} x_l, \\ y = -L_{fl}^T x_f, \end{cases} \quad (5)$$

whose observability is equivalent to that of the following system:

$$\begin{cases} \dot{x}_f = L_f x_f + L_{fl} x_l, \\ y = L_{fl}^T x_f, \end{cases} \quad (6)$$

where  $y \in \mathbb{R}^n$  represents the output, whose corresponding output matrix is denoted as  $L_{fl}^T$ , and contains all the state vectors measured by the leaders. Therefore, the classical concept of observability of a standard LTI system can be used to solve the problem of the observable state variables that are measurable by the leaders.

Observability is a fundamental problem of networked MASs, which focuses on whether the whole system state can be reconstructed by the output-observed agents' evolution behaviors. Then, observability concerns on the identical zero output brought by the nonzero initial states. In other words, such multiagent network is observable if there exists no unobservable state. For the convenience of theoretical demonstration and calculation, we only need to study the observability of system (6) in the following.

**Definition 1** (see [35]). Given  $x(t_0) = x_0$ , a state  $x_0$  of system (6) is unobservable if  $y(t) = 0$  for any  $t \in [t_0, t_f]$ , where finite time  $t_f > t_0$ .

According to known results in linear time-invariant system theory, we can derive the following proposition.

**Proposition 1.** For system (6), the following statements are equivalent:

- (1) System (6) is observable.
- (2) Observability matrix  $Q_o = \begin{bmatrix} L_{fl}^T \\ L_{fl}^T L_f \\ \vdots \\ L_{fl}^T (L_f)^{n-1} \end{bmatrix}$  has full rank.
- (3) Matrix  $\begin{bmatrix} \lambda I - L_f \\ L_{fl}^T \end{bmatrix}$  has full rank for  $\forall \lambda \in \mathbb{R}$ .
- (4) Matrix  $L_f$  has no nonzero right eigenvectors orthogonal to all rows of  $L_{fl}^T$ . Namely, for all eigenvalues  $(\lambda_i, i = 1, \dots, n)$  of matrix  $L_f$ , if

$$\begin{aligned} L_f \alpha &= \lambda \alpha, \\ L_{fl}^T \alpha &= 0, \end{aligned} \quad (7)$$

both hold, then the right eigenvector  $\alpha = 0$ .

Together with some fundamental propositions of the graph Laplacian for an undirected graph, the following results can be obtained.

**Theorem 1.** The first-order MAS (6) is observable iff  $L$  and  $L_f$  do not share common eigenvalues.

*Proof.* Similar proof can be seen from Lemma 2.2 in reference [37], omitted here.  $\square$

**Remark 1.** It is noted from Theorem 1 that the observability of networked MASs relies only on the information links between agents. Theorem 1 provides a simpler and more easily checkable method because it only involves the couplings of the entire network (between all agents) and sub-network (only among follower agents). It is also easy to compute the eigenvalues of the coupling matrices of the entire network and the subnetwork by MATLAB. In addition, Theorem 1 is the theoretical basis and analytic technique on observability.

**Theorem 2.** The first-order MAS (6) is observable iff there exists no eigenvector of  $L$  taking zero on the element corresponding to the leader agent.

*Proof (necessity).* By contradiction, supposed  $\lambda$  is the eigenvalue of  $L$ , whose corresponding eigenvector of  $L$  is denoted as  $\xi \triangleq \begin{bmatrix} \xi_f \\ \xi_l \end{bmatrix}$ , in which the elements corresponding to the leader agent are zero, that is,  $\xi_l = 0$ , then we can have

$$L\xi = \begin{bmatrix} L_f & L_{fl} \\ L_{fl}^T & L_l \end{bmatrix} \begin{bmatrix} \xi_f \\ 0 \end{bmatrix} = \begin{bmatrix} L_f \xi_f \\ L_{fl}^T \xi_f \end{bmatrix} = \lambda \begin{bmatrix} \xi_f \\ 0 \end{bmatrix}, \quad (8)$$

and then,  $L_f \xi_f = \lambda \xi_f$  and  $L_{fl}^T \xi_f = 0$ . Therefore, system (6) is unobservable from Proposition 1 (6), which is in contradiction with the fact that system (6) is observable. Then, the proof of the necessity is completed. *Sufficiency.* By contradiction, assume that system (6) is unobservable; from Theorem 1, we can know that  $L$  and  $L_f$  share one common eigenvalue at least, say  $\lambda$ , which corresponds to the eigenvector  $\xi$  satisfying

$$\xi = M \xi_f = \begin{bmatrix} \xi_f \\ 0 \end{bmatrix}, \quad (9)$$

where  $M \triangleq [I_f \ 0]^T$  and  $\xi$  and  $\xi_f$  are the eigenvectors of  $L$  and  $L_f$  corresponding to the common eigenvalue  $\lambda$ , respectively. Moreover, we can know from equation (4) that

$$L\xi = \begin{bmatrix} L_f & L_{fl} \\ L_{fl}^T & L_l \end{bmatrix} \begin{bmatrix} \xi_f \\ 0 \end{bmatrix} = \lambda \begin{bmatrix} \xi_f \\ 0 \end{bmatrix}, \quad (10)$$

which contradicts to the fact there exists no eigenvector of  $L$  taking zero on the element corresponding to the leader agent. Thus, the proof of the sufficiency is completed.  $\square$

**Theorem 3.** *The first-order MAS (6) is observable if*

- (i) *The eigenvalues of  $L_f$  are all distinct;*
- (ii) *All the row vectors of  $L_{f_l}^T$  are not orthogonal to at least one eigenvector of  $L_f$  simultaneously.*

*Proof.*  $L_f$  can be shown as  $L_f = UDU^{-1}$  if the eigenvalues of  $L_f$  are all distinct, where  $D \triangleq \text{diag}(\lambda_1, \dots, \lambda_n)$  with  $\lambda_i$  being the eigenvalues of  $L_f$  for  $i = 1, \dots, n$ , as well as  $U$  consists of the eigenvectors of  $L_f$ . The observability matrix  $Q_o$  can be presented as

$$\begin{aligned} Q_o &= \begin{bmatrix} L_{f_l}^T \\ L_{f_l}^T L_f \\ \vdots \\ L_{f_l}^T (L_f)^{n-1} \end{bmatrix} = \begin{bmatrix} L_{f_l}^T U U^{-1} \\ L_{f_l}^T U D U^{-1} \\ \vdots \\ L_{f_l}^T (U D U^{-1})^{n-1} \end{bmatrix} \\ &= \begin{bmatrix} L_{f_l}^T U U^{-1} \\ L_{f_l}^T U D U^{-1} \\ \vdots \\ L_{f_l}^T U D^{n-1} U^{-1} \end{bmatrix} = \begin{bmatrix} L_{f_l}^T U \\ L_{f_l}^T U D \\ \vdots \\ L_{f_l}^T U D^{n-1} \end{bmatrix} U^{-1}. \end{aligned} \quad (11)$$

$\square$

Because  $U^{-1}$  is nonsingular, we only consider the matrix

$$\tilde{Q}_o^T = [L_{f_l}^T U : L_{f_l}^T U D : \dots : L_{f_l}^T U D^{n-1}], \quad (12)$$

whose rank is completely determined by matrix  $L_{f_l}^T U$  since the diagonal matrix  $D$  is nonsingular. If all the row vectors of  $L_{f_l}^T$  are not orthogonal to at least one eigenvector of  $L_f$  simultaneously, then all the elements of at least one column of  $L_{f_l}^T U$  are nonzero so that observability matrix  $Q_o$  has full rank. Therefore, the first-order multiagent system (6) is observable.

From Theorem 3, we have obtained the following result immediately.

**Theorem 4.** *The first-order MAS (6) is observable if*

- (i) *The eigenvalues of  $L_f$  are all distinct;*
- (ii) *Every column of  $L_{f_l}^T U$  has at least one nonzero element, where  $U$  consists of the eigenvectors of  $L_f$ .*

*Proof.* From condition (i) and Theorem 3, we can have

$$\text{rank} \begin{bmatrix} \lambda_i I_n - D \\ L_{f_l}^T U \end{bmatrix} = \text{rank} \begin{bmatrix} \lambda_i - \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_i - \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_i - \lambda_n \\ b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{21} & \dots & b_{1n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{l1} & b_{l2} & \dots & b_{ln} \end{bmatrix}, \quad (13)$$

where  $D = \text{diag}(\lambda_1, \lambda_1, \dots, \lambda_n)$  with  $\lambda_i$  ( $i = 1, 2, \dots, n$ ) being the eigenvalues of  $L_f$  and  $L_{f_l}^T U = [b_{ij}] \in \mathbb{R}^{l \times n}$ . Since  $\lambda_i \neq \lambda_j$  for  $i \neq j$ , and every column of  $L_{f_l}^T U$  has at least one nonzero element, then

$$\text{rank} \begin{bmatrix} \lambda_i I_n - D \\ L_{f_l}^T U \end{bmatrix} = n. \quad (14)$$

According to Proposition 1 (3), this conclusion holds.  $\square$

*Remark 2.* It is noted from Proposition 1 (2) the observability of first-order MASs relies only on the information links between agents, that is,  $(L_f, L_{f_l}^T)$ . Theorem 1 mainly illustrates the influence of the relationship between the entire network among all agents (from leader agents to follower agents and among follower agents) and the sub-network among follower agents on the observability of the system. Theorem 2 shows how to choose the leaders to make the whole system observable. Theorem 3 and Theorem 4 explain the topological relations in Theorem 1 in more detail. Therefore, Theorem 1 is the theoretical basis and analytic technique on observability of Theorems 2–4.

In fact, the observability of first-order MASs cannot be directly extended to the second-order or higher-order MASs. It is of physical and theoretical significance to study the observability for MASs of second-order integrator or high-order integrator agents since first-order system is too simple to depict the dynamic characteristics accurately in the actual physical systems. Therefore, the model with second-order integrator or high-order integrator agents is beneficial for the research of cooperation in real world. However, for the observability of MASs with different orders, the observability problem brings new characteristics and research difficulties, involving in how to model, design the control protocols, select the number of leaders, and locate the leaders, as well as define the observability and so on. Specially, the leaders' positions and speeds have important impacts on observability. These factors make the observability of MASs of second-order integrator or high-order integrator agents a nontrivial new problem. Due to lack of research tools, there are very few results for the observability of MASs with second-order or high-order being available in the existing literatures. Therefore, the results for the first-order case cannot be directly extended to the second-order or high-order MASs.

**3.2. Observability of Second-Order MASs.** A second-order MAS consists of  $(n + l)$  agents, in which the former  $n$  agents are regarded as followers and the remaining  $l$  agents are regarded as leaders, and its dynamic equation is depicted by

$$\begin{aligned} \dot{x}_i &= v_i, \\ \dot{v}_i &= u_i, \\ i &\in \mathcal{L}_{n+l}, \end{aligned} \quad (15)$$

with

$$u_i = c_1 \sum_{j \in \mathcal{N}_i} a_{ij} (x_j - x_i) + c_2 v_i, \quad (16)$$

where  $x_i \in \mathbb{R}^n$  is the position state and  $v_i \in \mathbb{R}^n$  is the velocity state of agent  $i$ , respectively;  $u_i \in \mathbb{R}^q$  is the input;  $c_1$  and  $c_2$  are nonzero feedback gains.

Let  $x_f \triangleq [x_1, \dots, x_n]^T$ ,  $x_l \triangleq [x_{n+1}, \dots, x_{n+l}]^T$ ,  $v_f \triangleq [v_1, \dots, v_n]^T$ , and  $v_l \triangleq [v_{n+1}, \dots, v_{n+l}]^T$ . Based on the partition of follower agents and leader agents, the dynamics (15) and (16) can be rewritten as follows:

$$\begin{bmatrix} \dot{x}_f \\ \dot{x}_l \\ \dot{v}_f \\ \dot{v}_l \end{bmatrix} = \begin{bmatrix} 0 & 0 & I_n & 0 \\ 0 & 0 & 0 & I_n \\ c_1 L_f & c_1 L_{fl} & c_2 I_n & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_f \\ x_l \\ v_f \\ v_l \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \\ u_{\text{ext}} \end{bmatrix}, \quad (17)$$

where  $u_{\text{ext}} = [u_{\text{ext}}^1, \dots, u_{\text{ext}}^l]^T \in \mathbb{R}^l$  is the external control input. Specially, the dynamic equation (17) of followers can be described as

$$\begin{cases} \dot{\bar{x}}_f = \begin{bmatrix} \dot{x}_f \\ \dot{v}_f \end{bmatrix} = \begin{bmatrix} 0 & I_n \\ c_1 L_f & c_2 I_n \end{bmatrix} \begin{bmatrix} x_f \\ v_f \end{bmatrix} + \begin{bmatrix} 0 \\ c_1 L_{fl} \end{bmatrix} x_l \triangleq \bar{L}_f \bar{x}_f + \bar{L}_{fl} x_l, \\ y = \bar{L}_{fl}^T \begin{bmatrix} x_f \\ v_f \end{bmatrix} \triangleq \bar{L}_{fl}^T \bar{x}_f, \end{cases} \quad (18)$$

where  $\bar{x}_f \triangleq [x_f, v_f]^T$ ,  $\bar{L}_f \triangleq \begin{bmatrix} 0 & I_n \\ c_1 L_f & c_2 I_n \end{bmatrix}$ , and  $\bar{L}_{fl} \triangleq \begin{bmatrix} 0 \\ c_1 L_{fl} \end{bmatrix}$ .

For the observability of second-order multiagent systems, we can obtain the following results.

**Theorem 5.** *The second-order MAS (18) is observable iff  $(L_f, L_{fl}^T)$  is observable.*

*Proof.* Supposed  $\lambda$  is the eigenvalue of matrix  $\bar{L}_f$ , which corresponds to the right eigenvector of  $\bar{L}_f$  denoted as  $\beta = [\beta_1 \beta_2]^T$ , then we can have  $\bar{L}_f [\beta_1 \beta_2]^T = \lambda [\beta_1 \beta_2]^T$ , and then

$$\begin{cases} \lambda \beta_1 = \beta_2, \\ \lambda \beta_2 = c_1 L_f \beta_1 + c_2 \beta_2. \end{cases} \quad (19) \quad \square$$

Consequently,  $L_f \beta_1 = (\lambda^2 - \lambda c_2 / c_1) \triangleq u \beta_1$  (since  $c_1 \neq 0$ ), where  $u$  is the eigenvalue of  $L_f$  with the corresponding right eigenvector  $\beta_1$ , which is the root of equation  $s^2 - c_2 s - c_1 u = 0$  with  $\beta_2 = s \beta_1$ . In the following, we will show the fact that  $(\bar{L}_f, \bar{L}_{fl}^T)$  is observable iff  $(L_f, L_{fl}^T)$  is observable.

**3.2.1. Sufficiency.** Supposed  $(L_f, L_{fl}^T)$  is unobservable, then there must exist an eigenvalue  $u$  of  $L_f$ , whose corresponding right eigenvector is  $\beta_1 \neq 0$ , such that  $L_f \beta_1 = u \beta_1$  and  $L_{fl}^T \beta_1 = 0$ . Therefore, we can assert that the eigenvalue of  $\bar{L}_f$  satisfies equation  $s^2 - c_2 s - c_1 u = 0$ , and the corresponding right eigenvector of  $\bar{L}_f$  is  $[\beta_1 \beta_2]^T$  with  $\beta_2 = s \beta_1$ . Then,  $\bar{L}_{fl}^T [\beta_1 \beta_2]^T = 0$ , which contradicts that  $(\bar{L}_f, \bar{L}_{fl}^T)$  is observable. The proof of the sufficiency is completed.

**3.2.2. Necessity.** Supposed  $(\bar{L}_f, \bar{L}_{fl}^T)$  is unobservable, then there must exist an eigenvalue  $\lambda$  of  $\bar{L}_f$ , whose corresponding nonzero right eigenvector is  $[\beta_1 \beta_2]^T$ , such that  $\bar{L}_f [\beta_1 \beta_2]^T = \lambda [\beta_1 \beta_2]^T$  and  $\bar{L}_{fl}^T [\beta_1 \beta_2]^T = 0$ . Therefore, we can also assert that  $L_f$  has an eigenvalue  $u = (\lambda^2 - \lambda c_2 / c_1)$  with the corresponding nonzero right eigenvector  $\beta_1$ . Then,  $L_f \beta_1 = u \beta_1$  and  $L_{fl}^T \beta_1 = 0$ , which contradicts that  $(L_f, L_{fl}^T)$  is observable. The proof of the necessity is completed.

Note that, from Theorem 5, the nonzero feedback gains  $c_1$  and  $c_2$  can be designed to ensure that equation  $s^2 - c_2 s - c_1 u = 0$  has roots. Therefore, we can obtain the following result.

**Theorem 6.** *The second-order MAS (18) is observable if system (18) satisfies*

- (i)  $\text{rank}[(\lambda_i^2 - \lambda_i c_2) I_n - c_1 L_f] = n$ ,
- (ii)  $\text{rank}[\lambda_i c_1 L_{fl}^T] = n$ , or
- (iii)  $\text{rank} \begin{bmatrix} (\lambda_i^2 - \lambda_i c_2) I_n - c_1 L_f \\ \lambda_i c_1 L_{fl}^T \end{bmatrix} = n$ .

*Proof.* From Proposition 1 (4), we can have

$$\begin{aligned} \text{rank} \begin{bmatrix} \lambda_i I_{2n} - \bar{L}_f \\ \bar{L}_{fl}^T \end{bmatrix} &= \text{rank} \begin{bmatrix} \lambda_i I_n & -I_n \\ -c_1 L_f & (\lambda_i - c_2) I_n \\ 0 & c_1 L_{fl}^T \end{bmatrix} \\ &= \text{rank} \begin{bmatrix} 0 & -I_n \\ (\lambda_i^2 - \lambda_i c_2) I_n - c_1 L_f & (\lambda_i - c_2) I_n \\ \lambda_i c_1 L_{fl}^T & c_1 L_{fl}^T \end{bmatrix}, \end{aligned} \quad (20)$$

where  $\lambda_i$  ( $i = 1, 2, \dots, 2n$ ) are the eigenvalues of  $\bar{L}_f$ . Obviously, system (18) is observable if system (18) satisfies one of the conditions (i), (ii), and (iii).  $\square$

**Remark 3.** Notice from Theorem 5 that the observability of the second-order MAS depends on the system couplings  $(L_f, L_{fl}^T)$ . However, from Theorem 6, we can select appropriate nonzero feedback gains  $c_1$  and  $c_2$  so that the observability of the second-order MAS depends only on the information topology between followers or the information topology from leaders to followers. This makes it easier to study the observability through designing the topological relationship of such system.

**3.3. Observability of High-Order MASs.** A high-order MAS consists of  $(n + l)$  agents, in which the former  $n$  agents are regarded as followers and the remaining  $l$  agents are regarded as leaders, and its dynamic equation is described by

$$\dot{x}_i^{(1)} = x_i^{(2)}, \dots, \dot{x}_i^{(m-1)} = x_i^{(m)}, \dot{x}_i^{(m)} = u_i, \quad i \in \mathcal{L}_{n+l}, \quad (21)$$

with

$$u_i = \sum_{j \in \mathcal{N}_i} a_{ij} \sum_{k=0}^{m-1} r_k (x_j^{(k+1)} - x_i^{(k+1)}), \quad (22)$$

where  $x_i^{(k+1)}$  ( $i = 1, \dots, n + l$ ) is the  $k$ -th order of  $x_i^{(1)}$ ,  $m$  is the order number, and  $r_0, \dots, r_{m-1}$  are nonzero feedback gains.

Let  $\tilde{x}_f \triangleq [x_1^{(1)}, \dots, x_n^{(1)}, \dots, x_1^{(m)}, \dots, x_n^{(m)}]^T$  and  $\tilde{x}_l \triangleq [x_{n+1}^{(1)}, \dots, x_{n+l}^{(1)}, \dots, x_1^{(m)}, \dots, x_{n+l}^{(m)}]^T$ . Based on the partition of follower agents and leader agents, the dynamics (21) and (22) can be rewritten as follows:

$$\begin{cases} \dot{\tilde{x}}_f = \tilde{L}_f \tilde{x}_f + \tilde{L}_{fl} \tilde{x}_l, \\ y = \tilde{L}_{fl}^T \tilde{x}_f, \end{cases} \quad (23)$$

where

$$\begin{aligned} \tilde{L}_f &= \begin{bmatrix} 0 & 1 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \\ 0 & 0 & \dots & 0 \end{bmatrix}_{m \times m} \otimes I_n + \begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \\ r_0 & r_1 & \dots & r_{m-1} \end{bmatrix}_{m \times m} \otimes L_f, \\ \tilde{L}_{fl} &= \begin{bmatrix} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \\ r_0 & r_1 & \dots & r_{m-1} \end{bmatrix}_{m \times m} \otimes L_{fl}. \end{aligned} \quad (24)$$

For the observability of high-order MASs, we can have the following result.

**Theorem 7.** *The high-order MAS (23) is observable iff  $(L_f, L_{fl}^T)$  is observable.*

*Proof.* Supposed  $\lambda$  is the eigenvalue of  $\tilde{L}_f$ , whose corresponding right eigenvector is denoted as  $\beta = [\beta_1 \beta_2 \dots \beta_m]^T$ , then  $\tilde{L}_f [\beta_1, \beta_2, \dots, \beta_m]^T = \lambda [\beta_1, \beta_2, \dots, \beta_m]^T$ , which follows that

$$\begin{cases} \beta_2 = \lambda \beta_1, \\ \beta_3 = \lambda \beta_2, \\ \vdots \\ \beta_m = \lambda \beta_{m-1}, \\ r_0 L_f \beta_1 + r_1 L_f \beta_2 + \dots + r_{m-1} L_f \beta_m = \lambda \beta_m. \end{cases} \quad (25)$$

Consequently,

$$\begin{cases} \beta_2 = \lambda \beta_1, \\ \beta_3 = \lambda^2 \beta_1, \\ \vdots \\ \beta_m = \lambda^{m-1} \beta_1, \\ r_0 L_f \beta_1 + r_1 L_f \lambda \beta_1 + \dots + r_{m-1} L_f \lambda^{m-1} \beta_1 = \lambda^m \beta_1, \\ (r_0 + r_1 \lambda + \dots + r_{m-1} \lambda^{m-1}) L_f \beta_1 = \lambda^m \beta_1. \end{cases} \quad (26)$$

Therefore, we can have the polynomial  $f(\lambda) = r_0 + r_1 \lambda + \dots + r_{m-1} \lambda^{m-1} \neq 0$ . Otherwise, if  $f(\lambda) = 0$ , it implies that  $\beta_1 = 0$  or  $\lambda = 0$ . If  $\beta_1 = 0$ , then  $\beta_1 = \beta_2 = \beta_3 = \dots = \beta_m = 0$ ; and if  $\lambda = 0$ , then  $r_0 = 0$  which contradicts the fact that  $r_0$  is a nonzero gain. Let  $u = \lambda^m / r_0 + r_1 \lambda + \dots + r_{m-1} \lambda^{m-1}$ , then  $L_f \beta_1 = u \beta_1$ , where  $u$  is the root of equation  $s^m - u r_{m-1} s^{m-1} - \dots - u r_0 = 0$  with

$$\begin{cases} \beta_2 = s \beta_1, \\ \beta_3 = s^2 \beta_1, \\ \vdots \\ \beta_m = s^{m-1} \beta_1. \end{cases} \quad (27)$$

Next, we will show the fact that  $(\tilde{L}_f, \tilde{L}_{fl}^T)$  is observable iff  $(L_f, L_{fl}^T)$  is observable. The following proof is similar to that of Theorem 5, omitted here.

*Remark 4.* This paper has investigated not only the observability of first-order MASs but also the observability of second-order and high-order cases. Higher-order systems have more complex and higher dimensional system matrices, so it is more difficult to establish the conditions on the observability and more complex to calculate using observability criteria.

## 4. Simulations

**4.1. Example 1.** A five-agent first-order MAS is considered, where agents 1–3 and agents 4–5 are regarded as followers and leaders, respectively. The network topology is described in Figure 2. The system matrices are as follows:

$$\begin{aligned} L_f &= \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}, \\ L_{fl}^T &= \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}. \end{aligned} \quad (28)$$

By MATLAB, we can compute and  $\lambda(L_f) = \{0.5858, 2.0000, 3.4142\}$  and  $L_{fl}^T U = \begin{bmatrix} -0.5000 & 0.7071 & 0.5000 \\ -0.5000 & -0.7071 & 0.5000 \end{bmatrix}$ . Obviously, it is easy to find that  $\lambda(L_f)$  are all distinct, and all elements of every column of  $L_{fl}^T U$  are nonzero. From Theorem 4, such system is observable.




FIGURE 2: The topology with 5 agents.




FIGURE 3: A regular triangle array.




FIGURE 4: The topology with 7 agents.




FIGURE 5: A straight line array.




FIGURE 6: Position errors.




FIGURE 7: Velocities.

Figure 3 describes the initial states, trajectories, and final states of the followers, where the random initial states of all followers (the black stars) are to be finally steered to a regular triangle array (the black circles).

**4.2. Example 2.** A seven-agent second-order MAS is considered, where agents 1–5 and agents 6–7 are regarded as followers and leaders, respectively. The network topology is described in Figure 4. Let  $c_1 = c_2 = 1$  for the sake of simplicity. The system matrices are as follows:

$$\bar{L}_f = \begin{bmatrix} 0 & I_5 \\ L_f & I_5 \end{bmatrix} = \left[ \begin{array}{ccccc|c} & & & & & I_5 \\ 2 & -1 & -1 & 0 & 0 & \\ -1 & 3 & 0 & -1 & 0 & \\ -1 & 0 & 2 & -1 & 0 & I_5 \\ 0 & -1 & -1 & 3 & -1 & \\ 0 & 0 & 0 & -1 & 2 & \end{array} \right], \quad (29)$$

$$\bar{L}_{fl}^T = \begin{bmatrix} 0 & L_{fl}^T \end{bmatrix} = \left[ \begin{array}{c|ccccc} 0 & 0 & -1 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 0 & -1 \end{array} \right].$$

By computing,

$$\lambda(\bar{L}_f) = \{-1.7389, -1.3028, -1.1391, -0.8292, -0.2308, 1.2308, 2.7389, 1.8292, 2.3028, 2.1391\}. \quad (30)$$

For each  $\lambda$ , we have  $\text{rank}[(\lambda^2 + \lambda)I_5 - L_f] = 5$ . According to Theorem 6, the second-order multiagent system (18) is observable.

Since the second-order multiagent system (18) is observable, the followers finally achieve the expected states. Figure 5 describes the initial states, trajectories, and final states of the followers, where the random initial states of all followers (the black stars) are to be finally steered to a straight line array (the black circles). Arrows represent the direction of each agent's movement. Figures 6 and 7 are the position errors and velocities of the followers, respectively.

## 5. Conclusion

This paper has studied the observability of first-order/second-order/high-order MASs with relative protocol on

fixed topology, respectively, and brings the three kinds of networks into a unified framework. It has proved that the observability of MASs is depended only on structural pair  $(L_f, L_{f1}^T)$ . However, for second-order or high-order MASs, it is very hard to deal with the observability because of complexity of the topology structure and lack of theoretical research tools. Our main result shows an advantage of a unified scheme for the observability of MASs and increasing the theoretical studying on the observability concept for MASs. Future research studies are directed on observability of MASs with switching topology, time delays, and observability of general linear discrete-time and continuous-time MASs, switched MASs and heterogeneous MASs.

## Data Availability

Our paper does not use the data information.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China under grant nos. 61773023, 61773416, and 61873318, the Frontier Research Funds of Applied Foundation of Wuhan under grant no. 2019010701011421, the Natural Science Foundation of Hubei Province of China under grant no. 2018CFA058, the Fundamental Research Funds for the Central Universities under grant no. HUST: 2019kfyXKJC022, and the Program for HUST Academic Frontier Youth Team under grant no. 2018QYTD07.

## References

- [1] C. W. Reynolds, "Flocks, herds and schools: a distributed behavioral model," *ACM SIGGRAPH Computer Graphics*, vol. 21, no. 4, pp. 25–34, 1987.
- [2] J. K. Parrish, S. V. Viscido, and D. Grünbaum, "Self-organized fish schools: an examination of emergent properties," *The Biological Bulletin*, vol. 202, no. 3, pp. 296–305, 2002.
- [3] H. Su, Y. Ye, Y. Qiu, Y. Cao, and M. Z. Q. Chen, "Semi-global output consensus for discrete-time switching networked systems subject to input saturation and external disturbances," *IEEE Transactions on Cybernetics*, vol. 49, no. 11, pp. 3934–3945, 2019.
- [4] X. Wang and H. Su, "Self-triggered leader-following consensus of multi-agent systems with input time delay," *Neurocomputing*, vol. 330, pp. 70–77, 2019.
- [5] R. Olfati-Saber and R. M. Murray, "Consensus problems in networks of agents with switching topology and time-delays," *IEEE Transactions on Automatic Control*, vol. 49, no. 9, pp. 1520–1533, 2004.
- [6] R. Luo, H. Su, and Y. Zeng, "Chaos control and synchronization via switched output control strategy," *Complexity*, vol. 2017, Article ID 6125102, 11 pages, 2017.
- [7] X. Wang and H. Su, "Consensus of hybrid multi-agent systems by event-triggered/self-triggered strategy," *Applied Mathematics and Computation*, vol. 359, pp. 490–501, 2019.
- [8] B. Liu, T. Chu, L. Wang, and G. Xie, "Controllability of a leader-follower dynamic network with switching topology," *IEEE Transactions on Automatic Control*, vol. 53, no. 4, pp. 1009–1013, 2008.
- [9] L. Wang, F. Jiang, G. Xie, and Z. Ji, "Controllability of multi-agent systems based on agreement protocols," *Science in China Series F: Information Sciences*, vol. 52, no. 11, pp. 2074–2088, 2009.
- [10] B. Liu, T. Chu, L. Wang, Z. Zuo, G. Chen, and H. Su, "Controllability of switching networks of multi-agent systems," *International Journal of Robust and Nonlinear Control*, vol. 22, no. 6, pp. 630–644, 2012.
- [11] B. Liu, H. Su, R. Li, D. Sun, and W. Hu, "Switching controllability of discrete-time multi-agent systems with multiple leaders and time-delays," *Applied Mathematics and Computation*, vol. 228, pp. 571–588, 2014.
- [12] Z. Ji, H. Lin, and H. Yu, "Protocols design and uncontrollable topologies construction for multi-agent networks," *IEEE Transactions on Automatic Control*, vol. 60, no. 3, pp. 781–786, 2015.
- [13] B. Liu, Y. Han, F. Jiang, H. Su, and J. Zou, "Group controllability of discrete-time multi-agent systems," *Journal of the Franklin Institute*, vol. 353, no. 14, pp. 3524–3559, 2016.
- [14] Y. Guan and L. Wang, "Structural controllability of multi-agent systems with absolute protocol under fixed and switching topologies," *Science in China Series F: Information Sciences*, vol. 60, no. 9, Article ID 092203, 2017.
- [15] B. Liu, H. Su, F. Jiang, Y. Gao, L. Liu, and J. Qian, "Group controllability of continuous-time multi-agent systems," *IET Control Theory & Applications*, vol. 12, no. 11, pp. 1665–1671, 2018.
- [16] J. Qu, Z. Ji, C. Lin, and H. Yu, "Fast consensus seeking on networks with antagonistic interactions," *Complexity*, vol. 2018, Article ID 7831317, 15 pages, 2018.
- [17] Y. Liu and H. Su, "Containment control of second-order multi-agent systems via intermittent sampled position data communication," *Applied Mathematics and Computation*, vol. 362, Article ID 124522, 2019.
- [18] Z. Ji and H. Yu, "A new perspective to graphical characterization of multiagent controllability," *IEEE Transactions on Cybernetics*, vol. 47, no. 6, pp. 1471–1483, 2017.
- [19] X. Liu, Z. Ji, T. Hou, and H. Yu, "Decentralized stabilizability and formation control of multi-agent systems with antagonistic interactions," *ISA Transactions*, vol. 89, pp. 58–66, 2019.
- [20] X. Liu and Z. Ji, "Controllability of multiagent systems based on path and cycle graphs," *International Journal of Robust and Nonlinear Control*, vol. 28, no. 1, pp. 296–309, 2018.
- [21] H. G. Tanner, "On the controllability of nearest neighbor interconnections," in *Proceedings of the 43rd IEEE Conference on Decision and Control*, vol. 3, pp. 2467–2472, Nassau, The Bahamas, December 2004.
- [22] M. Ji and M. Egerstedt, "Observability and estimation in distributed sensor networks," in *Proceedings of the 46th IEEE Conference on Decision and Control*, pp. 4221–4226, New Orleans, LA, USA, December 2007.
- [23] D. Zelazo and M. Mesbahi, "On the observability properties of homogeneous and heterogeneous networked dynamic systems," in *Proceedings of the 47th IEEE Conference on Decision and Control*, pp. 2997–3002, Cancun, Mexico, December 2008.
- [24] R. Lozano, M. W. Spong, J. A. Guerrero, and N. Chopra, "Controllability and observability of leader-based multi-agent

- systems,” in *Proceedings of the 47th IEEE Conference on Decision and Control*, pp. 3713–3718, Cancun, Mexico, December 2008.
- [25] M. Franceschelli, S. Martini, M. Egerstedt, A. Bicchi, and A. Giua, “Observability and controllability verification in multi-agent systems through decentralized Laplacian spectrum estimation,” in *Proceedings of the 49th IEEE Conference on Decision and Control*, pp. 5775–5780, Atlanta, GA, USA, December 2010.
- [26] G. Parlangeli and G. Notarstefano, “On the observability of path and cycle graphs,” in *Proceedings of the 49th IEEE Conference on Decision and Control*, pp. 1492–1497, Atlanta, GA, USA, December 2010.
- [27] A. Chapman, M. Nabi-Abdolyousefi, and M. Mesbahi, “On the controllability and observability of Cartesian product networks,” in *Proceedings of the 51st IEEE Conference on Decision and Control*, pp. 80–85, Maui, Hawaii, USA, December 2012.
- [28] A. Y. Kibangou and C. Commault, “Algebraic characterization of observability in distance-regular consensus networks,” in *Proceedings of the 52nd IEEE Conference on Decision and Control*, pp. 1313–1318, Florence, Italy, December 2013.
- [29] A. Y. Kibangou and C. Commault, “Observability in connected strongly regular graphs and distance regular graphs,” *IEEE Transactions on Control of Network Systems*, vol. 1, no. 4, pp. 360–369, 2014.
- [30] N. O’Clery, Y. Yuan, G. Stan, and M. Barahona, “Observability and coarse graining of consensus dynamics through the external equitable partition,” *Physical Review E*, vol. 88, no. 4, Article ID 042805, 2013.
- [31] C. Liu and C. Li, “Reachability and observability of switched linear systems with continuous-time and discrete-time subsystems,” *International Journal of Control, Automation and Systems*, vol. 11, no. 1, pp. 200–205, 2013.
- [32] L. Tian, Y. Guan, and L. Wang, “Controllability and observability of switched multi-agent systems,” *International Journal of Control*, vol. 92, no. 8, pp. 1742–1752, 2019.
- [33] L. Sabattini, C. Secchi, and C. Fantuzzi, “Controllability and observability preservation for networked systems with time varying topologies,” in *Proceedings of the Preprints of the 19th World Congress, The International Federation of Automatic Control*, pp. 1837–1842, Cape Town, South Africa, August 2014.
- [34] Z. Lu, L. Zhang, and L. Wang, “Observability of multi-agent systems with switching topology,” *IEEE Transactions on Circuits and Systems II: Express Briefs*, vol. 64, no. 11, pp. 1317–1321, 2017.
- [35] L. Tian, Y. Guan, and L. Wang, “Controllability and observability of multi-agent systems with heterogeneous and switching topologies,” *International Journal of Control*, pp. 1–12, 2018.
- [36] Y.-Y. Liu, J.-J. Slotine, and A.-L. Barabasi, “Observability of complex systems,” *Proceedings of the National Academy of Sciences*, vol. 110, no. 7, pp. 2460–2465, 2013.
- [37] M. Ji and M. Egerstedt, “A graph-theoretic characterization of controllability for multi-agent systems,” in *Proceedings of the 2007 American Control Conference*, pp. 4588–4893, New York City, NY, USA, July 2007.

