Case Report

Management of an Unusual Maxillary Canine: A Rare Entity

**Jaya Nagendra Krishna Muppalla,¹ Krishnamurthy Kavuda,²
Rajani Punna,¹ and Amulya Vanapatla¹**

¹Kamineni Institute of Dental Sciences, Narketpally, Telangana 508254, India

²Awadh Dental College & Hospital, Jamshedpur, Jharkhand 831012, India

Correspondence should be addressed to Jaya Nagendra Krishna Muppalla; kittymjn@gmail.com

Received 14 July 2015; Accepted 29 November 2015

Academic Editor: Malka Ashkenazi

Copyright © 2015 Jaya Nagendra Krishna Muppalla et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Clinicians need to have intimate knowledge and thorough understanding of both pulp chamber and root canal anatomy. They should be aware of possibility of anatomical variations in the root canal system during endodontic treatment. Maxillary canines usually have single root and root canal but rarely may have single root with two root canals. This case describes a lengthier maxillary canine with two root canals.

1. Introduction

Thorough knowledge and understanding of pulp chamber and root canal system anatomy are essential for successful root canal therapy. Familiarity with variations in tooth anatomy and characteristic features in various racial groups can aid location and negotiation of canals [1]. Missed extra root canals are major reason for endodontic failure [2].

Pulp canal system is complex with branching and divisions throughout the root length. Vertucci (1984) classified the root canal configurations of human permanent teeth into various types ranging from single to three separate distinct canals [3].

Permanent maxillary canines are more commonly single rooted, single canal teeth. Presence of two root canals is a rarity [4]. Majority of them join in apical third and exit as single apical foramen [5].

This case report presents a permanent maxillary canine having two root canals exiting as single foramen.

2. Case Report

A 35-year-old female patient reported to dental clinic with a chief complaint of spontaneous pain from four days in maxillary left anterior region. Subjective symptoms include sharp, severe, continuous, throbbing pain and are aggravated

by taking hot foods and relieved by medication. Past medical history was noncontributory.

Oral examination revealed a deep carious lesion involving maxillary left canine (23, FDI System). The teeth was asymptomatic to palpation and tested negative using electrical pulp tester and gave an exaggerated severe response to cold test. No mobility was seen and tested negative for percussion. Radiographic examination revealed an abnormal root canal morphology. A diagnosis of acute irreversible pulpitis of 23 was established and endodontic treatment was planned.

Following local anesthesia with 2% lidocaine, the tooth was isolated with rubber dam [Hygienic; Coltene Whale-dent], and an endodontic access was made on palatal side with #1014 round bur and endo-Z carbide bur. The vital pulp tissue was extirpated and initially two canal orifices were located. Working length of 31 mm was measured by using K flex files #15 of length 31 mm (Figure 1). The palatal canal joined the buccal canal in apical third of root canal (Vertucci type II canal configuration). Crown down root canal preparation was done; the coronal and middle thirds were prepared using Gates Glidden drills #1–3 [Tulsa Dental, Dentsply] and apical preparation by hand K flex files [Dentsply] to size #40. Copious irrigation with 5.25% sodium hypochlorite, 17% EDTA were performed after use of each file. 2% chlorhexidine was used as final irrigant. The canals were dried with paper points and obturated with gutta-percha and AH Plus Sealer

FIGURE 1: Radiograph showing working length of maxillary canine with two separate canals.

FIGURE 2: Radiograph showing master cone gutta-percha.

(Figure 2) [Dentsply De Trey GmbH, Konstanz, Germany] using a lateral condensation technique (Figure 3). The patient was asymptomatic during the 3-month follow-up.

3. Discussion

Debridement of root canal to remove pulpal remnants, bacteria, and their byproducts before obturation is primary requisite for successful endodontic treatment. Being unable to locate and fill a canal results in failure of root canal therapy. Therefore, it is imperative to have knowledge of anatomic variations as endodontic success is related to canal debridement [6].

The diagnostic difficulty and possible canal superimposition on radiographic examination should be kept in mind when examining such cases. When locating extra canals, identification of periodontal ligament space that often projects onto root surface resembling a canal should be differentiated. Vertucci (1984) classified root canals according

FIGURE 3: Postoperative radiograph obturated with gutta-percha and AH Plus Sealer.

to number of canals present and their configuration into eight types [7].

Anatomic anomaly observed at first appointment should be checked for similar anomaly of tooth on the other side [8]. In the present case, radiographic image showed bilateral Vertucci type II [2-1] configuration.

Çalışkan et al. [9] studied root canal number, configuration, and ramifications of permanent teeth in Turkish population. They reported percentage of Vertucci type III [1-2-1] and type V [1-2] as 4.35 and 2.17, respectively. Alapati et al. [10] and Onay and Ungor [11] reported a maxillary right canine with type II canal configuration and Weisman [12] also reported a birooted maxillary left canine.

In the present case, the maxillary canines had an unusual root length of 31 mm which necessitated use of lengthier k flex files of 31 mm removing silicone directional stopper before use. They showed canal configuration of type II similar to that reported by Alapati et al. and Onay and Ungor. Two distinct canal orifices were located in labial/palatal direction which joined in apical third, forming a type II configuration.

Teeth with type II configuration during treatment may pose problems. The canal that is in line with the main passage is usually amenable to adequate enlarging and obturation procedures; the preparation and filling of the other canal are often extremely difficult [13].

A thorough knowledge of root canal anatomy and operator skill are essential for endodontic success. Careful clinical examination with radiographs from several different angles may lead to suspicion or identification of additional canals and leads to higher possible success [3].

4. Conclusion

Several variations exist in the root canal system and clinicians should be aware of the variations for complete infection removal and prevention of reinfection. Special care with careful endodontic exploration, different angle radiographs, and magnification with surgical microscope aids in detection and treatment of extra canals.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] P. Krasner and H. J. Rankow, "Anatomy of the pulp-chamber floor," *Journal of Endodontics*, vol. 30, no. 1, pp. 5–16, 2004.
- [2] M. Hulsmann and E. Schafer, "Problems in gaining access to the root canal system," in *Problems in Endodontics: Etiology, Diagnosis and Treatment*, H. Michael and S. Edgar, Eds., pp. 145–172, Quintessence Publishing, Berlin, Germany, 1st edition, 2009.
- [3] N. Bolla and S. Kavuri, "Maxillary canine with two root canals," *Journal of Conservative Dentistry*, vol. 14, no. 1, pp. 80–82, 2011.
- [4] R. C. Burns and L. S. Buchanan, "Tooth morphology and access openings," in *Pathways of the Pulp*, S. Cohen and R. C. Burns, Eds., pp. 128–178, Mosby, Boston, Mass, USA, 6th edition, 1994.
- [5] S. V. Ravi, E. Simon, K. Thomas, and A. Ravi, "Maxillary canine with two root canals: a case report," *Indian Journal of Dental Research*, pp. 69–71, 2012.
- [6] S. Seltzers and I. B. Bender, "The Interrelationship of pulp and periodontal disease," in *The Dental Pulp*, S. Seltzers and I. B. Bender, Eds., pp. 303–323, All India Publishers and Distributors, Chennai, India, 1st edition, 2000.
- [7] F. J. Vertucci, "Root canal anatomy of the human permanent teeth," *Oral Surgery, Oral Medicine, Oral Pathology*, vol. 58, no. 5, pp. 589–599, 1984.
- [8] D.-R. Shin, J.-M. Kim, D.-S. Kim, S.-Y. Kim, P. V. Abbott, and S.-H. Park, "A maxillary canine with two separated root canals: a case report," *Journal of Korean Academy of Conservative Dentistry*, vol. 36, no. 5, pp. 431–435, 2011.
- [9] M. K. Çalışkan, Y. Pehlivan, F. Sepetçioğlu, M. Türkün, and S. Ş. Tuncer, "Root canal morphology of human permanent teeth in a Turkish population," *Journal of Endodontics*, vol. 21, no. 4, pp. 200–204, 1995.
- [10] S. Alapati, E. I. Zaatar, M. Shyama, and N. Al-Zuhair, "Maxillary canine with two root canals," *Medical Principles and Practice*, vol. 15, no. 1, pp. 74–76, 2006.
- [11] O. E. Onay and M. Unger, "Maxillary Canines with two root canals," *Hacettepe Üniversitesi Diş Hekimliği Fakültesi'nin Dergisi*, vol. 32, pp. 20–24, 2008.
- [12] M. I. Weisman, "A rare occurrence: a bi-rooted upper canine," *Australian Endodontic Journal*, vol. 26, no. 3, pp. 119–120, 2000.
- [13] G. Anuraag, S. Anurag, B. Ridhi, and A. Kunal, "Maxillary canine with two root canals: a case report," *Journal of Dental Sciences & Oral Rehabilitation*, pp. 32–34, 2013.