Case Report

Idiopathic Central Precocious Puberty Associated with 11 Mb *De Novo* Distal Deletion of the Chromosome 9 Short Arm

Mariangela Cisternino,¹ Erika Della Mina,² Laura Losa,¹ Alexandra Madè,¹ Giulia Rossetti,¹ Lorenzo Andrea Bassi,¹ Giovanni Pieri,¹ Baran Bayindir,² Jole Messa,² Orsetta Zuffardi,^{2,3} and Roberto Ciccone²

¹ Department of Pediatrics, IRCCS Policlinico San Matteo, University of Pavia, Piazzale Golgi 19, 27100 Pavia, Italy

² Department of Molecular Medicine, University of Pavia, Via Forlanini 6, 27100 Pavia, Italy

³ IRCCS C. Mondino National Neurological Institute Foundation, Via Mondino 2, 27100 Pavia, Italy

Correspondence should be addressed to Mariangela Cisternino; m.cisternino@smatteo.pv.it

Received 6 June 2013; Accepted 11 July 2013

Academic Editors: D. J. Bunyan, C.-W. Cheng, P. Morrison, and M. Velinov

Copyright © 2013 Mariangela Cisternino et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We report a girl with a *de novo* distal deletion of 9p affected by idiopathic central precocious puberty and intellectual disability. Genome-wide array-CGH revealed a terminal deletion of about 11 Mb, allowing to define her karyotype as 46; XX, del(9)(p23-pter). To our knowledge, this is the second reported case of precocious puberty associated with 9p distal deletion. A third case associates precocious puberty with a more proximal 9p deletion del(9)(p12p13.3). In our case, more than 40 genes were encompassed in the deleted region, among which, DMRT1 which is gonad-specific and has a sexually dimorphic expression pattern and ERMP1 which is required in rats for the organization of somatic cells and oocytes into discrete follicular structures. Although we cannot exclude that precocious puberty in our del(9p) patient is a coincidental finding, the report of the other two patients with 9p deletions and precocious puberty indeed suggests a causative relationship.

1. Introduction

Central precocious puberty (CPP) is classically defined by the appearance of sexual secondary characteristics before the age of 8 years in girls and 9 years in boys [1]. It is caused by a premature activation of the hypothalamus-pituitary-gonadal axis. CPP may be either idiopathic or associated with occult intracranial lesion, mainly hypothalamic hamartoma or astrocytoma and noncancerous CNS disorders [2, 3]. This condition may cause early epiphyseal maturation with compromised final height as well as psychological stress [4, 5].

Chromosome 9p deletion syndrome (OMIM#158170) is a well-recognized entity, caused by a constitutional monosomy of a portion of 9p of different sizes in different patients. It was first described by Alfi et al. in 1973 [6]. Until now, approximately 180 cases have been published [7].

The most common features of monosomy 9p syndrome, as described by Swinkels et al. [8], include developmental and psychomotor delay, trigonocephaly, flat midface, short palpebral fissures, highly arched eyebrows, low-set ears, short flat nose with anteverted nostrils, thin upper lip, long philtrum, high palate, micrognathia, short neck, nipple hypertelorism, tapering fingers, flat feet, hypotonia, and developmental sex disorders in XY subjects. The critical region for a consensus phenotype has been reported to be located in a 300 Kb region on 9p22.3 [8]. Approximately half of the cases are due to *de novo* deletions of 9p, the remaining ones to unbalanced translocations with a derivative 9p chromosome. Few cases have been reported with 9p distal deletion concomitant to 9q distal duplication, with some of them due to a parental chromosome 9 inversion.

To our knowledge, only one case of distal 9p deletion [del9(p22pter)] [9] and a second case of proximal 9p deletion

FIGURE 1: Pictures of the girl affected by 9p-syndrome. (a), (b): face and profile of the patient at the age of 7 years. (c), (d): face and profile of the patient at the age of 14 years. (e), (f): shortness of 4th metacarpal and metatarsal.

[del9(p12p13.3)] [10] have been reported in association with precocious puberty.

Here, we report a girl with distal 9p deletion with idiopathic central precocious puberty and mental impairment.

2. Case Presentation

The proband is a girl. We have been following her from the age of 7 years and 4 months. She is the only child of nonconsanguineous healthy Italian parents. At her birth, her mother was 29 years old, and her father was 31 years old. She was born at the 40th week of gestation after uneventful pregnancy and delivery. Her birth weight was 3350 g (50th centile), her length was 50 cm (50th centile), and her head circumference was 35 cm (75th centile). She had neonatal jaundice requiring one-day phototherapy. Her psychomotor development was delayed: she began to walk by the age of 24

months. She suffered from chronic constipation for 2 and a half years. At the age of 6 years, she presented a left inguinal hernia which was surgically reduced.

She has a moderate intellectual disability with good social adaptation, and presently, she is enrolled in a public school with the aid of a tutor.

At 7 years and 4 months of age, she was referred to our unit of Pediatric Endocrinology for evaluation of precocious puberty since pubarche, axillarche, and axillary sweating were noted at the age of 7 years, followed by unilateral thelarche and pubertal spurt. Clinical evaluation showed several facial anomalies (Figure 1), including low anterior hair line, low-set ears, synophrys, short nose, long philtrum, and wide mouth with thin vermilions of the lower and upper lip. In addition, geographic tongue, dental crowding, slightly arched palate, bilateral short 4th and 5th metacarpal and metatarsal, and signs of ungueal decalcification were noted. She was

128 cm tall (H-SDS +1.19), and her weight was 28.3 kg (90th percentile) with a BMI of 17.3 (BMI-SDS +1.35). Pubertal stage (according to Marshall and Tanner) [12] was unilateral B2, PH2, and A2. Bone age was of 10 years according to the Greulich and Pyle atlas [13]. Predictable adult height (according to the Bayley & Pinneau method) was of 154.5 cm (target height 160.2 cm).

Hormonal investigations showed basal luteinizing hormone (LH) of 0.1 IU/L and basal follicle-stimulating hormone (FSH) of 5.4 IU/L (normal prepubertal values: LH < 0.5 IU/L and FSH < 7 IU/L), and the serum estradiol (E2) level was slightly increased 26.2 pg/mL (normal prepubertal values < 15 pg/mL). The GnRH stimulation test showed an LH and FSH response consistent with a prepubertal stage (LH peak 3.7 IU/L, FSH peak 20.7 IU/L; normal prepubertal values: LH < 7 IU/L, FSH 18.2 ± 2.62 IU/L). Measurement of adrenal androgen levels showed dehydroepiandrosterone sulphate (DHEA-S) values of 0.6 mcg/mL which were consistent with adrenarche. Pelvic ultrasonography (US) showed no masses, and ovarian and uterine volumes were not enlarged and were normal for chronological age. Premature adrenarche was then diagnosed; however, because of the bone age advancement and the increased E2 levels, the girl was subsequently re-examined.

Eight months later, at the age of 8 years, the patient showed a progression of thelarche, which became bilateral, and of pubarche, whereas the growth rate remained higher than normal; the GnRH stimulation test showed a pubertal LH response (peak 11.8 IU/L), and the pelvic US showed increased size of the uterus (6 cm³) and of the ovaries (2 cm³, bilaterally). The uterus was pear-shaped, and the ratio between the fundal and cervical anteroposterior diameters was >1. The girl was then diagnosed as a central precocious puberty. Brain MRI did not show any abnormality, including in the hypothalamic and pituitary regions; therefore, the origin of the precocious puberty was considered idiopathic. A suppressive treatment of puberty with a GnRH analogue (3.75 mg per month of a depot suspension of leuprolide acetate) was administered until the age of 12 years. The menarche appeared at the age of 13.1 years (final height 161.5 cm, H-SDS -0.11), followed by irregular menses and polymenorrhea which required periodical iron supplementation. At the age of 14 years, she complained of a unilateral hypertrophy of the labia minora causing local pain, irritation, and psychosocial distress, which required labiaplasty.

Additional investigations, during years, showed normal hepatic, renal, and thyroid function; hormonal profile was normal except for elevated levels of prolactin (max. 57.4 ng/mL) which was probably functional as no hypophyseal adenoma was found with MRI. No treatment for hyperprolactinemia was necessary due to the lack of specific symptoms, that is, galactorrhea and amenorrhea.

EEG showed focal paroxysms in the right hemisphere, not requiring antiepileptic therapy.

3. Cytogenetics and Array-CGH

Standard karyotype revealed a normal female karyotype.

A terminal deletion of the short arm of chromosome 9 [46, XX.ish.del(9p) (pVYS234B-9)] was detected by subtelomeric fluorescence *in situ* hybridization (FISH) analysis (Vysis kit).

Conventional cytogenetics and FISH analysis of patient's parents demonstrated that the deletion occurred *de novo*.

Genome-wide array-CGH has been performed using the Human Genome CGH Microarray Kit 4x44K (Agilent Technologies, Santa Clara, CA, USA) according to the manufacturers' protocol [14]. The investigation was made using a female genomic DNA pool (Promega Ltd.) as reference, and results has been referred as UCSC hg18 (NCBI Build 36), March 2006.

This analysis revealed a terminal deletion of about 11 Mb in the short arm of chromosome 9 [del(9)(p23-pter)]. Proximal breakpoint is between 11,259549 Mb (last deleted) and 11,447340 Mb (first not deleted); the last oligomer present on the platform was localized at 204,367 Kb (Figure 2).

4. Discussion

We present a case of distal 9p deletion which came to our attention because of idiopathic central precocious puberty.

About 180 cases of monosomy 9p syndrome have already been described [7]. The cases that presented del(9p) as the sole anomaly usually have a *de novo* mutation [15], which was the case of our patient. In the majority of cases, the breakpoint occurs at 9p21 [16]; however, many clinical features are similar regardless of the length of the deletion [15].

Even if our patient is different from previous del(9p) cases in the absence of trigonocephaly, flat nasal bridge, and the large number of digital whorls, she has many features in common with classic 9p-syndrome such as developmental delay and moderate mental retardation, long philtrum, arched palate, low-set ears, low anterior hair line, cardiac defect, and inguinal hernia. Besides, she has some physical characteristics that are not classical features of 9p-syndrome such as short 4th metacarpal and metatarsal and geographic tongue (Table 1).

However, what we consider to be important in this case is the fact that our patient was affected by central precocious puberty, from the age of 7 years, requiring suppressive therapy. To our knowledge, this is the second reported case of precocious puberty associated to 9p distal deletion. The first case was a boy described in 1979 by Funderburk et al. [9] who carried a *de novo* [del9(p22->pter)]. The unusual features of the boy were precocious puberty from the age of 8 years and 10 months and hexadactyly. Precocious puberty was not mentioned in other cases with distal 9p deletion that should have already undergone pubertal development according to their age. Several del(9p) patients had anomalous external genitalia such as hypoplastic labia majora and prominent labia minora [9], as was observed in this case.

In our patient, more than 40 genes were encompassed in the deleted region. Some of these genes are involved in an insulin metabolism pathway. Increased insulin and IGF1 levels were found elevated in girls with CPP, suggesting a causal interrelation between CPP and insulin secretion [17, 18]. Unfortunately, the insulin secretion was not studied at

TABLE 1: Clinical features of 9p-syndrome, as described in OMIM web site [11], compared to the clinical features presented by our patient affected of 9p-syndrome due to 9p24.3-p23 deletion.

Category	Features	Present case
Head	Trigonocephaly	No
Face	Midface hypoplasia	No
	Long philtrum	Yes
	Micrognathia	No
Ears	Low-set ears	Yes
	Malformed ears	No
	Posteriorly angulated ears	Yes
Eyes	Upslanting palpebral fissures	Yes
	Hypertelorism	Yes
	Epicanthal folds	Yes
	Small palpebral fissures	Yes
	Myopia	No
	High-arched eyebrows	Yes
Nose	Flat nasal bridge	Yes
	Anteverted nares	No
	Choanal atresia	No
Mouth	Thin upper lip	Yes
	Microstomia	No
	High narrow palate	Yes
Neck	Short neck	No
Heart	Heart murmurs	Yes
	Congenital cardiac malformations	No
	Atrial septal defect	No
	Ventricular septal defect	No
	Patent ductus arteriosus	No
Breasts	Widely spaced nipples	No
Abdomen	Inguinal hernia	Yes
	Omphalocele	No
Skeletal	Scoliosis	No
	Tapering fingers	No
	Pes planus	No
Skin, nails, and hair	Pale skin	Yes
	Hyperconvex nails	No
	High arched eyebrows	No
CNS	Mental retardation	Yes
	Delayed psychomotor development	Yes
	Speech delay	Yes
	Hypotonia	Yes
		Central precocious puberty
		Short metacarpal
		Short metatarsal
		Wide mouth
		Dental crowding
		Geographic tongue
		Ungueal decalcification
		Low anterior hairline
		Synophrys

Obviously, we cannot exclude that precocious puberty in our del(9p) patient is a coincidental finding; although, the report of other two patients with 9p deletions and precocious

The association of moderate mental retardation and CPP has been described in patients carrying several genetic anomalies, other than the 9p distal deletion, detected with the CGH array technique and the fluorescence in situ hybridization analysis (FISH) [21–24]. The molecular basis of this association remains unknown, but it is likely that multiple gene aberrations are responsible for this association.

All authors declare that they do not have a direct financial relation with the companies mentioned in the paper that might lead to a conflict of interests for any of the authors.

[1] A. Rogol and R. M. Blizzard, "Variations and disorders of pubertal development," in *Wilkins' the Diagnosis and Treatment*

- of *Endocrine Disorders in Childhood and Adolescence*, M. S. Kappy, R. M. Blizzard, and C. J. Migeon, Eds., pp. 857–917, Charles Thomas, Springfield, Ill, USA, 1994.
- [2] M. Cisternino, T. Arrigo, A. M. Pasquino et al., “Etiology and age incidence of precocious puberty in girls: a multicentric study,” *Journal of Pediatric Endocrinology and Metabolism*, vol. 13, supplement 1, pp. 695–701, 2000.
 - [3] M. Chalumeau, C. G. Hadjiathanasiou, S. M. Ng et al., “Selecting girls with precocious puberty for brain imaging: validation of European evidence-based diagnosis rule,” *Journal of Pediatrics*, vol. 143, no. 4, pp. 445–450, 2003.
 - [4] R. Brauner, L. Adan, F. Malandry, and D. Zantleifer, “Adult height in girls with idiopathic true precocious puberty,” *Journal of Clinical Endocrinology and Metabolism*, vol. 79, no. 2, pp. 415–420, 1994.
 - [5] G. B. Kletter and R. P. Kelch, “Clinical review 60: Effects of gonadotropin-releasing hormone analog therapy on adult stature in precocious puberty,” *Journal of Clinical Endocrinology and Metabolism*, vol. 79, no. 2, pp. 331–334, 1994.
 - [6] O. Alfí, G. N. Donnell, and B. F. Crandall, “Deletion of the short arm of chromosome # 9 (46,9p-). A new deletion syndrome,” *Annales de Genetique*, vol. 16, no. 1, pp. 17–22, 1973.
 - [7] R. Onesimo, D. Orteschi, M. Scalzone et al., “Chromosome 9p deletion syndrome and sex reversal: novel findings and redefinition of the critically deleted regions,” *American Journal of Medical Genetics A*, vol. 158, no. 9, pp. 2266–2271, 2012.
 - [8] M. E. M. Swinkels, A. Simons, D. F. Smeets et al., “Clinical and cytogenetic characterization of 13 Dutch patients with deletion 9p syndrome: delineation of the critical region for a consensus phenotype,” *American Journal of Medical Genetics A*, vol. 146, no. 11, pp. 1430–1438, 2008.
 - [9] S. J. Funderburk, R. S. Sparkes, and I. Klisak, “The 9p- syndrome,” *Journal of Medical Genetics*, vol. 16, no. 1, pp. 75–79, 1979.
 - [10] G. Eshel, E. Lahat, O. Reish, and J. Barr, “Neurodevelopmental and behavioral abnormalities associated with deletion of chromosome 9p,” *Journal of Child Neurology*, vol. 17, no. 1, pp. 50–51, 2002.
 - [11] OMIM, “Online Mendelian Inheritance in Man,” <http://www.omim.org/clinicalSynopsis/158170>.
 - [12] W. A. Marshall and J. M. Tanner, “Variations in pattern of pubertal changes in girls,” *Archives of Disease in Childhood*, vol. 44, no. 235, pp. 291–303, 1969.
 - [13] W. W. Greulich and S. I. Pyle, *Radiographic Atlas of Skeletal Development of the Hand and Wrist*, Stanford University Press, Stanford, Calif, USA, 2nd edition, 1959.
 - [14] N. Resta, R. Giorda, R. Bagnulo et al., “Breakpoint determination of 15 large deletions in Peutz-Jeghers subjects,” *Human Genetics*, vol. 128, no. 4, pp. 373–382, 2010.
 - [15] J. L. Huret, C. Leonard, B. Forestier, M. O. Rethore, and J. Lejeune, “Eleven new cases of del(9p) and features from 80 cases,” *Journal of Medical Genetics*, vol. 25, no. 11, pp. 741–749, 1988.
 - [16] D. W. Bianchi, “Chromosome 9, partial monosomy 9p,” in *Birth Defects Encyclopedia*, M. L. Buyse, Ed., pp. 353–354, Blackwell Scientific, Malden, Mass, USA, 1990.
 - [17] K. Sørensen, L. Aksglaede, J. H. Petersen, A. M. Andersson, and A. Juul, “Serum IGF1 and insulin levels in girls with normal and precocious puberty,” *European Journal of Endocrinology*, vol. 166, no. 5, pp. 903–910, 2012.
 - [18] M. Cisternino, A. Borghesi, L. Bacchella et al., “Increased insulin response to intravenous glucose tolerance test in girls with central precocious puberty,” *Hormone Research*, vol. 58, no. S2, p. 300, 2002.
 - [19] A. D. Krentz, M. W. Murphy, A. L. Sarver, M. D. Griswold, V. J. Bardwell, and D. Zarkower, “DMRT1 promotes oogenesis by transcriptional activation of Stra8 in the mammalian fetal ovary,” *Developmental Biology*, vol. 356, no. 1, pp. 63–70, 2011.
 - [20] C. Garcia-Rudaz, F. Luna, V. Tapia et al., “FxnA, a novel gene differentially expressed in the rat ovary at the time of folliculogenesis, is required for normal ovarian histogenesis,” *Development*, vol. 134, no. 5, pp. 945–957, 2007.
 - [21] L. Mutesa, A. C. Hellin, M. Jamar, G. Pierquin, V. Bours, and A. Verloes, “Precocious puberty associated with partial trisomy 18q and monosomy 11q,” *Genetic Counseling*, vol. 18, no. 2, pp. 201–207, 2007.
 - [22] K. Buiting, D. Kanber, J. I. Martín-Subero et al., “Clinical features of maternal uniparental disomy 14 in patients with an epimutation and a deletion of the imprinted DLK1/GTL2 gene cluster,” *Human Mutation*, vol. 29, no. 9, pp. 1141–1146, 2008.
 - [23] H. Lybæk, K. H. Ørstavik, T. Prescott et al., “An 8.9Mb 19p13 duplication associated with precocious puberty and a sporadic 3.9Mb 2q23.3q24.1 deletion containing NR4A2 in mentally retarded members of a family with an intrachromosomal 19p-into-19q between-arm insertion,” *European Journal of Human Genetics*, vol. 17, no. 7, pp. 904–910, 2009.
 - [24] T. Koshio, S. Sakazume, H. Kawame et al., “De-novo balanced translocation between 7q31 and 10p14 in a girl with central precocious puberty, moderate mental retardation, and severe speech impairment,” *Clinical Dysmorphology*, vol. 17, no. 1, pp. 31–34, 2008.