Case Report

A Timely Intervention: Endoscopic Retrieval of a Swallowed Magnetized Activity Watch

Jason S. Radowsky, Joseph S. Lee, and Andrew T. Schlussel

Department of General Surgery, Brian Allgood Army Community Hospital, Yongsan 96205, Republic of Korea

Correspondence should be addressed to Andrew T. Schlussel; andrew.t.schlussel@gmail.com

Received 12 October 2015; Revised 24 December 2015; Accepted 29 December 2015

Academic Editor: Yoshihiro Moriwaki

Copyright © 2016 Jason S. Radowsky et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The accidental ingestion of a foreign object often presents a difficult scenario for the clinician. This includes not only the decision to retrieve the material but also the appropriate technique to use. We present the case of a young asymptomatic girl who swallowed a magnetic activity watch, which was then successfully retrieved with an endoscopic snare. To our knowledge, this is the first documented case of salvaging an operational watch from the stomach using an endoscopic technique.

1. Introduction

Foreign body ingestion is a challenging medical dilemma for both surgeons and gastroenterologists, with the greatest incidence seen in the psychiatric, elderly, and pediatric populations. The mechanism of ingestion varies based on the cohort studied. The psychiatric population may have an altered decision-making capability or will swallow objects for a secondary gain [1]. The elderly often have altered mental status, decreased oropharyngeal sensation and control, and ill-fitting dental appliances. The vast majority of cases, > 80%, occur in children. However, only 10–20% require endoscopic removal, with < 1% mandating an operation to remove the foreign body [2, 3]. Younger children demonstrate the greatest risk as their senses are developing and begin to explore their own environment, frequently placing objects in their mouth [4]. The object ingested, anatomical location, and age of the patient are important factors when making the clinical decision to pursue extraction through an invasive technique or to manage the patient expectantly in hopes of transit through the gastrointestinal tract [5]. An endoscopic or operative approach to extract dentures, batteries, or magnets is strongly recommended, as these objects portend a greater risk of intestinal laceration, perforation, obstruction, or luminal necrosis [4, 6, 7].

As one may expect, there is a paucity of data in the literature regarding the extraction of an ingested watch or

object with similar dimensions. One case report has discussed the extraction of a watch through an open gastrostomy, three cases have utilized an endoscopic approach for retrieval of an impacted watch in the proximal esophagus, and one report discussed successive esophageal balloon dilation, to dislodge the timepiece, pushing it distally into the stomach to allow for passage [8–11]. To our knowledge this is the first documented case of a watch extraction from the stomach using a flexible endoscope.

2. Case Report

This is a case of a healthy, well-developed, 13-year-old girl who was reported to have accidentally swallowed a Shine Misfit fitness tracking watch (Misfit Wearables, Burlingame, CA, USA). While swimming, she removed the magnetized functional portion of the watch from its band and placed this in her mouth at which time it was swallowed by mistake. At the time of presentation, the object was identified in the stomach on an emergency department X-ray at our community-sized hospital. The patient was initially managed conservatively and was informed to return the following day for serial imaging (Figure 1). After approximately 30 hours, the watch was still retained in the stomach. Due to the failure of distal progression through the pylorus and concern for obstruction at the ileocecal valve, the decision was made to proceed with an endoscopic intervention in the operating

FIGURE 3: Successful end-on lassoing in the groove of the watch with the snare.

consent for the use of the images and information in this case report.

3. Discussion

The accidental ingestion of a foreign object is far more common in pediatric, elderly, psychiatric, and incarcerated patients when compared to the adolescent or healthy adult population. Once in the stomach, 80% of these objects will eventually pass out of the body; therefore, intervention remains uncommon as long as the patient is asymptomatic [13]. The decision to proceed with endoscopic retrieval, in this case, was prompted by a failure in the progression of the watch for greater than 24 hours. In addition, there was an uncertainty to the maintenance of the watch's structural integrity when exposed to a low pH for an extended period of time. The device is advertised as being waterproof to 50 meters; however, this did not allay the surgeon or patients' concern for battery exposure and the risk of a caustic perforation [12]. Several studies have shown that sharp foreign bodies, severe symptoms, long duration from ingestion to endoscopy, and existence of mucosal injury are significant risk factors predictive of complications related to removal of foreign bodies [14, 15]. We believe that early detection and a timely endoscopic removal were two factors that contributed to successful retrieval for this case.

As this retrieval occurred at a small community hospital, a limited variety of endoscopic tools were available. While there are no published guidelines for which endoscopic snare or grasper to utilize during foreign body recovery, the American Society for Gastrointestinal Endoscopy (ASGE) provides recommendations regarding the dimensions, configurations, and indications for many commercially available devices [16]. In addition, the ASGE has published intervention timing guidelines based on patient symptoms, object type and size, and anatomic location. Although a magnetized digital watch is not included in these published categories, disk batteries and coins larger than 2 cm are recommended to be extracted 1-2 days following ingestion [17].

In conclusion, the retrieval of a magnetized watch from the stomach with a flexible endoscope is a safe and efficacious procedure. It is critical that the physician must consider

patient size, as well as the device size and design, to formulate an appropriate management plan. The patient should understand the endoscopic and operative risks and should be counseled on the need for a timely intervention when observation has failed.

Copyright Protection

Our team is comprised of military service members and employees of the US Government. This work was prepared as part of our official duties. Title 17 U.S.C. 105 provides that "Copyright protection under this title is not available for any work of the United States Government." Title 17 U.S.C. 101 defines a US Government work as a work prepared by a military service member or employee of the US Government as part of that person's official duties.

Disclaimer

This is an original work by the authors. The opinions expressed are the authors' and the authors' alone. They do not necessarily reflect the opinion of the United States Government, the United States Department of Defense, or Brian Allgood Army Community Hospital.

Conflict of Interests

The authors declare no conflict of interests regarding the publication of this paper.

Authors' Contribution

Conception and design were carried out by Jason S. Radowsky. Creation of media was done by Andrew T. Schlusell. Drafting of the paper was carried out by Jason S. Radowsky. Critical revision was done by Joseph S. Lee and Andrew T. Schlusell. Supervision was done by Andrew T. Schlusell.

Acknowledgment

The authors would like to gratefully thank SGT Sky S. Park, LPN, who acted as their endoscopic technician and who contributed his experience and assistance to safely and efficiently complete this intervention.

References

- [1] I. C. Grimes, B. J. Spier, L. R. Swize, M. J. Lindstrom, and P. R. Pfau, "Predictors of recurrent ingestion of gastrointestinal foreign bodies," *Canadian Journal of Gastroenterology*, vol. 27, no. 1, pp. e1-e4, 2013.
- [2] D. C. Little, S. R. Shah, S. D. St Peter et al., "Esophageal foreign bodies in the pediatric population: our first 500 cases," *Journal of Pediatric Surgery*, vol. 41, no. 5, pp. 914-918, 2006.
- [3] W. Cheng and P. K. H. Tam, "Foreign body ingestion in children: experience with 1,265 cases," *Journal of Pediatric Surgery*, vol. 34, no. 10, pp. 1472-1476, 1999.

- [4] S. J. Sharpe, L. M. Rochette, and G. A. Smith, "Pediatric battery-related emergency department visits in the United States, 1990–2009," *Pediatrics*, vol. 129, no. 6, pp. 1111–1117, 2012.
- [5] I. Rahman, P. Patel, P. Boger et al., "Therapeutic upper gastrointestinal tract endoscopy in Pediatric Gastroenterology," *The World Journal of Gastrointestinal Endoscopy*, vol. 7, no. 3, pp. 169–182, 2015.
- [6] S. N. Bandyopadhyay, S. Das, S. K. Das, and A. Mandal, "Impacted dentures in the oesophagus," *Journal of Laryngology and Otology*, vol. 128, no. 5, pp. 468–474, 2014.
- [7] M. Strickland, D. Rosenfield, and A. Fecteau, "Magnetic foreign body injuries: a large pediatric hospital experience," *Journal of Pediatrics*, vol. 165, no. 2, pp. 332–335, 2014.
- [8] E. B. Dubois, "Watch in the stomach," *Journal of the American Medical Association*, vol. 245, no. 17, p. 1731, 1981.
- [9] A. Almaç and M. Gönüllü, "Wrist watch in the esophagus," *Journal of Otolaryngology*, vol. 17, no. 4, p. 200, 1988.
- [10] N. Aggarwal and D. Sinha, "An amazing case of working wrist watch in the esophagus," *Indian Journal of Otolaryngology and Head and Neck Surgery*, vol. 58, no. 1, pp. 105–106, 2006.
- [11] Y. T. Jeon, J. Chun, C. W. Song et al., "Endoscopic removal of sharp foreign bodies impacted in the esophagus," *Endoscopy*, vol. 33, no. 6, pp. 518–522, 2001.
- [12] Misfit Wearables, Misfit Shine, March 2015, <http://misfit.com/products/shine>.
- [13] V. Selivanov, G. F. Sheldon, J. P. Cello, and R. A. Crass, "Management of foreign body ingestion," *Annals of Surgery*, vol. 199, no. 2, pp. 187–191, 1984.
- [14] Y.-K. Park, K.-O. Kim, J.-H. Yang, S.-H. Lee, and B.-I. Jang, "Factors associated with development of complications after endoscopic foreign body removal," *Saudi Journal of Gastroenterology*, vol. 19, no. 5, pp. 230–234, 2013.
- [15] J. H. Park, C. H. Park, J. H. Park et al., "Review of 209 cases of foreign bodies in the upper gastrointestinal tract and clinical factors of successful endoscopic removal," *The Korean Journal of Gastroenterology*, vol. 43, no. 4, pp. 226–233, 2004.
- [16] D. L. Diehl, D. G. Adler, J. D. Conway et al., "Endoscopic retrieval devices," *Gastrointestinal Endoscopy*, vol. 69, no. 6, pp. 997–1003, 2009.
- [17] S. O. Ikenberry, T. L. Jue, M. A. Anderson et al., "Management of ingested foreign bodies and food impactions," *Gastrointestinal Endoscopy*, vol. 73, no. 6, pp. 1085–1091, 2011.