Case Report

First Case of Lung Abscess due to *Salmonella enterica* Serovar Abony in an Immunocompetent Adult Patient

Vassiliki Pitiriga,¹ John Dendrinis,² Emanuel Nikitiadis,²
Georgia Vrioni,¹ and Athanassios Tsakris¹

¹Department of Microbiology, Medical School, National and Kapodistrian University of Athens, M. Asias 75, Goudi, 11527 Athens, Greece

²Metropolitan Hospital, Ethnarchou Makariou 9 & El. Venizelou 1, N. Faliro, 18547 Athens, Greece

Correspondence should be addressed to Vassiliki Pitiriga; siliapit@hotmail.com

Received 19 March 2016; Accepted 8 June 2016

Academic Editor: Lawrence Yamuah

Copyright © 2016 Vassiliki Pitiriga et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In healthy individuals, nontyphoidal *Salmonella* species predominantly cause a self-limited form of gastroenteritis, while they infrequently invade or cause fatal disease. Extraintestinal manifestations of nontyphoidal *Salmonella* infections are not common and mainly occur among individuals with specific risk factors; among them, focal lung infection is a rare complication caused by nontyphoidal *Salmonella* strains typically occurring in immunocompromised patients with prior lung disease. We describe the first case of a localized lung abscess formation in an immunocompetent healthy female adult due to *Salmonella enterica* serovar Abony. The patient underwent lobectomy and was discharged after full clinical recovery. This case report highlights nontyphoidal *Salmonellae* infections as a potential causative agent of pleuropulmonary infections even in immunocompetent healthy adults.

1. Introduction

In developed countries, nontyphoidal *Salmonellae* (NTS) strains are a leading cause of self-limiting enterocolitis in healthy population; they are estimated to cause 94 million cases of gastroenteritis and 115,000 deaths globally each year [1]. Up to 5% of patients will develop secondary bacteremia [2], with low attributable mortality (1–5%). Localized extraintestinal infections develop as secondary complications in approximately 5–10% of cases with NTS bacteremia [3] and occur predominately in a wide variety of immunocompromised individuals [4], including patients with severe underlying diseases [5], immunocompromised elderly patients [6, 7], or children [8]. Mainly they involve the gastrointestinal tract, endothelial surfaces [9], pericardium [10], meninges [11], lungs, joints, and bones [12], or soft tissues. Among them, pleuropulmonary NTS infection is an infrequent manifestation [13, 14] mostly occurring in immunocompromised patients with prior lung or pleural pathology [15].

In healthy immunocompetent individuals, extraintestinal complications caused by NTS remain uncommon [16–19]. We

present the first case of a lung abscess caused by *Salmonella* serovar Abony in an immunocompetent healthy young adult with no prior history of pulmonary disease or presence of any underlying disease.

2. Case Presentation

A 26-year-old female of Hellenic ethnicity was admitted to our outpatient clinic, reporting a 10-day history of low grade fever, chills, nausea, headache, urge to vomiting, and a dull pain over the right kidney area. Her past medical history was free of any chronic or acute infection or systemic disease, except for a prior admission to our hospital 1 year ago due to diarrhea not associated with any specific microorganism.

Her physical examination revealed temperature of 38°C, normal blood pressure (110/70 mmHg), and tachycardia (HR = 160 bpm). During palpation, a slight pain in right upper abdominal area was revealed with no other specific signs and symptoms. The respiratory examination was remarkable for dullness to percussion with decreased breath sounds over the lower right lung base. The remainder of the physical

FIGURE 1: (a) Chest radiograph on presentation; (b) chest CT scanning on day 3 of admission, showing a large abscess in the right lower lobe.

examination was unremarkable. Initial laboratory findings revealed a polymorphonuclear leucocytosis of $13,700/\text{mm}^3$ with 72% neutrophils and 18% lymphocytes, ESR of 29 mm/hr, and C-reactive protein of 5.31 mg/dL. No other pathological findings were indicated from the biochemical testing. Detailed investigations did not reveal any predisposing factors or evidence of an underlying immunodeficiency. More specifically, there was no evidence of malnutrition, no history of therapy with glucocorticoids or other immunosuppressive drugs, and no indication of immunoglobulin excess or deficiency through quantitative serum immunoglobulin tests, and blood tests were negative for chronic infections (HIV, viral hepatitis, etc.) or autoimmune disorders (antinuclear antibodies and other autoantibodies). Chest X-ray examination showed pneumonic infiltration in the lower one-third of the right hemithorax and laterally located dense appearance resembling left pleural effusion. Based on this evidence, the patient was diagnosed as having community-acquired pneumonia and after blood cultures were taken, antimicrobial treatment was initiated with intravenous ceftriaxone 1 gr/day + azithromycin 500 mg/day. Sputum, protected specimen brush (PSB) material of bronchial secretions, and three sets of blood specimens were also taken on admission for cultures which did not yield any pathogens.

Within the following two days, the patient's fever rose to 40°C despite the administration of antimicrobial therapy, and her condition deteriorated by developing dry cough, chest pain, total absence of breath sound during auscultation in right hemithorax, and dyspnea. Additionally, a strong right lumbar pain appeared. Three additional sets of blood cultures, taken while the patient was febrile, were negative.

On the third day, computed tomography (CT) scanning of the chest was performed and revealed a lung thick-walled abscess formation in the right lower lobe, with a surrounding inflammatory infiltrate, extended atelectasis, and pleural

effusion to the right lower lobe. Figures 1(a) and 1(b) exhibit the size and morphology of the lung abscess. A subsequent ultrasound examination of the patient's liver, carried out in order to examine whether there was any subdiaphragmatic extension or origin of the infection, did not reveal any relevant evidence.

The combination of ceftriaxone + azithromycin was consequently discontinued and replaced by moxifloxacin 400 mg/day + tazobactam plus piperacillin (0.5 + 4.0) gr $\times$ 3/day + clindamycin 600 mg $\times$ 3/day. PCR for tuberculosis was performed and anti-*Echinococcus* IgG and IgM antibody titers were measured; however all the results were negative.

The next days, her fever did not subside, CRP levels rose gradually to 16.85 mg/dL, and the pleural effusion continued to rise. Blood cultures remained all negative for any bacterial growth.

On the sixth day of admission, the patient underwent lobectomy, owing to the lack of response to antibiotic therapy, the deterioration of symptoms, and the difficulty in approaching the specific lobe area by thoracentesis. Approximately 700 mL of pleural fluid was collected and sent to laboratory for biochemical analysis, Gram stain, cultures, and antimicrobial profile. Biochemical analysis of the pleural fluid showed the following: glucose of 74 mg/dL, lactate dehydrogenase of 730 U/L, total protein of 3.9 g/dL, and white blood cell count of $15,400/\text{mm}^3$ with 80% polymorphs. Gram staining and cultures of the pleural fluid were negative. In addition, the aspiration of the abscess revealed yellowish pus (about 45 mL) that was also sent for laboratory analysis the same day. Cultures of the pus sample collected from the abscess yielded a Gram-negative aerobic rod identified as *Salmonella enterica* subsp. *enterica* serovar Abony. *Salmonella* isolate was identified to the genus level by both the automated Vitek-2 System (bioMérieux, Inc., Hazelwood, MO) and the API 20E (bioMérieux, Inc., Hazelwood, MO).

Serotyping of the isolate was performed using the somatic O and flagellar H antisera according to the Kauffman-White classification scheme (Difco Laboratories, Detroit, MI, USA). Molecular confirmation of *Salmonella* serotyping was carried out using the DNA microarray system Premi-Test® *Salmonella* (DSM Nutritional Products, Check-Points, Wageningen, Netherlands) [20]. Antimicrobial susceptibility testing was initially performed by the Vitek-2 System, according to the recommendations of the National Committee for Clinical Laboratory Standards [21] and confirmed by E-test (bioMérieux, Inc., Hazelwood, MO). The isolate was susceptible to commonly used antibiotics (ampicillin MIC of 0.75 µg/mL, ceftriaxone MIC of 0.085 µg/mL, cefotaxime MIC of 0.082 µg/mL, ceftazidime MIC of 0.115 µg/mL, ciprofloxacin MIC of 0.032 µg/mL, moxifloxacin MIC of 0.016 µg/mL, and trimethoprim-sulfamethoxazole MIC of 0.064 µg/mL). Based on the laboratory report, the antimicrobial therapy was changed on the eighth day after admission, to sulfamethoxazole/trimethoprim (800 + 160) mg × 2/day, moxifloxacin 400 mg/day, and clindamycin 600 mg × 3/day, along with supportive therapy.

The postoperative clinical condition of the patient improved noticeably. Six days after surgery, the patient's symptoms resolved and she was discharged on sulfamethoxazole/trimethoprim (800 + 160) mg × 2/day and ciprofloxacin 500 mg × 2/day, for 20 days.

3. Discussion

Even though the prevalence of invasive NTS in humans by means of bacteremia and extraintestinal infections is increasing worldwide among immunocompromised patients, particularly in developing countries, likely secondary to the high prevalence of coexisting malnutrition, malaria, and HIV infection, it remains uncommon in immunocompetent subjects [22]. Especially for NTS pleuropulmonary infection as a secondary manifestation in healthy individuals, only two reports exist in the literature documenting *Salmonella* Group B spp. as the primary cause of lung abscess in two immunocompetent female children [23, 24]. In another report of Genzen et al. [25], even though the 55-year-old man diagnosed with pulmonary *Salmonella* serovar Typhimurium infection was considered as immunocompetent, his medical history of chronic alcoholism and bronchitis should be taken into account as a significant predisposing factor for the invasive development of the infection. To our knowledge, this is the first report of lung abscess caused by NTS presenting in an immunocompetent healthy individual of the adult age group. Moreover, only one case report has been described in the literature documenting *Salmonella* serovar Abony to cause severe invasive disease, by means of disseminated intravascular coagulation, in an immunocompromised elderly patient [26]. We report for the first time an extraintestinal complication of lung abscess caused by *Salmonella* serovar Abony.

Lung infections by NTS may occur via several routes such as direct extension from a nearby infection, aspiration of gastric secretions, or hematogenous dissemination from the gastrointestinal tract. In the present case, our patient reported a history of diarrhea one year before the onset of

illness. At that time, she was admitted to our clinic with a history of loose stools for a period of the prior ten days. She was given treatment with ciprofloxacin for three days before admission. On examination, she was not febrile and the stools were not accompanied by mucus or blood, while vomiting was absent. Stool and blood cultures were negative, possibly due to the antibiotic treatment and she was discharged 4 days after without identification of the infectious causative agent. This incident might be connected with the existing pulmonary complication, as it provides an indication of a potential gastrointestinal infection by *Salmonella* Abony with a subsequent seeding of the pathogen to the lungs through bacteremia.

4. Conclusion

This case report indicates that NTS strains should be considered as a potential etiological agent of infection in the differential diagnosis of pleuropulmonary infection causes, even among immunocompetent healthy adults.

Competing Interests

The authors declare that they have no conflict of interests relevant to this paper.

References

- [1] S. E. Majowicz, J. Musto, E. Scallan et al., "The global burden of nontyphoidal *Salmonella* gastroenteritis," *Clinical Infectious Diseases*, vol. 50, no. 6, pp. 882–889, 2010.
- [2] E. L. Hohmann, "Nontyphoidal salmonellosis," *Clinical Infectious Diseases*, vol. 32, no. 2, pp. 263–269, 2001.
- [3] D. A. Pegues, M. E. Ohl, and S. I. Miller, "*Salmonella* species, including *Salmonella typhi*," in *Principles and Practice of Infectious Diseases*, G. L. Mandell, J. E. Bennett, and R. Dolin, Eds., pp. 2636–2654, Churchill Livingstone, Philadelphia, Pa, USA, 6th edition, 2005.
- [4] M. A. Gordon, "*Salmonella* infections in immunocompromised adults," *Journal of Infection*, vol. 56, no. 6, pp. 413–422, 2008.
- [5] B. Cabaret, M.-L. Couëc, M. Lorrot, E. Launay, and C. Gras-Le Guen, "Multifocal osteoarticular infection caused by *Salmonella* non typhi in a child with sickle cell disease," *Archives de Pédiatrie*, vol. 20, no. 4, pp. 398–402, 2013.
- [6] P.-L. Chen, C.-M. Chang, C.-J. Wu et al., "Extraintestinal focal infections in adults with nontyphoid *Salmonella* bacteraemia: predisposing factors and clinical outcome," *Journal of Internal Medicine*, vol. 261, no. 1, pp. 91–100, 2007.
- [7] R.-B. Hsu, Y.-G. Tsay, R. J. Chen, and S.-H. Chu, "Risk factors for primary bacteremia and endovascular infection in patients without acquired immunodeficiency syndrome who have nontyphoid salmonellosis," *Clinical Infectious Diseases*, vol. 36, no. 7, pp. 829–834, 2003.
- [8] E. Çiftçi, H. Güriz, A. D. Aysev, E. Ince, B. Erdem, and U. Doğru, "*Salmonella* bacteraemia in Turkish children: 37 cases seen in a university hospital between 1993 and 2002," *Annals of Tropical Paediatrics*, vol. 24, no. 1, pp. 75–80, 2004.
- [9] R.-B. Hsu and F.-Y. Lin, "Risk factors for bacteraemia and endovascular infection due to non-typhoid *Salmonella*: a reappraisal," *Quarterly Journal of Medicine*, vol. 98, no. 11, pp. 821–827, 2005.

- [10] D. Ortiz, E. M. Siegal, C. Kramer, B. K. Khandheria, and E. Brauer, "Nontyphoidal cardiac salmonellosis: two case reports and a review of the literature," *Texas Heart Institute Journal*, vol. 41, no. 4, pp. 401–406, 2014.
- [11] E. M. Molyneux, L. A. Mankhambo, P. Ajib et al., "The outcome of non-typhoidal *Salmonella* meningitis in Malawian children, 1997–2006," *Annals of Tropical Paediatrics*, vol. 29, no. 1, pp. 13–22, 2009.
- [12] A. Dutta and C. H. Allen, "Non-typhoidal *Salmonella* osteomyelitis in the midfoot of a healthy child and review of the literature," *Journal of Infectious Diseases and Therapy*, vol. 1, no. 2, article 107, 2013.
- [13] P.-L. Chen, C.-J. Wu, C.-M. Chang et al., "Extraintestinal focal infections in adults with *Salmonella enterica* serotype Choleraesuis bacteremia," *Journal of Microbiology, Immunology and Infection*, vol. 40, no. 3, pp. 240–247, 2007.
- [14] E. Moraitou, I. Karydis, D. Nikita, and M. E. Falagas, "Case report: parotid abscess due to *Salmonella enterica* serovar Enteritidis in an immunocompetent adult," *International Journal of Medical Microbiology*, vol. 297, no. 2, pp. 123–126, 2007.
- [15] C. Saturveithan, "Salmonella osteomyelitis in a one year old child without sickle cell disease: a case report," *Malaysian Orthopaedic Journal*, vol. 8, pp. 52–54, 2014.
- [16] J. C. Kam, S. Abdul-Jawad, C. Modi et al., "Pleural Empyema due to Group D Salmonella," *Case Reports in Gastrointestinal Medicine*, vol. 2012, Article ID 524561, 4 pages, 2012.
- [17] N. F. Crum, "Non-typhi *Salmonella* empyema: case report and review of the literature," *Scandinavian Journal of Infectious Diseases*, vol. 37, no. 11-12, pp. 852–857, 2005.
- [18] R. L. Hall, R. Partridge, N. Venkatraman, and M. Wiselka, "Invasive non-typhoidal *Salmonella* infection with multifocal seeding in an immunocompetent host: an emerging disease in the developed world," *BMJ Case Reports*, vol. 2013, 2013.
- [19] W.-J. Lin, C.-C. Wang, S.-N. Cheng, W.-T. Lo, and C.-H. Hung, "Hand abscess, phlebitis, and bacteremia due to *Salmonella enterica* serotype Augustenborg," *Journal of Microbiology, Immunology and Infection*, vol. 39, no. 6, pp. 519–522, 2006.
- [20] P. Andreoli, J. Thijssen, R. Anthony, P. Vos, and W. De Levita, "Fast method for detecting micro-organisms in food samples 2004," Patent WO2004106547 A3.
- [21] National Committee for Clinical Laboratory Standards, "Performance standards for antimicrobial susceptibility testing; 24th informational supplement," Tech. Rep. M100-S11, National Committee for Clinical Laboratory Standards, Wayne, Pa, USA, 2014.
- [22] E. Galanakis, M. Bitsori, S. Maraki, C. Giannakopoulou, G. Samonis, and Y. Tselentis, "Invasive non-typhoidal salmonellosis in immunocompetent infants and children," *International Journal of Infectious Diseases*, vol. 11, no. 1, pp. 36–39, 2007.
- [23] R. Thapa, A. Ghosh, D. Mallick, and B. Biswas, "Lung abscess secondary to non-typhoidal salmonellosis in an immunocompetent girl," *Singapore Medical Journal*, vol. 50, no. 10, pp. 1033–1035, 2009.
- [24] B. Adhisivam, S. Mahadevan, and C. Udaykumar, "Lung abscess caused by *Salmonella*," *Indian Journal of Pediatrics*, vol. 73, no. 5, pp. 450–451, 2006.
- [25] J. R. Genzen, D. M. Towle, J. D. Kravetz, and S. M. Campbell, "*Salmonella typhimurium* pulmonary infection in an immunocompetent patient," *Connecticut Medicine*, vol. 72, no. 3, pp. 139–142, 2008.
- [26] S. C. Glover, C. C. Smith, and I. A. Porter, "Fatal *Salmonella* septicaemia with disseminated intravascular coagulation and renal failure," *Journal of Medical Microbiology*, vol. 15, no. 1, pp. 117–121, 1982.