

Case Report

Catheter-Related ESBL-Producing *Leclercia adecarboxylata* Septicemia in Hemodialysis Patient: An Emerging Pathogen?

Roa Sultana Alosaimi ¹ and Mai Muhmmmed kaaki²

¹Department of Medicine, Ministry of Health, Jeddah, Saudi Arabia

²Department of Microbiology, King Abdul-Aziz Medical City, Jeddah, Saudi Arabia

Correspondence should be addressed to Roaa Sultana Alosaimi; dr.roaa@live.com

Received 19 October 2019; Accepted 6 January 2020; Published 22 January 2020

Academic Editor: Raul Colodner

Copyright © 2020 Roaa Sultana Alosaimi and Mai Muhmmmed kaaki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We report a multidrug-resistant strain of *Leclercia adecarboxylata* which was responsible for a catheter-related bacteremia, in a 50-year-old female with an end-stage renal disease on hemodialysis. The isolated strain was an extended beta-lactamase producer. Based on a literature review of *L. adecarboxylata*, there have been only two reports of extended beta-lactamase producer strains. To our knowledge, this is the first case reported in Saudi Arabia.

1. Introduction

Leclercia adecarboxylata is a motile, Gram-negative, oxidase-negative bacterium, which was first described by Leclerc in 1962 as *Escherichia adecarboxylata* [1]. Based on nucleic acid and protein electrophoretic techniques, “*E. adecarboxylata*” was separated from the “*Enterobacter agglomerans*” complex [2] to which it was assigned temporarily and renamed *Leclercia adecarboxylata* [3]. This bacterium was isolated from environmental samples, including water and soil [4]. However, a cumulative report of human infections by *L. adecarboxylata* mostly in immunocompromised patients suggested that the organism is an opportunistic bacterium and the infections caused by it are underestimated. It has been reported to cause septicemia [5], wound infection [6, 7], urinary tract infection [8, 9], posttraumatic polymicrobial infection [10], soft tissue infection [11], and peritonitis [12]. The majority of clinical isolates of *L. adecarboxylata* were susceptible to commonly used antibiotics [4], and multidrug-resistant isolates have rarely been reported [8]. Here, we report a case of extended-spectrum beta-lactamase-producing *L. adecarboxylata* from a patient with catheter-related septicemia.

2. Case Report

A 50-year-old female patient with a known case of diabetes mellitus and end-stage renal disease due to hypertension was started on hemodialysis in March 2014. Permanent tunneled catheters were placed into the right internal jugular vein as the patient was not fit for AVF creation due to vascular occlusions.

On 7 April 2019, at the end of a dialysis session, the patient developed fever (39°C) and chills. Fever was documented at 39°C, was associated with chills and rigors, and was not responding to antipyretics. In the emergency department, the patient reported no change in weight, loss of appetite, or night sweats. She denied any history of respiratory, gastrointestinal, neurological symptoms. On examination, the temperature was 39°C, the blood pressure 150/75 mm Hg, the heart rate 110 beats per minute, and the respiratory rate 18 breaths per minute. The patient was fully conscious and oriented. The remainder of the general medical examination was normal.

She was admitted to our nephrology unit with possible source catheter-related blood stream infection. She was started on ceftazidime 2 gm IV once daily and vancomycin 1 gm IV Q HD as per the protocol. The white blood cell

TABLE 1: *Leclercia adecarboxylata* antibiotic susceptibility profile.

Antimicrobial agents	MIC (ug/ml)*	Susceptibility
Amoxicillin/clavulanate	4	R
Ampicillin	≥32	R
Ceftriaxone	≥64	R
Ciprofloxacin	≤0.25	S
Gentamicin	≤1	S
Amikacin	≤2	S
Meropenem	≤0.25	S
Trimethoprim/sulfamethoxazole	≤20	S
Piperacillin/tazobactam	≤4	R

MIC = minimum inhibitory concentration; S = sensitive; R = resistant.
*Vitek 2 automatic system using an AST GN card.

count was 13.3×10^3 cells, C-reactive protein was 278 mg/dl, and procalcitonin was 52.27. The electrocardiogram revealed sinus tachycardia, and chest radiography was negative.

Blood culture from venous and central catheters was sampled separately, and the causative microorganism was identified as a Gram-negative, motile, lactose-fermenting bacillus, which was identified by the Vitek MS automatic identification system using a GN card as *Leclercia adecarboxylata*. Antibiotic sensitivity testing using Vitek 2 showed that it is an extended-spectrum beta-lactamase producer. The results are summarized in Table 1 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3421816/table/T1/>). The *L. adecarboxylata* strain was resistant to most beta-lactams, including narrow, expanded, and broad-spectrum cephalosporins. But it was susceptible to all quinolones and carbapenems tested.

The treatment regimen was changed to meropenem 500 mg IV once daily and gentamicin based on antibiotic susceptibility tests (ASTs) from blood cultures. This targeted therapy was successful, and the patient became afebrile. The patient received meropenem and gentamicin for two weeks, and then she was discharged in good condition.

3. Discussion

L. adecarboxylata is a Gram-negative bacillus belonging to the Enterobacteriaceae family. The association of *L. adecarboxylata* with catheter (particularly tunneled CVC) related septicemia is increasingly reported [13]. In 2013, De Mauri and colleagues reported that there were 15 reports of adult patients with catheter-related septicemia due to *L. adecarboxylata* [5]. Bacterial resistance to antibiotics is increasing worldwide in healthcare settings and in the community. The dissemination of extended-spectrum beta-lactamase Enterobacteriaceae (ESBL-E) is alarming [14]. Antibiotic-resistant *L. adecarboxylata* strains have been reported in six different cases. Of these cases, only two were extended beta-lactamase producer isolates. The first case from a patient with acute myeloid leukaemia produced ESBL. This strain encoded SHV-type beta-lactamases [15]. The second case was ESBL producing a multidrug-resistant *L. adecarboxylata* strain harbouring blaTEM-1, blaCTX-M-3, and int11 cassette (dfrA12-orfF-aadA2) genes in a 47-year-old female with breast cancer [16]. Although *L. adecarboxylata* has been recognized as a relatively insignificant

human pathogen due to its low virulence and high antibiotic susceptibility, multidrug-resistant strains can become life-threatening human bacterial pathogens by acquiring genetic determinants, including blaSHV, blaTEM-1, blaCTX-M group 1, and int11 genes. Of note, *L. adecarboxylata* is an uncommon isolate in microbiology laboratories, but it will likely be more often identified with new diagnostic techniques.

In conclusion, we report the third case of catheter-related septicemia due to ESBL producing a multidrug-resistant *L. adecarboxylata* strain in a 50-year-old female with end-stage renal disease.

Data Availability

The authors declare that this article does not show the name or data of the patient.

Consent

Written informed consent was obtained from the patient.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] H. Leclerc, "Biochemical study of pigmented Enterobacteriaceae," *Annales de l'Institut Pasteur*, vol. 102, pp. 726–741, 1962.
- [2] D. Izard, J. Mergaert, F. Gavini et al., "Separation of *Escherichia adecarboxylata* from the 'Erwinia herbicola-Enterobacter agglomerans' complex and from the other Enterobacteriaceae by nucleic acid and protein electrophoretic techniques," *Annales de l'Institut Pasteur*, vol. 136, pp. 151–168, 1985.
- [3] K. Tamura, R. Sakazaki, Y. Kosako, and E. Yoshizaki, "Leclercia adecarboxylata Gen. Nov., Comb. Nov., formerly known as *Escherichia adecarboxylata*," *Current Microbiology*, vol. 13, no. 4, pp. 179–184, 1986.
- [4] I. Stock, S. Burak, and B. Wiedemann, "Natural antimicrobial susceptibility patterns and biochemical profiles of *Leclercia adecarboxylata* strains," *Clinical Microbiology and Infection*, vol. 10, no. 8, pp. 724–733, 2004.
- [5] A. De Mauri, D. Chiarinotti, S. Andreoni, G. L. Molinari, N. Conti, and M. De Leo, "Leclercia adecarboxylata and catheter-related bacteraemia: review of the literature and outcome with regard to catheters and patients," *Journal of Medical Microbiology*, vol. 62, no. 10, pp. 1620–1623, 2013.
- [6] E. H. Hurley, E. Cohen, J. A. Katarincic, and R. K. Ohnmacht, "Leclercia Adecarboxylata Infection in an immunocompetent child," *Rhode Island Medical Journal*, vol. 98, no. 9, pp. 41–44, 2015.
- [7] Z. Temesgen, D. R. Toal, and F. R. Cockerill III, "Leclercia adecarboxylata Infections: case report and review," *Clinical Infectious Diseases*, vol. 25, no. 1, pp. 79–81, 1997.
- [8] A. Mkanera, M. Conde, M. A. Diallo et al., "A multi-drug resistance pattern of a," *International Journal of Biological and Chemical Sciences*, vol. 12, no. 4, p. 1550, 2018.
- [9] H. Sawamura, Y. Kawamura, M. Yasuda et al., "A clinical isolate of *Leclercia adecarboxylata* from a patient of

- pyelonephritis,” *Journal of the Japanese Association for Infectious Diseases*, vol. 79, no. 10, pp. 831–835, 2014.
- [10] J. D. Forrester, J. Adams, and R. G. Sawyer, “*Leclercia adedecarboxylata* bacteremia in a trauma patient: case report and review of the literature,” *Surgical Infections*, vol. 13, no. 1, pp. 63–66, 2012.
- [11] C. A. Botero-García, C. H. Gómez, J. S. Bravo, and L. A. Pescador Vargas, “*Leclercia adedecarboxylata*, a rare cause of soft tissue infections in immunocompromised patients, case report and review of the literatura,” *Infectio*, vol. 22, no. 4, p. 223, 2018.
- [12] H. M. Kim, C. Y. Chon, S. H. Ahn et al., “Fatal spontaneous bacterial peritonitis by *Leclercia adedecarboxylata* in a patient with hepatocellular carcinoma,” *International Journal of Clinical Practice*, vol. 62, no. 8, pp. 1296–1298, 2008.
- [13] M. F. Fernandez-Ruiz, L. Lopez-Medrano, A. Garcia-Sanchez et al., “Successful management of tunneled hemodialysis catheter-related bacteremia by *Leclercia adedecarboxylata* 166 without catheter removal: report of two cases,” *International Journal of Infectious Diseases*, vol. 13, pp. e517–e518, 2009.
- [14] J. D. Pitout and K. B. Laupland, “Extended-spectrum β -lactamase-producing Enterobacteriaceae: an emerging public-health concern,” *The Lancet Infectious Diseases*, vol. 8, no. 3, pp. 159–166, 2008.
- [15] A. Mazzariol, J. Zuliani, R. Fontana, and G. Cornaglia, “Isolation from blood culture of a *Leclercia adedecarboxylata* strain producing an SHV-12 extended-spectrum beta-lactamase,” *Journal of Clinical Microbiology*, vol. 41, no. 4, pp. 1738–1739, 2003.
- [16] G. W. Shin, M. J. You, H. S. Lee, and C. S. Lee, “Catheter-related bacteremia caused by multidrug-resistant *Leclercia adedecarboxylata* in a patient with breast cancer,” *Journal of Clinical Microbiology*, vol. 50, no. 9, pp. 3129–3132, 2012.



Hindawi

Submit your manuscripts at
www.hindawi.com

