Case Report

Idiopathic Acquired Leukonychia in a 34-Year-Old Patient

Maria Rita Bongiorno and Mario Aricò

Department of Dermatology, University of Palermo, Via del Vespro 131, 90127 Palermo, Italy

Correspondence should be addressed to Maria Rita Bongiorno, istderm@unipa.it

Received 27 January 2009; Accepted 5 March 2009

Recommended by Jeffrey Weinberg

We present a rare case of a 34-year-old patient with persistent, progressive, acquired leukonychia totalis and partialis. Idiopathic acquired leukonychia is a rare chromatic disorder of the nail not associated with other abnormalities and discernible etiology. Our case report did not link the inheritance of leukonychia with diverse clinical syndromes. To our knowledge, only five cases of idiopathic, acquired, true total leukonychia were found in literature. This case was the sixth patient with asymptomatic idiopathic, white fingernails, and toenails without a hereditary cause.

Copyright © 2009 M. R. Bongiorno and M. Aricò. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Leukonychia refers to whiteness of the nails which can occur either in patches or involving the entire nail [1]. The condition can be either true leukonychia, with involvement of the nail plate, or pseudoleukonychia, caused by subungual and nailbed abnormalities.

Leukonychia may be acquired or inherited. Acquired leukonychia is frequently associated with trauma, drugs such as chemotherapeutic agents, [2] systemic or local infections—such as typhoid fever, hepatic cirrhosis, ulcerative colitis or leprosy, hypocalcaemia, and, very commonly, minor trauma. True leukonychia can be inherited as an isolated condition or as one of several other reported syndromes. There are a number of autosomal, dominantly inherited leukonychia syndromes, including leukonychia totalis, in which leukonychia occurs in combination with kidney stone and sebaceous cysts [3, 4], as well as leukonychia, with sensory-neural deafness and knuckle pads, known as the Bart-Pumphrey syndrome [5, 6] (Table 1).

Classification of the so-called true leukonychia may also be based on the distribution of white blotches, known as leukonychia punctata, leukonychia striata, leukonychia partialis, or leukonychia totalis. Idiopathic true leukonychia is a much rarer condition, with only a few reported [7–

11] cases. The following is the case history of a man with persistent, progressive idiopathic leukonychia.

2. Case Presentation

The 34-year-old male patient was admitted in the Department of Dermatology, University of Palermo (Italy), complaining of color changes on the nails. The symptoms were observed synchronously and had apparently been present for 11 years. When the patient was 23, he developed simultaneous leukonychia partialis of both the fingers and toenails. There was no evidence of atopy, lichen planus, alopecia areata, or psoriasis.

The patient was without a previous medical history. He reported not having taken any drug specific and not to be exposed to chemical agents. He also had no peripheral neurovascular disorders. On examination, striata and total leukonychia of all twenty nails was found (Figure 1). Further clinical examination revealed soft nails with slow growth, without other cutaneous or visceral abnormalities.

A progression over the years from leukonychia partialis to leukonychia totalis was demonstrated. On examination, the patient had no central nervous system, eye, ear, hair, teeth, or skin abnormalities.

There was no family history of nail disease, the patient was born to nonconsanguineous parents and had two older

FIGURE 1: (a)-(b) Both hands had opaque nail plates characteristic of leukonychia totalis. (c)-(d) The patient toenails exhibited leukonychia totalis and partialis.

TABLE 1: Classification of true leukonychia.

(A) Hereditary
(1) Isolated
(2) Associated with
(a) Bart-Pumphrey syndrome
(b) Bauer syndrome
(c) Heimler syndrome
(d) Knuckle pads-leukonychia-deafness syndrome
(e) Keratoderma, hypotrichosis and leukonychia totalis syndrome
(f) Lowry-Wood syndrome
(g) FLOTCH syndrome
(h) Congenital keratosis palmaris et plantaris, deafness and total leukonychia
(i) LEOPARD syndrome
(B) Acquired
(1) Idiopathic
(2) Associated with
(a) Trauma
(b) Drugs
(c) Systemic infections
(d) Local infections
(e) Inflammatory disease.

siblings, a brother and sister, without any features of the disease. There was no family history of atopy, psoriasis, lichen

planus, alopecia areata, or any other illnesses. Repeated potassium hydroxide preparations and fungal cultures of the white nails were negative.

3. Discussion

Leukonychia is a whitening of the nail plate. It was first described by Mees in 1919, as an associated finding in arsenic intoxication [12].

The physiologic mechanism leading to this phenomenon is not entirely clear. According to Newton’s theorem, a surface appears white when it reflects the radiation of visible light. This mechanism can be proposed in explaining leukonychia. Because true leukonychia is thought to be due to abnormal matrix keratinization, with persistent parakeratosis and keratohyaline granules in the nail plate, parakeratosis and dissociation of the keratin bundles may play a role in the modification of the solar light reflection by the ungual plates.

Several studies have provided evidence for the association of total leukonychia with diverse clinical syndromes, including leukonychia with palmoplantar keratoderma and atrophic fibrosis, pili torti [13], congenital hypoparathyroidism, hypoparathyroidism, onychorrhexis, and cataracts and the LEOPARD syndrome [1]. Total leukonychia has also been associated with peptic ulcer disease and cholelithiasis as well as with keratoderma and hypotrichosis [14].

Our case report did not link the inheritance of leukonychia with any of the above-mentioned syndromes. To our

TABLE 2: True acquired idiopathic leukonychia.

Authors	Presentations	Active disease (years)
Claudel et al. [8]	Leukonychia totalis and partialis	2
Grossman et al. [1]	Leukonychia partialis to a combined partialis and totalis	3
Stewart et al. [7]	Leukonychia totalis and partialis	Unrelated
Park et al. [10]	Leukonychia partialis to leukonychia totalis	13
Butterworth [11]	Leukonychia totalis and partialis	Unrelated
Our case	Leukonychia partialis to leukonychia totalis	11

knowledge, only five cases of idiopathic, acquired, true total leukonychia were found in literature (Table 2). This case was the sixth patient with asymptomatic idiopathic, white fingernails and toenails without a hereditary cause.

- [14] K. R. Stevens, P. F. Leis, S. Peters, S. Baer, and I. Orengo, "Congenital leukonychia," *Journal of the American Academy of Dermatology*, vol. 39, no. 3, pp. 509–512, 1998.

References

- [1] M. Grossman and R. K. Scher, "Leukonychia: review and classification," *International Journal of Dermatology*, vol. 29, no. 8, pp. 535–541, 1990.
- [2] A. Yoruk and H. Yukselgungor, "Chemotherapy induced transverse leukonychia in children," *International Journal of Dermatology*, vol. 42, no. 6, pp. 468–469, 2003.
- [3] L. L. Bushkell and R. J. Gorlin, "Leukonychia totalis, multiple sebaceous cysts, and renal calculi. A syndrome," *Archives of Dermatology*, vol. 111, no. 7, pp. 899–901, 1975.
- [4] R. S. Medansky and J. M. Fox, "Hereditary leukonychia totalis," *Archives of Dermatology*, vol. 82, no. 3, pp. 412–414, 1960.
- [5] R. S. Bart and R. E. Pumphrey, "Knuckle pads, leukonychia and deafness. A dominantly inherited syndrome," *The New England Journal of Medicine*, vol. 276, no. 4, pp. 202–207, 1967.
- [6] K. R. Ong, S. Visram, S. McKaig, and L. A. Brueton, "Sensorineural deafness, enamel abnormalities and nail abnormalities: a case report of Heimler syndrome in identical twin girls," *European Journal of Medical Genetics*, vol. 49, no. 2, pp. 187–193, 2006.
- [7] L. Stewart, E. Young, and H. W. Lim, "Idiopathic leukonychia totalis and partialis," *Journal of the American Academy of Dermatology*, vol. 13, no. 1, pp. 157–158, 1985.
- [8] C. D. Claudel, J. A. Zic, and A. S. Boyd, "Idiopathic leukonychia totalis and partialis in a 12-year-old patient," *Journal of the American Academy of Dermatology*, vol. 44, no. 2, supplement, pp. 379–380, 2001.
- [9] V. Bettoli and A. Tosti, "Leukonychia totalis and partialis: a single family presenting a peculiar course of the disease," *Journal of the American Academy of Dermatology*, vol. 15, no. 3, p. 535, 1986.
- [10] H.-J. Park, C.-N. Lee, J.-E. Kim, E. Jeong, J.-Y. Lee, and B.-K. Cho, "A case of idiopathic leukonychia totalis and partialis," *British Journal of Dermatology*, vol. 152, no. 2, pp. 401–402, 2005.
- [11] T. Butterworth, "Leukonychia partialis. A phase of leukonychia totalis," *Cutis*, vol. 29, no. 4, pp. 363–364, 1982.
- [12] R. A. Mees, "Een verschijnsel bij polyneuritis arsenicosa," *Nederlands Tijdschrift voor Geneeskunde*, vol. 1, pp. 391–396, 1919.
- [13] T. A. Giustina, T. Y. Woo, J. P. Campbell, and C. N. Ellis, "Association of pili torti and leukonychia," *Cutis*, vol. 35, no. 6, pp. 533–534, 1985.