Case Report

Eosinophilic Enteritis with Ascites in a Patient with Overlap Syndrome

Spyros Aslanidis,¹ Athina Pyrpasopoulou,¹ Kostas Soufleris,¹ Eirini Kazantzidou,² and Stella Douma¹

¹2nd Propedeutic Department of Internal Medicine, Hippokration General Hospital, 54643 Thessaloniki, Greece

²Radiology Department, Hippokration General Hospital, 54643 Thessaloniki, Greece

Correspondence should be addressed to Spyros Aslanidis, saslan@med.auth.gr

Received 27 January 2009; Accepted 30 April 2009

Recommended by Robert A. Eisenberg

Gastrointestinal involvement is frequent in patients with systemic lupus erythematosus (SLE). Eosinophilic gastroenteritis, however, has only rarely been described in rheumatological conditions, despite its reported connection to autoimmune diseases, such as hypereosinophilic syndrome, vasculitides, and systemic mastoidosis. It presents typically with abdominal pain and diarrhea and is only exceptionally associated with ascites. Diagnosis can be problematic, as several other clinical conditions (malignancies, infection/tuberculosis, and inflammatory bowel diseases) have to be ruled out. It is basically a nonsurgical disease, with excellent recovery on conservative treatment. We report the rare case of a young woman with overlap syndrome who presented with abdominal pain and ascites. The diagnosis of eosinophilic enteritis was made based on clinical, radiological, and laboratory criteria. The patient was treated with corticosteroids with excellent response.

Copyright © 2009 Spyros Aslanidis et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Gastrointestinal involvement is frequent in patients with systemic lupus erythematosus (SLE). In a series of 201 patients with childhood-onset SLE the most frequent symptom was abdominal pain. It was mostly related to lupus involvement, especially ascites, and pancreatitis, more rarely to treatment-induced events or infection and never to events unrelated to SLE [1]. Eosinophilic gastroenteritis has only rarely been described in rheumatological conditions, such as rheumatoid arthritis and systemic lupus erythematosus [2–6], despite its reported connection to autoimmune diseases, such as hypereosinophilic syndrome, vasculitides, and systemic mastoidosis. It presents typically with abdominal pain and diarrhea and is only exceptionally associated with ascites. Diagnosis can be difficult, as several other clinical conditions (malignancies, infection/tuberculosis, and inflammatory bowel diseases) have to be ruled out. Differential diagnosis from lupus enteritis (gastrointestinal vasculitis), a common cause of acute abdominal pain associated with small bowel wall thickening and ascites, involves exclusion of mesenteric ischemia [7].

We report the rare case of a young woman with overlap syndrome who presented with abdominal pain and ascites. Based on clinical, radiological, and laboratory criteria, the diagnosis of eosinophilic enteritis was made. The patient was treated conservatively with corticosteroids with excellent response.

2. Case Presentation

A 24-year-old woman presented with abdominal pain and vomiting. She was previously well till 24 hours earlier when she developed diffuse abdominal pain, nausea, and vomiting, with gradual worsening. On clinical examination, the patient was afebrile, tachycardic (108/min), with diffuse tenderness, guarding of the entire abdomen and abdominal distension; there were no active bowel sounds.

The patient's medical history included heterozygous β -thalassemia and a 6-year history of connective tissue disease with overlapping features of systemic lupus erythematosus and limited scleroderma, with positive antinuclear-ANA, antiribonucleoprotein-anti-RNP, anti-DNA and anti- β 2-glycoprotein antibodies for which she was currently on

FIGURE 1: Computed tomography scan of the abdomen performed. (a) On day 3 of the patient's admission, 4 days after initiation of symptoms. Arrows: thickening/dilatation of the intestinal wall, stars: ascitic fluid. (b) Four days after initiation of treatment, with marked improvement of the intestinal morphology, and moderate reduction of the amount of ascitic fluid.

methotrexate (10 mg/week), hydroxychloroquine (400 mg/day) and low dose aspirin. Two months prior to this episode she reported similar symptomatology, which responded well to "classical" treatment (ranitidine and metoclopramide).

On the 3rd day after admission her findings persisted; computed tomography (CT) scanning of the abdomen revealed dilatation and marked contrast-enhanced thickening of the small intestinal wall, mainly ileum; other parts of the small intestine (jejunum) and the large intestine were unaffected. A substantial amount of ascitic fluid was also present (Figure 1(a)). Large vessels were patent, with no signs of occlusion, venous thrombosis, intestinal ischemia, or intestinal obstruction; no pathological lymph nodes were observed.

The patient was referred to our hospital for further evaluation and treatment.

Diagnostic paracentesis of the ascitic fluid showed low SAAG (serum/ascites albumin gradient), elevated lactic

dehydrogenase (LDH), and presence of eosinophils (23% of 800 WBC/ μ L, 52% neutrophils, and 15% lymphocytes) in the ascitic fluid without peripheral eosinophilia; the number of peripheral eosinophils rose to 13% (1100/ μ L) on the 4th day of her hospitalization. The PPD (tuberculin, purified protein derivative) skin test and the cultures of the ascitic fluid were negative.

The diagnosis of eosinophilic enteritis was made based on clinical, laboratory, and radiological criteria. Endoscopy was not undertaken because of (a) the patient's severe clinical condition (peritonism and marked intestinal dilatation), (b) the difficulty in accessing the affected area (ileum), and (c) the strong probability of negative biopsies in this particular (subserosal) type of involvement [8].

The patient was treated with IV prednisolone (1 mg/kg daily for 7 days and gradual tapering) and IV cefotaxime, with rapid resolution of her symptoms and considerable clinical improvement. A subsequent scan, performed 4 days after initiation of treatment showed significant normalization of the morphology of the intestinal wall with moderate reduction of the ascitic fluid (Figure 1(b)). Five months later, the patient was in excellent clinical condition under a maintenance dose of methylprednisolone (2 mg every other day), soon to be discontinued.

3. Discussion

Eosinophilic gastroenteritis is a heterogeneous and uncommon disease, affecting children as well as adults, whose pathogenesis and etiology are not well understood. The diagnosis of eosinophilic gastroenteritis is based on a high degree of suspicion and represents a challenge to the physician, not only because of its rarity, but also because of the variety of clinical manifestations. No definite criteria exist; diagnosis is based on a combination of clinical, radiological, and histological findings. Clinically, the presence of gastrointestinal symptoms, the lack of evidence of parasitic or extraintestinal eosinophilic disease and the exclusion of intestinal lymphoma, Crohn's disease or other tumors, nodular and irregular thickening of the folds usually in the distal stomach and proximal small bowel on CT [9, 10], and the histologically proven intense eosinophilic infiltration in any of the different layers of at least one site of the gastrointestinal tract suffice for the diagnosis [11–13]. Eosinophilic gastroenteritis is basically a nonsurgical disease, with excellent recovery on conservative treatment [14, 15]. Steroids are usually indicated as first line therapy for patients with nonallergic eosinophilic gastroenteritis and provide prompt and effective relief of symptoms within a few days to weeks. Most experts recommend doses similar to those used in inflammatory bowel disease (1–2 mg/kg per day), tapered over a period of 3–4 months to induce remission. Relapses however are frequently noted during steroid tapering or after discontinuation, necessitating repeated steroid therapy [15]. Therefore, long-term follow-up is warranted, despite initial rapid improvement, since there is no agreement on the duration of treatment.

References

- [1] O. Richer, T. Ulinski, I. Lemelle, et al., "Abdominal manifestations in childhood-onset systemic lupus erythematosus," *Annals of the Rheumatic Diseases*, vol. 66, no. 2, pp. 174–178, 2007.
- [2] W. F. Ng, P. Cohen, A. Hepburn, S. Hamdulay, M. Carpani, and J. C. Mason, "A case of eosinophilic enteritis and rheumatoid arthritis," *Rheumatology*, vol. 44, no. 12, pp. 1585–1586, 2005.
- [3] P. R. Sunkureddi, N. Luu, S. Y. Siao, W. W. Tang, and B. A. Baethge, "Eosinophilic enteritis with systemic lupus erythematosus," *Southern Medical Journal*, vol. 98, no. 10, pp. 1049–1052, 2005.
- [4] D. A. Barbie, A. A. Mangi, and G. Y. Lauwers, "Eosinophilic gastroenteritis associated with systemic lupus erythematosus," *Journal of Clinical Gastroenterology*, vol. 38, no. 10, pp. 883–886, 2004.
- [5] K. DeSchryver-Keckskemeti and R. E. Clouse, "A previously unrecognized subgroup of "eosinophilic gastroenteritis". Association with connective tissue diseases," *American Journal of Surgical Pathology*, vol. 8, no. 3, pp. 171–180, 1984.
- [6] A. L. Buchman, D. Wolf, and T. Gramlich, "Eosinophilic gastrojejunitis associated with connective tissue disease," *Southern Medical Journal*, vol. 89, no. 3, pp. 327–330, 1996.
- [7] C.-K. Lee, M. S. Ahn, E. Y. Lee, et al., "Acute abdominal pain in systemic lupus erythematosus: focus on lupus enteritis (gastrointestinal vasculitis)," *Annals of the Rheumatic Diseases*, vol. 61, no. 6, pp. 547–550, 2002.
- [8] Y. Q. Hsu and C. Y. Lo, "A case of eosinophilic gastroenteritis," *Hong Kong Medical Journal*, vol. 4, no. 2, pp. 226–228, 1998.
- [9] X. Zheng, J. Cheng, K. Pan, K. Yang, H. Wang, and E. Wu, "Eosinophilic enteritis: CT features," *Abdominal Imaging*, vol. 33, no. 2, pp. 191–195, 2008.
- [10] K. M. Horton, F. M. Corl, and E. K. Fishman, "CT of nonneoplastic diseases of the small bowel: spectrum of disease," *Journal of Computer Assisted Tomography*, vol. 23, no. 3, pp. 417–428, 1999.
- [11] M. Y. Yun, Y. U. Cho, I. S. Park, et al., "Eosinophilic gastroenteritis presenting as small bowel obstruction: a case report and review of the literature," *World Journal of Gastroenterology*, vol. 13, no. 11, pp. 1758–1760, 2007.
- [12] A. Charalabopoulos, K. Charalabopoulos, M. Avuzuklidou, et al., "Eosinophilic gastroenteritis: presentation of two patients with unusual affect of terminal ileum and caecum with manifestations of acute abdomen and literature review," *International Journal of Clinical Practice*, vol. 58, no. 4, pp. 413–416, 2004.
- [13] M.-J. Chen, C.-H. Chu, S.-C. Lin, S.-C. Shih, and T.-E. Wang, "Eosinophilic gastroenteritis: clinical experience with 15 patients," *World Journal of Gastroenterology*, vol. 9, no. 12, pp. 2813–2816, 2003.
- [14] M. E. Rothenberg, "Eosinophilic gastrointestinal disorders (EGID)," *Journal of Allergy and Clinical Immunology*, vol. 113, no. 1, pp. 11–28, 2004.
- [15] S. Khan, "Eosinophilic gastroenteritis," *Best Practice and Research: Clinical Gastroenterology*, vol. 19, no. 2, pp. 177–198, 2005.