Case Report

Schnitzler's Syndrome: A Case Report

Gabriel Tinoco,¹ Rehan Kanji,² and Deepthi Moola³

¹ Division of Hospital Medicine, Department of Medicine, University of Miami Miller School of Medicine, 1120 NW 14th Street, Suite 1185 CRB, Miami, FL 33136, USA

² American University of Antigua, 1 Battery Park Plaza, New York, NY 10004, USA

³ Department of Obstetrics and Gynecology, TriHealth, 619 Oak Street, Cincinnati, OH 45206, USA

Correspondence should be addressed to Gabriel Tinoco; gtinoco@med.miami.edu

Received 16 February 2013; Accepted 23 May 2013

Academic Editor: Abhay R. Satoskar

Copyright © 2013 Gabriel Tinoco et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Schnitzler's syndrome is an extremely rare entity that poses a challenge for the clinician not only due to its difficult diagnosis but also due to its management. In this article we report a new case and briefly review the current treatment options.

1. Introduction

Schnitzler's syndrome is an extremely rare and not completely understood entity. It is characterized by a chronic recurrent urticarial rash with leukocytoclastic vasculitis, monoclonal IgM gammopathy, intermittent fevers, arthralgia, bone pain, lymphadenopathy, skeletal hyperostosis, and occasionally hepato- or splenomegaly [1–7].

To date only around 100 cases [1, 2, 8] have been reported since the first described case in 1972 [9].

This autoinflammatory syndrome clinically presents with a diverse constellation of symptoms making its initial diagnosis very challenging. Its early recognition is key due to the increased risk of lymphoproliferative disorders (10–15% per year [2]) and its tremendous impact on the quality of life of the patients with this entity.

Due to its obscure pathophysiology, the treatment remains not well defined, being steroids the first line of treatment [3, 10]. However different treatment modalities, with different range of responses, have been reported including NSAIDs, [3] thalidomide [11, 12], methotrexate [13], cyclosporine [14], cyclophosphamide [15], psoralen plus ultraviolet [16], interferon [17], and recently biological therapies such as TNF α blockade [18, 19], Anakinra [12, 20–24], Rituximab [25, 26], and Tocilizumab [27, 28] (Table 2).

2. Case Report

A 49-year-old Caucasian female presented with generalized arthralgias and recurrent diffuse pruritic maculopapular lesions over her face, torso, and upper extremities. The patient suffered outbreaks with variable intensity almost in a daily basis over the past five years with no identifiable triggering factors.

The patient mentioned that the lesions cleared up in 24 hours, leaving no marks or scars, while new lesions appeared daily.

Her past medical history was significant for transfusion related hepatitis C, diabetes mellitus type 2, depression, chronic pancreatitis, and malnutrition.

Previous surgeries included pancreatic cyst removal at age 11, tonsillectomy/adenoidectomy, tubal ligation, and appendectomy.

She had a 35 pack/year history of smoking, with remote history of inhaled crack cocaine use. She denied alcohol or other illicit drug use.

Review of systems revealed unintended weight loss of about 20 pounds over a 6-month period, diffuse abdominal pain, generalized weakness, night sweats, and subjective recurrent low-grade fevers not associated with the onset of her skin lesions.

TABLE 1: Laboratories at admission.

(a)			
WBC	48.6		
Differential	97.8% <i>N</i>		
Hemoglobin	13		
Hematocrit	40.9		
Platelets	321		
(b)			
D-dimer	565	BNP	489
CK	<20	CKMB	<0.3
Trop I	<0.1	Lactic Acid	2.6
Amylase	<30	Lipase	42
Alk. phos	259	AST/ALT	11/14
PT	15.8	PTT	37.3
INR	1.3	ESR	80
(c)			
Sodium	125	Potassium	4.9
Chloride	92	Bicarbonate	27
BUN	16	Creatinine	0.6
Glucose	230	Calcium	8.1
Phosphorous	4.0	Magnesium	1.5

Her family history was noncontributory (father deceased at the age of 70 because of heart disease, mother had diabetes mellitus and coronary artery disease, and her brother and daughter had good health).

The initial physical examination revealed diffuse pruritic maculopapular lesions over her face, thorax, and upper extremities (Figure 1), sparing her palms, soles, and mucous membranes. Also mild hepatosplenomegaly was found.

Her laboratory studies showed marked leukocytosis of 48.6 with neutrophils of 97.8% with elevated D-dimer and erythrocyte sedimentation rate (ESR) (Table 1).

She was started on broad-spectrum antibiotics that were discontinued 72 hours after the admission when infections were ruled out.

Antinuclear antibodies, rheumatoid factor, anticitrullinated peptide antibodies, human immunodeficiency virus enzyme-linked immunosorbent assay (ELISA), Venereal Disease Research Laboratory (VDRL), hepatitis B surface antigen (HBsAg), hepatitis B core antigen (HBcAg), hepatitis B e antigen (HBeAg), and cryoglobulins were found to be negative or within normal limits.

Serum protein electrophoresis showed monoclonal gammopathy; SPEP showed gamma region asymmetric spike consistent with monoclonal gammopathy. UPEP showed oligoclonal gammopathy involving IgG, IgM kappa, and lambda (IgM: 552 mg/L).

Biopsy of her skin lesions showed neutrophil rich cells admixed with eosinophils consistent with chronic urticaria.

TABLE 2: Common therapeutic strategies for Schnitzler's syndrome.

Treatment	Efficacy	References
Corticosteroids	High doses may achieve complete remission in about 39% of the cases.	[1–3, 7, 8, 11, 12, 14–17, 29, 30]
NSAID's	Partial or temporary remission of the symptoms in about 15% of the cases.	[2, 3, 6–8, 11, 12, 29, 31]
Antihistamines	Partial or temporary remission of the symptoms.	[2, 3, 8, 12, 16, 29, 30]
Cyclophosphamide, methotrexate	Partial remission in occasional cases.	[2, 8, 13, 15]
Anakinra	Complete remission in few cases.	[2, 8, 12, 20–24]

A right groin excisional lymph node biopsy showed reactive lymphoid hyperplasia with plasmacytosis.

Bone marrow biopsy showed a hypercellular marrow with plasmacytosis and myeloid predominance and was reported as negative for malignancy.

Based on the clinical presentation and laboratory results the diagnosis of Schnitzler syndrome was made. As proposed by Lipsker et al. [3] the diagnosis of this entity requires 2 major criteria and a minimum of 2 minor criteria. She had 2 major criteria (chronic urticarial rash/intermittent fever and monoclonal IgM) plus at least 5 minor criteria (arthralgias, bone pain, lymphadenopathy, hepato- and/or splenomegaly, and elevated ESR and/or leukocytosis).

She was started on high dose of corticosteroids, nonsteroidal anti-inflammatory drugs, and antihistamines with partial remission of her symptoms and she was discharged home (Figure 2).

Anakinra was added to her regimen for several weeks with an excellent response; however due to economical constraints treatment was discontinued, and her symptoms partially recurred.

3. Conclusions

Schnitzler's syndrome is an underdiagnosed disease with only around 100 cases reported in the literature. Having eastern European origin [1, 2, 7, 8] and being a woman (women are more frequently affected than man (1.6:1) [2]) are strong associations to this entity.

The mortality risk is not increased in patients with Schnitzler's syndrome (survival of 94% after 15 years of the diagnosis) [2]; however there is a well-established increased risk of malignancy (specially lymphoproliferative disease such as Waldenström macroglobulinemia and IgM myeloma) of 10-year risk of 15% [2, 31].

FIGURE 1: Urticaria as seen on the skin of the back, thigh, and arm of patient (from left to right).

FIGURE 2: Progression of WBC count throughout the hospital course.

Systemic steroids remain as the first line of treatment [3, 8, 10], being effective to less than a half of the patients [8]. The data available demonstrated that the treatment of Schnitzler syndrome is not standardized, and there are promising results with new biological therapies such as the interleukin-1 receptor antagonist Anakinra [12, 20–24], Rituximab [25, 26], and Tocilizumab [27, 28].

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images.

Conflict of Interests

The authors declare that they have no conflict of interests.

Authors’ Contribution

Gabriel Tinoco, Rehan Kanji, and Deepthi Moola reviewed and analyzed the patient data and wrote the paper. All authors read and approved the final paper.

Acknowledgments

The authors would like to acknowledge Richard B. Williams M.D., Barry Materson M.D., and Joseph Rosenblatt M.D. for their support in the elaboration of this paper.

References

- [1] R. Akimoto, M. Yoshida, R. Matsuda, K. Miyasaka, and M. Itoh, "Schnitzler's syndrome with IgG κ gammopathy," *Journal of Dermatology*, vol. 29, no. 11, pp. 735–738, 2002.
- [2] H. D. de Koning, E. J. Bodar, J. W. M. van der Meer, and A. Simon, "Schnitzler syndrome: beyond the case reports: review and follow-up of 94 patients with an emphasis on prognosis and treatment," *Seminars in Arthritis and Rheumatism*, vol. 37, no. 3, pp. 137–148, 2007.
- [3] D. Lipsker, Y. Veran, F. Grunenberger, B. Cribier, E. Heid, and E. Grosshans, "The Schnitzler syndrome: four new cases and review of the literature," *Medicine*, vol. 80, no. 1, pp. 37–44, 2001.
- [4] M. Janier, D. Bonvalet, M.-F. Blanc et al., "Chronic urticaria and macroglobulinemia (Schnitzler's syndrome): report of two cases," *Journal of the American Academy of Dermatology*, vol. 20, no. 2 I, pp. 206–211, 1989.
- [5] L. Schnitzler, D. Hurez, and J. L. Verret, "Chronic urticaria—osteo-condensation—macroglobulinemia," *Annales de Dermatologie et de Venereologie*, vol. 116, no. 8, pp. 547–550, 1989.
- [6] L. Borradori, M. Rybojad, A. Puissant, A. Dallot, O. Verola, and P. Morel, "Urticarial vasculitis associated with a monoclonal IgM gammopathy: Schnitzler's syndrome," *British Journal of Dermatology*, vol. 123, no. 1, pp. 113–118, 1990.
- [7] A. Morita, S. Sakakibara, M. Yokota, and T. Tsuji, "A case of urticarial vasculitis associated with macroglobulinemia (Schnitzler's syndrome)," *Journal of Dermatology*, vol. 22, no. 1, pp. 32–35, 1995.
- [8] E. Tinazzi, A. Puccetti, G. Patuzzo, M. Sorletto, A. Barbieri, and C. Lunardi, "Schnitzler syndrome, an autoimmune-autoinflammatory syndrome: report of two new cases and review of the literature," *Autoimmunity Reviews*, vol. 10, no. 7, pp. 404–409, 2011.
- [9] L. Schnitzler, "Lésions urticariennes chroniques permanentes (érythème pétaoloïde?)," *Journée Dermatologique d'Angers, Cas cliniques*, no. 46 B, 1972.
- [10] P. Goupille, P. Pizzuti, E. Diot, F. Jattiot, J.-L. Guilmot, and J.-P. Valat, "Schnitzler's syndrome (urticaria and macroglobulinemia) dramatically improved with corticosteroids," *Clinical and Experimental Rheumatology*, vol. 13, no. 1, pp. 95–98, 1995.
- [11] M. Worm and G. Kolde, "Schnitzler's syndrome: successful treatment of two patients using thalidomide," *British Journal of Dermatology*, vol. 148, no. 3, pp. 601–602, 2003.
- [12] H. D. De Koning, E. J. Bodar, A. Simon, J. C. H. van der Hilst, M. G. Netea, and J. W. M. van der Meer, "Beneficial response to anakinra and thalidomide in Schnitzler's syndrome," *Annals of the Rheumatic Diseases*, vol. 65, no. 4, pp. 542–544, 2006.
- [13] S. Bozeman and S. Lewis, "Schnitzler syndrome successfully treated with methotrexate," *Annals of Allergy, Asthma and Immunology*, vol. 101, no. 2, article 219, 2008.
- [14] M. Pascual-Lopez, A. Hernandez-Nunez, J. Sanchez-Perez et al., "Schnitzler's syndrome with monoclonal IgG kappa gammopathy: good response to cyclosporin," *Journal of the European Academy of Dermatology and Venereology*, vol. 16, no. 3, pp. 267–270, 2002.
- [15] D. Peterlana, A. Puccetti, E. Tinazzi, S. Simeoni, and C. Lunardi, "Schnitzler's syndrome treated successfully with intravenous pulse cyclophosphamide," *Scandinavian Journal of Rheumatology*, vol. 34, no. 4, pp. 328–330, 2005.
- [16] G. Cianchini, L. Colonna, F. Bergamo, C. Angdo, and P. Puddu, "Efficacy of psoralen-UV-A therapy in 3 cases of Schnitzler syndrome," *Archives of Dermatology*, vol. 137, no. 11, pp. 1536–1537, 2001.
- [17] N. E. C. Schartz, S. Buder, H. Sperl et al., "Report of a case of Schnitzler's syndrome treated successfully with interferon alpha 2b," *Dermatology*, vol. 205, no. 1, pp. 54–56, 2002.
- [18] N. E. Aikawa, C. A. A. Silva, E. Bonfá, and J. F. D. Carvalho, "Schnitzler's syndrome improvement after anti-TNF-alpha therapy," *Joint Bone Spine*, vol. 77, no. 5, article 491, 2010.
- [19] R. Thonhofer, E. Uitz, and W. Graninger, "Schnitzler's syndrome—exacerbation after anti-TNF treatment," *Rheumatology*, vol. 46, no. 6, pp. 1041–1042, 2007.
- [20] E. Besada and H. Nossent, "Dramatic response to IL1-RA treatment in longstanding multidrug resistant Schnitzler's syndrome: a case report and literature review," *Clinical Rheumatology*, vol. 29, no. 5, pp. 567–571, 2010.
- [21] S. W. Schneider, M. Gaubitz, T. A. Luger, and G. Bonsmann, "Prompt response of refractory Schnitzler syndrome to treatment with anakinra," *Journal of the American Academy of Dermatology*, vol. 56, supplement 5, pp. S120–S122, 2007.
- [22] C. Schuster, B. Kränke, E. Aberer, E. Arbab, G. Sturm, and W. Aberer, "Schnitzler syndrome: response to anakinra in two cases and a review of the literature," *International Journal of Dermatology*, vol. 48, no. 11, pp. 1190–1194, 2009.
- [23] P. A. Frischmeyer-Guerrerio, R. Rachamalla, and S. S. Saini, "Remission of Schnitzler syndrome after treatment with anakinra," *Annals of Allergy, Asthma and Immunology*, vol. 100, no. 6, pp. 617–619, 2008.
- [24] H. Wastiaux, S. Barbarot, V. Gagey-Caron et al., "Schnitzler syndrome: a dramatic improvement with anakinra," *Journal of the European Academy of Dermatology and Venereology*, vol. 23, no. 1, pp. 85–87, 2009.
- [25] K. M. Ramadan, H. A. Eswedi, and M. R. El-Agnaf, "Schnitzler syndrome: a case report of successful treatment using the anti-CD20 monoclonal antibody rituximab," *British Journal of Dermatology*, vol. 156, no. 5, pp. 1072–1074, 2007.
- [26] N. Cascavilla, M. Bisceglia, and G. D'Arena, "Successful treatment of Schnitzler's syndrome with anakinra after failure of rituximab trial," *International Journal of Immunopathology and Pharmacology*, vol. 23, no. 2, pp. 633–636, 2010.
- [27] C. Ding and G. Jones, "Anti-interleukin-6 receptor antibody treatment in inflammatory autoimmune diseases," *Reviews on Recent Clinical Trials*, vol. 1, no. 3, pp. 193–200, 2006.
- [28] N. Kluger, D. Bessis, and B. Guillot, "Tocilizumab as a potential treatment in Schnitzler syndrome," *Medical Hypotheses*, vol. 72, no. 4, pp. 479–480, 2009.
- [29] S. S. Berdy and K. J. Bloch, "Schnitzler's syndrome: a broader clinical spectrum," *Journal of Allergy and Clinical Immunology*, vol. 87, no. 4, pp. 849–854, 1991.
- [30] J. P. Clauvel, J. C. Brouet, and F. Danon, "Chronic urticaria with monoclonal IgM—a report of five cases," *Clinical Immunology and Immunopathology*, vol. 25, no. 3, pp. 348–353, 1982.
- [31] W. Lim, K. H. Shumak, M. Reis et al., "Malignant evolution of Schnitzler's syndrome—chronic urticaria and IgM monoclonal gammopathy: report of a new case and review of the literature," *Leukemia and Lymphoma*, vol. 43, no. 1, pp. 181–186, 2002.