Case Report

Polymicrobial Bacteremia Involving *Comamonas testosteroni*

Jose Orsini,^{1,2} Eric Tam,¹ Naomi Hauser,¹ and Salil Rajayer¹

¹Department of Medicine, New York University School of Medicine, Woodhull Medical and Mental Health Center, Brooklyn, NY 11206, USA

²Division of Critical Care Medicine, Department of Medicine, New York University School of Medicine, Woodhull Medical and Mental Health Center, Brooklyn, NY 11206, USA

Correspondence should be addressed to Jose Orsini; jose.orsini@woodhullhc.nychhc.org

Received 9 November 2014; Accepted 16 December 2014; Published 28 December 2014

Academic Editor: W. Zidek

Copyright © 2014 Jose Orsini et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Comamonas spp. are uncommon isolates in microbiology laboratories and have been rarely observed as an infectious agent in clinical practice. They have widespread environmental distribution and have been isolated from water, soil, and plants as well as from some hospital devices such as intravenous catheters and water contained in humidifier reservoirs used in respiratory treatment. The genus *Comamonas* originally contained the following species: *acidovorans*, *testosteroni*, *kerstersii*, *terrigena*, *denitrificans*, and *nitrativorans*. It now contains 17 species, while *acidovorans* spp. have been reclassified as *Delftia acidovorans*. In spite of its uncommon human pathogenesis, there are few reports on the aggressive manner of it as an opportunistic pathogen, mostly related to *testosteroni* spp. We present a case of polymicrobial bacteremia involving *Comamonas testosteroni*. The aim of this case report is to alert clinicians to the potential diagnosis of bloodstream infections caused by uncommon pathogens.

1. Introduction

Comamonas testosteroni, formerly known as *Pseudomonas testosteroni*, is an aerobic, motile, nonspore forming, ubiquitous Gram-negative organism. Although it has low virulence potency, there are few reports on its involvement in human infections. It became clinically important after 1987, when reports began accumulating on human infections such as cellulitis [1], peritonitis [2], endocarditis [3], meningitis [4], endophthalmitis [5], tenosynovitis [6], and pneumonia [7]. However, cases of bloodstream infections caused by *Comamonas testosteroni* have been infrequently reported [2, 8–12]. The name of *testosteroni* was given due to its characteristic of utilizing carbon from the metabolism of testosterone [13], although this property has also been demonstrated by some *Pseudomonas* spp. and fungi [14]. Some strains of *Comamonas testosteroni* have acquired the plasmid-mediated *bla*NDM1, gene that confers broad spectrum antimicrobial resistance thus potentially hampering treatment options in the event where this bacterium causes human infections [15]. In this report, we present an uncommon case of polymicrobial bacteremia involving *Comamonas testosteroni* with a detailed review of the literature.

2. Case Report

An 80-year-old African American female, morbidly obese, was brought to emergency department complaining of generalized body aches (predominantly in shoulders) and nondocumented fevers at home for about 1 week. Her past medical history was significant for systemic arterial hypertension, diabetes mellitus, hiatal hernia, osteoarthritis, and cholelithiasis (s/p cholecystectomy). On arrival to the hospital, her vital signs were as follows: blood pressure of 150/77 mmHg, a regular heart rate of a 106 beats/minute, a respiratory rate of 21 breaths/minute, a temperature of a 100.8°F, and an oxygen saturation of 96% breathing ambient air.

Remarkable laboratory findings on admission were a white blood cell (WBC) count of 14, 500/mm³ (4.8–10.8), a sodium level of 129 mmol/L (135–147), a potassium level of 5.5 mmol/L (3.5–5.3), and a creatinine level of 2.0 mg/dL (0.8–2.0). C-reactive protein (CRP) and erythrocyte sedimentation rate (ESR) levels were both elevated at 152.7 mg/L (1.0–4.0) and 130 mm/h (0–20), respectively. Her urine analysis showed positive leukoesterase and a WBC count of 10–15/hpf (0–5). Coagulation and liver profiles were within normal limits. Bilateral shoulder X-ray demonstrated degenerative changes,

with a right rotator cuff tendinitis but no fractures. Chest radiography revealed no infiltrates.

She was admitted to the medicine ward with the presumptive diagnosis of urinary tract infection (UTI), and empiric antimicrobial therapy was initiated with ceftriaxone 2 grams intravenously daily. Blood cultures drawn on admission grew methicillin-sensitive *Staphylococcus aureus*, and antimicrobial therapy was streamlined to nafcillin 2 grams intravenously every 4 hours. Transthoracic echocardiogram (TTE) showed no vegetations. On day 3 of admission, her hospital course was complicated by worsening leukocytosis and a transient episode of hypotension which responded to intravenous fluids therapy. At that time, a new set of blood cultures was obtained from which *Comamonas testosteroni* was isolated (sensitive to ceftazidime, carbapenems, piperacillin/tazobactam, and trimethoprim/sulfamethoxazole). Antimicrobial therapy was then escalated to cefazolin 1 gram intravenously every 8 hours plus doripenem 250 mg intravenously every 8 hours. Abdomen, pelvis, and chest computed tomography (CT) were obtained, demonstrating the presence of diverticulosis but no abscess formation or mass. Repeated blood cultures were negative, and the patient was discharged home to complete a total of 4-week course of intravenous antimicrobial therapy.

3. Discussion

This is one of fewer than 15 cases of *Comamonas testosteroni* bacteremia reported in the literature and the first reported case of *Comamonas testosteroni* and *Staphylococcus aureus* polymicrobial bloodstream infection, according to a MEDLINE search using the words “*Comamonas*” and “bacteremia” and “bloodstream infection.” Approximately 35 cases of infections involving *Comamonas testosteroni* have been reported in the medical literature. Majority of infections are community-acquired rather than nosocomial [3, 5], and most of the patients reported had some degree of immunosuppression such as malignancy, chronic liver disease, and end-stage renal disease requiring hemodialysis [1, 11, 12]. Bacterial translocation from the gastrointestinal tract seems to play an important role in the pathogenesis of infections caused by *Comamonas* spp. [9, 16].

Based on results from previous reports, majority of the 14 reported patients with bacteremia caused by *Comamonas testosteroni* were males, their median age was 47 years (range, newborn–89), only one patient died, and predisposing factors for infection were not identified in two of them. Most of the patients responded well to appropriate antimicrobial therapy [12]. *Comamonas* spp. isolates are usually susceptible to aminoglycosides, fluoroquinolones, carbapenems, piperacillin-tazobactam, cephalosporins, and trimethoprim-sulfamethoxazole [1, 4].

The patient discussed in this report did not have any obvious source of infection. She did not have central venous catheters, and the abdominal imaging did not evidence any acute inflammatory process or mass. Her immunosuppressive state was limited to diabetes mellitus, which might have played a role as a possible predisposing factor. Therefore,

the authors hypothesized that the most likely source might have been the right shoulder rotator cuff tendinitis. Although this hypothesis cannot be proven, it is supported by the fact that the patient reported shoulder pain, the CRP and ESR levels were markedly elevated, and the right shoulder imaging showed rotator cuff tendinitis. In 2000, Isotalo et al. reported a case of polymicrobial tenosynovitis, involving an organism most likely to be related to *Comamonas* spp. based on biochemical factors [6]. If proven, our patient may have been only the second case of tenosynovitis caused by *Comamonas* spp. reported in the literature. We also postulate that the second likely mode of bacteremia in this case could have been dissemination of *Comamonas testosteroni* from the bowel following gastrointestinal translocation, given the finding of extensive diverticulosis demonstrated in the abdominal imaging.

Polymicrobial bacteremia involving *Comamonas testosteroni* have been reported in association with *Streptococcus parasanguis* and *Ralstonia pickettii* [17]. Two other cases of polymicrobial bacteremia involving *Comamonas* spp. (*Delftia acidovorans* (formerly known as *Comamonas acidovorans*) and *Comamonas kerstersii*) were previously reported in association with *Streptococcus agalactiae* and *Bacteroides fragilis*, respectively [17]. To our knowledge, this is the first case of bloodstream infection involving *Comamonas testosteroni* and *Staphylococcus aureus*.

In this report, we have highlighted an unusual cause of bloodstream infection. *Comamonas* spp. should be kept in mind as a rare cause of bacteremia.

Conflict of Interests

The authors report no conflict of interests regarding the publication of this paper.

References

- [1] T.-L. Tsui, S.-M. Tsao, K.-S. Liu et al., “*Comamonas testosteroni* infection in Taiwan: reported two cases and literature review,” *Journal of Microbiology, Immunology and Infection*, vol. 44, no. 1, pp. 67–71, 2011.
- [2] D. J. Barbaro, P. A. Mackowiak, S. S. Barth, and P. M. Southern Jr., “*Pseudomonas testosteroni* infections: eighteen recent cases and a review of the literature,” *Reviews of Infectious Diseases*, vol. 9, no. 1, pp. 124–129, 1987.
- [3] G. R. Cooper, E. D. Staples, K. A. Iczkowski, and C. J. Clancy, “*Comamonas (Pseudomonas) testosteroni* endocarditis,” *Cardiovascular Pathology*, vol. 14, no. 3, pp. 145–149, 2005.
- [4] B. Arda, S. Aydemir, T. Yamazhan, A. Hassan, A. Tünger, and D. Serter, “*Comamonas testosteroni* meningitis in a patient with recurrent cholesteatoma,” *Acta Pathologica, Microbiologica et Immunologica Scandinavica*, vol. 111, no. 4, pp. 474–476, 2003.
- [5] A. K. Reddy, S. I. Murthy, S. Jalali, and U. Gopinathan, “Post-operative endophthalmitis due to an unusual pathogen, *Comamonas testosteroni*,” *Journal of Medical Microbiology*, vol. 58, no. 3, pp. 374–375, 2009.
- [6] P. A. Isotalo, D. Edgar, and B. Toye, “Polymicrobial tenosynovitis with *Pasteurella multocida* and other Gram negative bacilli after a Siberian tiger bite,” *Journal of Clinical Pathology*, vol. 53, no. 11, pp. 871–872, 2000.

- [7] F. Franzetti, M. Cernuschi, R. Esposito, and M. Moroni, "Pseudomonas infections in patients with AIDS and AIDS-related complex," *Journal of Internal Medicine*, vol. 231, no. 4, pp. 437–443, 1992.
- [8] M. D. Smith and J. D. Gradon, "Bacteremia due to *Comamonas* species possibly associated with exposure to tropical fish," *Southern Medical Journal*, vol. 96, no. 8, pp. 815–817, 2003.
- [9] M. Gul, P. Ciragil, E. Bulbuloglu, M. Aral, S. Alkis, and F. Ezberci, "*Comamonas testosteroni* bacteremia in a patient with perforated acute appendicitis," *Acta Microbiologica et Immunologica Hungarica*, vol. 54, no. 3, pp. 317–321, 2007.
- [10] J. M. Abraham and G. L. Simon, "*Comamonas testosteroni* bacteremia: a case report and review of the literature," *Infectious Diseases in Clinical Practice*, vol. 15, no. 4, pp. 272–273, 2007.
- [11] W. Nseir, J. Khateeb, M. Awawdeh, and M. Ghali, "Catheter-related bacteremia caused by *Comamonas testosteroni* in a hemodialysis patient," *Hemodialysis International*, vol. 15, no. 2, pp. 293–296, 2011.
- [12] S. Farshad, F. Norouzi, M. Aminshahidi, B. Heidari, and A. Alborzi, "Two cases of bacteremia due to an unusual pathogen, *Comamonas testosteroni* in Iran and a review literature," *Journal of Infection in Developing Countries*, vol. 6, no. 6, pp. 521–525, 2012.
- [13] M. Horinouchi, T. Yamamoto, K. Taguchi, H. Arai, and T. Kudo, "Meta-cleavage enzyme gene *tesB* is necessary for testosterone degradation in *Comamonas testosteroni* TA441," *Microbiology*, vol. 147, no. 12, pp. 3367–3375, 2001.
- [14] J. Tamaoka, D.-M. Ha, and K. Komagata, "Reclassification of *Pseudomonas acidovorans* den Dooren de Jong 1926 and *Pseudomonas testosteroni* Marcus and Talalay 1956 as *Comamonas acidovorans* comb. nov. and *Comamonas testosteroni* comb. nov., with an emended description of the genus *Comamonas*," *International Journal of Systematic Bacteriology*, vol. 37, no. 1, pp. 52–59, 1987.
- [15] X. Wang, W. Liu, D. Zou et al., "High rate of New Delhi Metallo- β -Lactamase 1-producing bacterial infection in China," *Clinical Infectious Diseases*, vol. 56, no. 1, pp. 161–162, 2013.
- [16] H. Hagiya, T. Murase, J. Sugiyama et al., "*Delftia acidovorans* bacteremia caused by bacterial translocation after organophosphorus poisoning in an immunocompetent adult patient," *Journal of Infection and Chemotherapy*, vol. 19, no. 2, pp. 338–341, 2013.
- [17] O. Opota, B. Ney, G. Zanetti, K. Jatton, G. Greub, and G. Prod'homme, "Bacteremia caused by *Comamonas kerstersii* in a patient with diverticulosis," *Journal of Clinical Microbiology*, vol. 52, no. 3, pp. 1009–1012, 2014.