

Case Report

Superior Dislocation of the Patella in a Young Patient without Osteophytes: A Case Report with Discussion about Differential Diagnosis

Wander Edney de Brito ,^{1,2} Gustavo Constantino Campos,³ João Batista de Miranda,³ and Alessandro Rozim Zorzi ,^{1,3,4}

¹São Leopoldo Mandic School of Medicine, Campinas, Brazil

²Hospital Municipal Mário Gatti, Campinas, Brazil

³State University of Campinas (UNICAMP), Campinas, Brazil

⁴Hospital Israelita Albert Einstein, São Paulo, Brazil

Correspondence should be addressed to Alessandro Rozim Zorzi; arzorzi@hc.unicamp.br

Received 3 October 2018; Revised 14 January 2019; Accepted 31 January 2019; Published 27 March 2019

Academic Editor: John Kortbeek

Copyright © 2019 Wander Edney de Brito et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We report a case of superior dislocation of the patella in a young woman without degenerative changes. We retrospectively analyzed the clinical and imaging data obtained from the patient. This article describes a rare case of patellar dislocation following a bicycle fall in a 19-year-old woman without any history of patellofemoral complaints. Our literature search yielded 28 case reports; however, most reports describe older individuals with osteoarthritis. Only two reports have previously described this lesion in young patients without osteophytes, but some features, like an increase of the patella tilt, may raise doubts about whether it would be better to classify them as a vertical dislocation of the patella, another quite rare lesion, or just as a variant of a superior dislocation.

1. Introduction

Superior dislocation of the patella (SDP) is a rare condition that is often associated with degenerative changes and the presence of osteophytes, with peak incidence observed in the sixth decade of life [1]. The distal articular edge of the patella engages in the proximal articular edge of the trochlea, locking the knee in full extension.

Although rare, this condition requires urgent treatment. Patients with SDP often present to the Emergency Department (ED) with severe pain and restriction of movement. It is important to not misdiagnose SDP as an acute rupture of the patellar tendon based on radiographic findings. Careful history taking and clinical examination are important to differentiate between these conditions [2]. SDP requires immediate closed reduction, whereas patellar tendon rupture requires surgical treatment in a nonemergency setting.

This case report describes one of the youngest known patients diagnosed with SDP.

2. Case Presentation

Ethical approval and informed consent to report this case and its figures were obtained from the Ethics Committee of the State University of Campinas (approval number 2.878.038/ID 95776318.8.0000.5404).

A 19-year-old woman fell from a bicycle and hit her knee against a street guide. She presented with severe pain and an anteromedial skin bruise on the left knee and her knee locked in full extension (Figure 1(a)).

Radiographs showed superior displacement of the patella with its inferior articular surface engaging the proximal articular surface of the trochlea (Figure 1(b)).

We administered 2% lidocaine hydrochloride intraarticularly to treat her pain. Superior-inferior and lateral-




FIGURE 1: Image shows the patient's left knee upon arrival at the ED, after a fall from a bicycle (a). It is possible to notice that there is no bulging in the anterior region of the knee found in the VDP. SDP is suspected based on the attachment of the distal end of the patella to the proximal end of the trochlea (arrow). The high riding patella needs to be distinguished from a patellar tendon rupture (b). Radiograph of the left knee obtained after the procedure shows successful reduction (c). ED: emergency department; SDP: superior dislocation of the patella.

lateral patellar manipulations were unsuccessful, and we subsequently grasped and anteriorized the patella in relation to the femur, which led to immediate reduction. A plain control radiograph was obtained following this procedure and showed complete reduction of the patellofemoral joint (Figure 1(c)).

To ensure patient comfort, the knee was immobilized for a week in an inguino malleolar orthosis at 20 degrees of flexion. Upon her return 10 days later, the patient was completely pain free and asymptomatic without any movement limitations.

3. Discussion

Watson-Jones first described SDP in 1946 [3]. All reports since then have described SDP in older patients presenting with trochlear and patellar osteophytes causing engagement of the osteophytes at the lower pole of the patella and at the proximal trochlea [4]. Other authors have reported similar cases in old patients with knee osteoarthritis, always with locking osteophytes.

In 2007, Saleemi et al. [5] first reported SDP in a young patient with no radiographic evidence of osteoarthritis and no osteophytes. In 2016, Kataoka et al. [6] described

the second case in a 19-year-old woman who developed SDP secondary to direct knee trauma. Our case is the third report describing SDP in young patient without osteophyte.

However, it is noteworthy that, in our case and in the two previously cited [5, 6], the patella presents an increased tilt in the lateral radiography, which does not occur in cases with presence of osteophyte. This can cause confusion with another very rare lesion, the vertical dislocation of the patella (VDP) [7, 8]. Would these cases be better classified as VDP? In this way, there would be no cases of SDP without osteophytes. According to van Egmond et al. [4], there is a pathognomonic signal of the SDP: the proximal part of the patella is tilted away from the femur. This is caused by the pull of the patella tendon and the simultaneous relaxation of the quadriceps tendon. Kataoka et al. [6] performed a CT scanning showing clearly that, despite the increased patella tilt and the absence of a higher patella height, there is the attachment of the articular surface of the distal patella in the proximal trochlea, which is a characteristic of the SDP. Another interesting fact is that, in the VDP images, it is possible to notice an exuberant protrusion in the anterior region of the knee [8, 9]. Based on these three arguments, we therefore chose to classify our case as SDP, but we agree that

there are differences in relation to the classical SDP described in patients with osteophytes.

Clinicians must consider SDP among the differential diagnosis in patients presenting to the ED with direct trauma to the knee associated with anterior knee pain. Initial radiographs may misdiagnose SDP as a ruptured patellar tendon [10, 11]. Careful examination is important to identify locking of the knee in full extension. Ultrasonography is a useful diagnostic aid in doubtful cases [11, 12].

Several maneuvers have been described to perform closed reduction of the patella. Usually, intra-articular administration of local anesthetics provides sufficient analgesia; however, sedation or regional blocks may be indicated. The knee should be positioned in hyperextension with the hip flexed to relax the rectus femoris muscle. The patella is secured between the operator's two digits. Gentle lateralization and medialization movements are performed until successful reduction occurs. If this maneuver does not yield results, forceful anteriorization of the patella can be attempted. However, open reduction is required in patients not responding to these aforementioned maneuvers [13, 14].

4. Conclusion

Although rare, SDP should be accurately diagnosed and treated in the ED and should not be misdiagnosed as patellar tendon rupture. SDP commonly occurs in individuals with osteoarthritis; however, young patients with direct trauma to the knee may also present with this condition.

Conflicts of Interest


The authors declare that there are no conflicts of interest regarding the publication of this article.

References


- [1] N. Jahangir and M. Umar, "Spontaneous superior patellar dislocation in young age: case report and reduction technique," *Journal of Surgical Case Reports*, vol. 2017, no. 3, p. 15, 2017.
- [2] R. K. Clift and W. El-Alami, "Superior patellar dislocation: the value of clinical examination and radiological investigation," *BMJ Case Reports*, vol. 2012, no. 1, Article ID bcr2012007571, 2012.
- [3] R. Watson-Jones, *Fractures and Joint Injuries*, Baltimore: Williams & Wilkins, Philadelphia, PA, USA, 1946.
- [4] P. W. van Egmond, M. C. Vermeulen, C. F. van Dijke, and H. C. A. Graat, "Superior dislocation of the patella: a pathognomonic finding and review of literature," *Skeletal Radiology*, vol. 46, no. 2, pp. 259–264, 2016.
- [5] A. J. Saleemi, A. Hussain, M. J. Iqbal, M. G. Thuse, and A. A. George, "Superior dislocation of patella in a rugby player: an update on a extremely rare condition and review of literature," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 15, no. 9, pp. 1112–1113, 2007.
- [6] T. Kataoka, N. Iizawa, and S. Takai, "Superior dislocation of the patella in a young woman without osteophytes: a case report," *Journal of Nippon Medical School*, vol. 83, no. 1, pp. 24–26, 2016.
- [7] O. Ofluoglu, D. Yasmin, R. Donthineni, and M. Yildiz, "Superior dislocation of the patella with early onset patellofemoral arthritis: a case report and literature review," *Knee Surgery, Sports Traumatology, Arthroscopy*, vol. 14, no. 4, pp. 350–355, 2006.
- [8] U. N. Udogwu and C. S. Sabatini, "Vertical patellar dislocation: a pediatric case report and review of the literature," *Orthopedic Reviews (Pavia)*, vol. 10, no. 3, p. 7688, 2018.
- [9] H. Ahmad Khan, A. Bashir Shah, and Y. Kamal, "Vertical patellar displacement: reduction by the push up and rotate method, a case report and literature review," *Trauma Monthly*, vol. 21, no. 5, article e24705, 2016.
- [10] H. W. Michael and J. C. Edward Jr., "Superior dislocation of the patella," *Journal of Trauma: Injury, Infection, and Critical Care*, vol. 17, no. 1, pp. 77–79, 1977.
- [11] R. S. Bassi and B. A. Kumar, "Superior dislocation of the patella; a case report and review of the literature," *Emergency Medicine Journal*, vol. 20, no. 1, pp. 97–98, 2003.
- [12] A. Boonrod, S. Sumanont, M. Boonard, and A. Boonrod, "Superior patellar dislocation misdiagnosed as patellar tendon rupture: the value of ultrasonography," *Case Reports in Orthopedics*, vol. 2016, Article ID 2037381, 4 pages, 2016.
- [13] J. P. Rao and M. A. Meese, "Irreducible superior dislocation of the patella requiring open reduction," *American Journal of Orthopedics*, vol. 26, no. 7, pp. 486–488, 1997.
- [14] S. Takai, N. Yoshino, and Y. Hirasawa, "Case report arthroscopic treatment of voluntary superior dislocation of the patella," *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, vol. 14, no. 7, pp. 753–756, 1998.



**The Scientific
World Journal**




Gastroenterology
Research and Practice



MEDIATORS
of
INFLAMMATION



Journal of
Diabetes Research



Disease Markers



Journal of
Immunology Research



PPAR Research




Hindawi

Submit your manuscripts at
www.hindawi.com




International Journal of
Endocrinology




BioMed
Research International



Journal of
Ophthalmology



Stem Cells
International




eCAM
Evidence-Based
Complementary and
Alternative Medicine




Journal of
Obesity



Journal of
Oncology




Computational and
Mathematical Methods
in Medicine




Behavioural
Neurology



Parkinson's
Disease



AIDS
Research and Treatment



Oxidative Medicine and
Cellular Longevity