Case Report

Subarachnoid Hemorrhage Secondary to Forceful Sneeze

Ali Zohair Nomani, Haris Majid Rajput, Mansoor Iqbal, Zakir Jan, Muhammad Irshad, Mazhar Badshah, and Rao Sohail Yasin Khan

Department of Neurology, Pakistan Institute of Medical Sciences, Islamabad 44080, Pakistan

Correspondence should be addressed to Ali Zohair Nomani; alin9432@gmail.com

Received 1 September 2014; Revised 27 December 2014; Accepted 28 December 2014

Academic Editor: N. Scott Litofsky

Copyright © 2015 Ali Zohair Nomani et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Subarachnoid hemorrhage (SAH) is a relatively less common but important neurological condition comprising 5% of all the cerebrovascular accidents. In most populations the reported incidence is 6-7 per 100,000 person-years and one-third of survivors become dependent. It is a serious but potentially treatable cause of neurological morbidity. Multiple authors have identified the most unusual novel associations and triggers of subarachnoid bleeds over the past decade. We herein report a rare case of subarachnoid hemorrhage leading to focal neurological deficit in a middle aged man secondary to forceful sneeze.

1. Introduction

Subarachnoid hemorrhage (SAH) is a relatively less common but important neurological condition comprising 5% of all the cerebrovascular accidents. In most populations the reported incidence is 6-7 per 100,000 person-years and one-third of survivors become dependent [1]. It has been found to be around 20 per 100,000 in Finland and Japan [1]. In India and Pakistan, incidence of SAH is still low as compared with that in Japan and the Western world. If early deaths due to SAH however are taken into consideration, the rate would turn out to be as high as 32 per 100,000 [2, 3]. Average case fatality rate has been found to be about 50% in population-based studies, with a trend towards gradual improvement. In India and Pakistan, incidence of SAH is around 5%. About 80% of the cases of SAH are due to rupture of intracranial aneurysms and the rest of 20% are nonaneurysmal [3-6].

Prevention of subarachnoid hemorrhage and indeed diagnosis and therapy requires understanding its pathophysiology. As such, it is important to know which activities can lead to SAH, as it is a serious but potentially treatable cause of neurological morbidity [7-14]. Common etiological associations include intracranial arterial aneurysms, head injuries, and arteriovenous malformations. The triggers that lead to SAH may at times be identifiable while none may be apparent at other times. A few of the recent case reports have

suggested novel causes/triggers of subarachnoid hemorrhage [9, 10, 15-23]. We herein report a rare case of intracranial subarachnoid bleed leading to focal neurological deficit in a middle aged man secondary to forceful sneeze. While authors in their review articles, cross-over studies, short communications, and case-control studies have identified sneeze as a potential trigger factor, we did not find any detailed case report describing solely sneeze leading to SAH through thorough database search at Pubmed, Medline, MedlinePlus, PubMed Central, and Pakmedinet. To our knowledge, our case is unique and the first of its kind to be reported from Pakistan.

2. Case Presentation

A 55-year-old gentleman presented to our stroke unit with 2-day history of sudden onset, severe, generalized headache following forceful sneeze. The headache occurred instantaneously following the sneeze while the patient was sitting on a chair watching television. It was the worst headache of patients' life. A consultation with local general practitioner (GP) within 10 minutes of the event revealed normal blood pressure (120/80 mm Hg) and pulse (86/min, regular) but the patient was having neckache and had 3 to 4 episodes of vomiting. This was almost instantaneously followed by right

FIGURE 1: First CT scan brain without contrast showing suspicion of subarachnoid bleed. (a) Hyperdensity involving the posterior most portion of falx cerebri and extending to right (white arrow)-(trickling trail of blood); (b) hyperdensity extending along falx cerebri posteriorly (white arrow long) and marginally to the right (white arrow short).

sided numbness and weakness of body while the patient was consulting his GP. Neither did he lose consciousness nor had he fits or fever but he had been irritable and aggressive since then. He was not addicted to any of the elicit or related drugs of abuse. Clinical examination revealed supple neck with right hemiparesis and Broca's aphasia. He was prescribed analgesics and advised Computed Tomography (CT) scan brain. He got his CT done at a local hospital on second day of event after which he was referred to our unit on the same day with suspicion of subarachnoid hemorrhage (Figure 1). We received the patient with above described findings after 46 hours of initial event and he got admitted in our stroke unit. His first CT brain was reviewed by consultant radiologist at our setup who reported the presence of trickling blood along the falx cerebri extending posteriorly and towards the right side into posterior fossa (Figure 1). In the light of suspicion, cerebrospinal fluid (CSF) examination was done to confirm the presence of xanthochromia. A subsequent CT scan brain was done on 4th day of event that revealed clear cut presence of SAH with developing hydrocephalus (Figure 2). A following CT angiography (CTA) revealed aneurysm of the right posterior communicating artery (Figure 3).

The patient was managed conservatively using analgesics, antiemetics, intravenous hydration, nimodipine, and follow-up rehabilitation program as he did not provide consent for neurosurgical intervention. The patient made an uneventful recovery with complete resolution of neurological deficit and was symptom-free on a follow-up visit at 8 weeks.

3. Discussion

Subarachnoid hemorrhage is an important cause of neurological deficits. It is mostly associated with ruptured intracranial aneurysms. The triggers responsible for rupture have been

studied by many researchers who have pointed out their significance as well as pathophysiology. Reported associations include coagulopathies (either pharmacologically-induced or resulting from systemic diseases), lumbar puncture for diagnostic or anesthesiological purposes, traumatic injuries, aortic coarctation, degenerative vascular diseases, herpes encephalitis, aneurysmal rupture following sexual intercourse, venous hypertension, cortical venous thrombosis, bee sting, epidural blood patch, Takayasu arteritis, brucella meningitis, cranial arterial dissections, contrast agent neurotoxicity, Wegner's granulomatosis, dengue fever, polyarteritis nodosa, moyamoya, and Conn's syndrome [9, 10, 23–34]. While authors have identified sneeze as a potential trigger factor, published cases describing SAH secondary to sneeze have rarely been reported. Matsuda et al. evaluated medical records of 513 patients with aneurysmal subarachnoid hemorrhage. They analyzed the factors precipitating aneurysmal rupture and found that incidence was higher between 6:00 AM to 9:00 AM and 6:00 PM to 9:00 PM when engaging in daily routines such as defecation/micturition, brushing teeth/washing face/dressing, eating/drinking, and taking a bath [35]. Vlak et al. identified independent risk factors for SAH in two of his publications including strenuous activities that resemble valsava. He identified eight triggers in a crossover study and reported that nose blowing is associated with a relative risk of 2.3 for ACA (anterior communicating artery), 6.6 for ICA (internal carotid artery), 6.2 for MCA (middle cerebral artery), 3.8 for ANT (anterior circulation: combining ACA, MCA, and ICA), and 1.9 for VBA (vertebrobasilar arteries) [36, 37]. While both suggested multiple trigger factors, none reported sneeze in their work. The most plausible explanation of SAH with sneeze relates to the cardiovascular and cerebrovascular effects of the valsava maneuver on MAP (mean arterial pressure) and ICP (intracranial pressure), thereby altering the transmural pressure gradient across a preexisting aneurysmal wall. This

FIGURE 2: Second CT scan brain without contrast showing clearly visible subarachnoid bleed. (a) Classical star shaped hyperdensity seen filling the subarachnoid space most apparently around the circle of Willis (white arrows); (b) hyperattenuating signal in bilateral sylvian fissures (white arrows); (c) trickle of blood seen extending into the ventricular system (black arrow) (also visible in (a)).

stands true for other triggers as well. Under normal circumstances, cerebral blood flow is maintained constant over a range of cerebral perfusion pressure values by cerebrovascular autoregulation while maintaining a specific narrow range of mean arterial pressure (MAP) [34, 38]. Changes in ICP and blood pressure may alter intracranial blood volume as a result of dilatation or constriction of cerebral blood vessels while volume-pressure changes and brain shifts are taking place (Figure 2). The overall result is a compromised arterial wall at the verge of rupture and aneurysms are the most vulnerable sight for this. The purpose of describing this event getting triggered by such a casual event as sneeze points out to the fact that even the simplest and apparently harmless activities can damage a potentially weak point in the cranial vasculature [38, 39].

4. Conclusion

SAH can be triggered by the simplest and apparently harmless activities if there is a potentially weak point in the cranial vasculature. As such, family screening should be duly offered

to immediate family members for unruptured, untreated intracranial aneurysms. Moreover, patients and their relatives should be educated about the possible risk factors/triggers for SAH.

Disclosure

This work has been carried out without any grants or funds. It has been completed without external financial support and the expenses whatsoever for the purpose of this study have been contributed solely by the authors themselves and no one else.

Conflict of Interests

The authors declare no conflict of interests.

Authors' Contribution

All the authors contributed equally to this work. They performed the literature search, did data collection, analyzed

FIGURE 3: CTA brain showing PCom artery aneurysm. (a) CTA: an aneurysm arising from origin of right posterior communicating artery (PCom) (white arrow); (b) CTA: the aneurysm measured 11 mm in larger dimension with a broad-base/neck of 5 mm and a dome of 3.5 mm (white arrows); (c) vascular reconstruction: right PCom aneurysm on vascular reconstruction imaging (arrowhead) with prominent ipsilateral PCom and correspondingly small/hypoplastic ipsilateral P1 of PCA (posterior cerebral artery).

the data, wrote the paper, and reviewed the final paper. All the authors meet the criteria for authorship as established by ICMJE.

References

- [1] J. van Gijn, R. S. Kerr, and G. J. Rinkel, "Subarachnoid haemorrhage," *The Lancet*, vol. 369, no. 9558, pp. 306–318, 2007.
- [2] I. A. Raja and M. A. Javaid, "Aneurysm surgery in Pakistan," *Neurologia Medico-Chirurgica*, vol. 38, supplement, pp. 134–137, 1998.
- [3] S. N. Bhagwati, "Incidence of subarachnoid hemorrhage from aneurysmal rupture in India," *Neurologia Medico-Chirurgica*, vol. 38, pp. 128–130, 1998.
- [4] M. Wasay, I. A. Khatri, and S. Kaul, "Stroke in South Asian countries," *Nature Reviews Neurology*, vol. 10, no. 3, pp. 135–143, 2014.
- [5] K. Sher, S. Shah, and S. Kumar, "Etiologic patterns of ischaemic stroke in young adults," *Journal of the College of Physicians and Surgeons Pakistan*, vol. 23, no. 7, pp. 472–475, 2013.
- [6] B. Persson, L. Toss, and P. Ortenwall, "Fatal sneezing?" *Läkartidningen*, vol. 74, no. 47, article 4222, 1977.
- [7] M. R. Reynolds, J. T. Willie, G. J. Zipfel, and R. G. Dacey Jr., "Sexual intercourse and cerebral aneurysmal rupture: potential mechanisms and precipitants: a review," *Journal of Neurosurgery*, vol. 114, no. 4, pp. 969–977, 2011.
- [8] K. W. Lee, F. Y. Tsai, and C. Y. Cheng, "Venous hypertension may be a factor in aneurysmal rupture. A case report," *Neuroradiology Journal*, vol. 26, no. 3, pp. 311–314, 2013.
- [9] R. Verma, R. Sahu, and R. Lalla, "Subarachnoid haemorrhage as the initial manifestation of cortical venous thrombosis," *BMJ Case Reports*, 2012.
- [10] E. M. Nyberg, M. Imran Chaudry, A. S. Turk, A. M. Spiotta, D. Fiorella, and R. D. Turner, "Report of two cases of a rare cause of subarachnoid hemorrhage including unusual presentation and an emerging and effective treatment option," *Journal of NeuroInterventional Surgery*, vol. 5, no. 5, p. e30, 2013.
- [11] S. Dikici, L. Y. Aydin, A. Saritas, O. Kudas, and H. Kandis, "An unusual presentation of bee sting: subarachnoid hemorrhage," *The American Journal of Emergency Medicine*, vol. 30, no. 8, pp. 1663.e5–1663.e6, 2012.
- [12] Z. I. Hasiloglu, S. Albayram, H. Ozer, D. C. Olgun, H. Selcuk, and M. Y. Kaynar, "Cranial subarachnoid hemorrhage as an unusual complication of epidural blood patch," *Clinical Neurology & Neurosurgery*, vol. 113, no. 8, pp. 689–692, 2011.

- [13] S.-J. Hyun, S.-N. Hwang, T.-K. Nam, S.-W. Park, and J.-S. Byun, "Takayasu's arteritis complicated with subarachnoid hemorrhage and hematomyelia—case report," *Neurologia Medico-Chirurgica*, vol. 51, no. 2, pp. 119–122, 2011.
- [14] Y. Tonomura, H. Kataoka, N. Yata, M. Kawahara, K. Okuchi, and S. Ueno, "A successfully treated case of herpes simplex encephalitis complicated by subarachnoid bleeding: a case report," *Journal of Medical Case Reports*, vol. 4, no. 1, article 310, 2010.
- [15] D. Valade and A. Ducros, "Acute headache in the emergency department," *Handbook of Clinical Neurology*, vol. 97, pp. 173–181, 2010.
- [16] J. K. H. Woo, D. Malfair, T. Vertinsky, M. K. S. Heran, and D. Graeb, "Intracranial transthecal subarachnoid fat emboli and subarachnoid haemorrhage arising from a sacral fracture and dural tear," *British Journal of Radiology*, vol. 83, no. 985, pp. e18–e21, 2010.
- [17] L. K. Bittencourt, F. Palma-Filho, R. C. Domingues, and E. L. Gasparetto, "Subarachnoid hemorrhage in isolated cortical vein thrombosis: a rare presentation of an unusual condition," *Arquivos de Neuro-Psiquiatria*, vol. 67, no. 4, pp. 1106–1108, 2009.
- [18] M. Tønssas, S. H. Johnsen, M. Seip, and P. Eldevik, "Diagnostiske utfordringer hos en mann med subaraknoidalblødning," *Tidsskrift for den Norske Lægeforening*, vol. 129, no. 23, pp. 2490–2492, 2009.
- [19] S.-J. Lee, Y.-Y. Lin, C.-W. Hsu, S.-J. Chu, and S.-H. Tsai, "Intraventricular hematoma, subarachnoid hematoma and spinal epidural hematoma caused by lumbar puncture: an unusual complication," *American Journal of the Medical Sciences*, vol. 337, no. 2, pp. 143–145, 2009.
- [20] H. Korri, A. Awada, Y. Ali, and J. Choucair, "Brucellar meningitis complicated by aneurysmal subarachnoid hemorrhage," *Revue Neurologique*, vol. 164, no. 12, pp. 1052–1055, 2008.
- [21] P. F. Hsieh, Y.-C. Lee, and M.-H. Chang, "Unilateral carotid and vertebral artery dissections and contralateral subarachnoid hemorrhage in a postpartum patient," *Acta Neurologica Taiwanica*, vol. 17, no. 2, pp. 94–98, 2008.
- [22] P. Ceruti, R. Morbio, M. Marraffa, and G. Marchini, "Simultaneous bilateral acute angle-closure glaucoma in a patient with subarachnoid hemorrhage," *Journal of Glaucoma*, vol. 17, no. 1, pp. 62–66, 2008.
- [23] C. Bradford and A. McElduff, "An unusual cause of hypocalcaemia: magnesium induced inhibition of parathyroid hormone secretion in a patient with subarachnoid haemorrhage," *Critical Care and Resuscitation*, vol. 8, no. 1, pp. 36–39, 2006.
- [24] M. Domenicucci, A. Ramieri, S. Paolini et al., "Spinal subarachnoid hematomas: our experience and literature review," *Acta Neurochirurgica*, vol. 147, no. 7, pp. 741–750, 2005.
- [25] G. J. Kaptain, G. Lanzino, and N. F. Kassell, "Subarachnoid haemorrhage: epidemiology, risk factors, and treatment options," *Drugs Aging*, vol. 17, no. 3, pp. 183–199, 2000.
- [26] S. Sharp, J. Stone, and R. Beach, "Contrast agent neurotoxicity presenting as subarachnoid hemorrhage," *Neurology*, vol. 52, no. 7, pp. 1503–1505, 1999.
- [27] D. N. Cruz and A. S. Segal, "A patient with Wegener's granulomatosis presenting with a subarachnoid hemorrhage: case report and review of CNS disease associated with Wegener's granulomatosis," *American Journal of Nephrology*, vol. 17, no. 2, pp. 181–186, 1997.
- [28] O. Patey, L. Ollivaud, J. Breuil, and C. Lafaix, "Unusual neurologic manifestations occurring during dengue fever infection," *American Journal of Tropical Medicine and Hygiene*, vol. 48, no. 6, pp. 793–802, 1993.
- [29] C. W. Barton, "Subarachnoid hemorrhage presenting as acute chest pain: a variant of le coup de poignard," *Annals of Emergency Medicine*, vol. 17, no. 9, pp. 977–978, 1988.
- [30] C. Simonetti and F. Bechini, "Subarachnoid hemorrhage as initial manifestation of polyarteritis nodosa," *Rivista di Neurologia*, vol. 58, no. 5, pp. 180–182, 1988.
- [31] M. D. du Trevou and J. R. van Dellen, "Moyamoya disease as a cause of subarachnoid haemorrhage. A case report," *South African Medical Journal*, vol. 71, no. 11, pp. 725–727, 1987.
- [32] A. Bryer and J. L. Miller, "Conn's syndrome presenting as a subarachnoid haemorrhage. A case report," *South African Medical Journal*, vol. 62, no. 8, pp. 249–250, 1982.
- [33] D. W. Oxley, "An unusual tear-gas gun fatality," *Journal of Forensic Sciences*, vol. 22, no. 3, pp. 606–609, 1977.
- [34] J. N. Walton, "Subarachnoid haemorrhage of unusual aetiology," *Neurology*, vol. 3, no. 7, pp. 517–543, 1953.
- [35] M. Matsuda, K. Watanabe, A. Saito, K.-I. Matsumura, and M. Ichikawa, "Circumstances, activities, and events precipitating aneurysmal subarachnoid hemorrhage," *Journal of Stroke and Cerebrovascular Diseases*, vol. 16, no. 1, pp. 25–29, 2007.
- [36] M. H. M. Vlak, G. J. E. Rinkel, P. Greebe, J. G. van der Bom, and A. Algra, "Trigger factors for rupture of intracranial aneurysms in relation to patient and aneurysm characteristics," *Journal of Neurology*, vol. 259, no. 7, pp. 1298–1302, 2012.
- [37] M. H. M. Vlak, G. J. E. Rinkel, P. Greebe, and A. Algra, "Risk of rupture of an intracranial aneurysm based on patient characteristics: a case-control study," *Stroke*, vol. 44, no. 5, pp. 1256–1259, 2013.
- [38] J. L. Trimble and D. R. Kockler, "Statin treatment of cerebral vasospasm after aneurysmal subarachnoid hemorrhage," *The Annals of Pharmacotherapy*, vol. 41, no. 12, pp. 2019–2023, 2007.
- [39] A. L. Kühn, J. S. Balam, and I. Q. Grunwald, "Current management and treatment of cerebral vasospasm complicating SAH," *CNS and Neurological Disorders—Drug Targets*, vol. 12, no. 2, pp. 233–241, 2013.