Case Report

Fallopian Tube Herniation: An Unusual Complication of Surgical Drain

Lipi Sharma, Alpana Singh, Sruthi Bhaskaran, A. G. Radhika, and Gita Radhakrishnan

UCMS and GTBH, Dilshad Garden, New Delhi 110095, India

Correspondence should be addressed to Lipi Sharma, lip.sharma14@gmail.com

Received 29 March 2012; Accepted 24 June 2012

Academic Editors: J. C. Canterino and K. Nasu

Copyright © 2012 Lipi Sharma et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Surgical drains have been used since time immemorial, but their use is not without complications. By presenting this case we aim to describe an uncommon complication of herniation of fallopian tube following the simple procedure of surgical drain removal. **Case Presentation.** This case describes a 23-year G2P1L1 who underwent an emergency cesarean section for obstructed labor with intraperitoneal drain insertion. The patient had an uneventful postoperative period, drain was removed on day 4, and she was discharged. She presented on day 8 with the complaint of soakage of drain site dressing. On examination an edematous, tubular structure with early sign of necrosis was seen coming out of drain site and a provisional diagnosis of appendix herniation was made. On emergency laparotomy fallopian tube was seen coming out through the drain site and salphingectomy was done. **Conclusion.** Drains are not a substitute for good surgical technique. Although herniation of intestine, omentum, appendix, gall bladder, and ovary have been reported, we could not find any case of fallopian tube herniation in the literature searched by us.

1. Background

Surgeons use drain to prevent intra-abdominal fluid collection, both therapeutically and prophylactically. Surgical drains, as useful as they are, have been noted not to be without complications; such as, hemorrhage, infection, tissue damage, pain, blockage, and herniation of viscera [1]. In the case illustrated below, fallopian tube (a rare possibility) herniated out of surgical drain site, which due to delay in detection lead to salphingectomy.

2. Case Report

A 23-year-old, unbooked, G2P1L1 presented to the obstetric emergency at term with history of pain in abdomen since morning and leaking per vaginum since 6 hrs. On examination she was of average built. Her pulse was 106/min and blood Pressure was 100/70 mm Hg. On per abdominal examination her uterus corresponded to 34 weeks size, the lower uterine segment was stretched out, and fetus was in transverse lie. The uterus was having mild contractions and fetal heart sound was absent. On per vaginum examination

vagina was hot and dry, cervix was 5 cms dilated, fully effaced, and membranes were absent with fetal hand felt in vagina. An emergency caesarean section was done in view of transverse lie with hand prolapse in obstructed labor with intrauterine fetal demise.

Intraoperatively it was found that due to obstruction uterine musculature had become edematous. Due to continuous ooze from lower uterine segment, a drain was inserted from a separate incision in right iliac fossa and placed intraperitoneally in pouch of Douglas. The patient developed mild cough on day 2; otherwise her postoperative period was uneventful. The drain output had reduced to 25 mL by day 4, drain was removed and she was discharged with advice to come on day 8 for stitch removal. At the time of stitch removal patient complained that her drain site dressing was soaked. On examination a 4 × 5 cm pinkish brown, tubular, edematous mass with slough at base was seen coming out through drain site which had become strangulated with early changes of necrosis. It was nontender and nonreducible. The patient gave history of normal bowel movement. On examination her vitals were stable, bowel sounds were present, and there were no signs of peritonitis. Following surgical opinion

FIGURE 1: Right Fallopian tube: swollen, edematous with early signs of necrosis.

patient was taken for emergency laparotomy on suspicion of intestinal or appendicular herniation from drain site.

Intraoperatively the herniating viscus was the right fallopian tube which has become necrosed due to strangulation, appendix and gut were normal. Right salpingectomy was done and tissue was sent for histopathology. The histopathology reported it as fallopian tube with submucosal edema, congestion, and features of acute inflammation (Figure 1). Postoperative period was uneventful and patient was discharged.

3. Discussion

Drains have been used in surgical practice since time immemorial. Indications for surgical drain use may be therapeutic (to evacuate existing collection of fluid) or prophylactic (to prevent the collection of fluid). This fluid may be blood, bile, pus, urine, serum/lymph, pancreatic secretion, and bowel anastomotic leaks. Various reports of fallopian tube as a content of hernia sac (inguinal and femoral hernia) [2], and herniation of fallopian tube posthysterectomy (abdominal and vaginal) [3] are present. There are reports of herniation of intestine [4], appendix [5], omentum [6], gall bladder [7] (single report), and ovary [8] (single report) from surgical drain site but, on reviewing indexed English literature we could not find any report of fallopian tube herniation. What lead to fallopian tube herniation from surgical drain site in this patient and was it preventable?

The incidence of herniation of viscera from port site in laparoscopic surgery is reported to be .65%–2.5% [9]. Do the same factors govern herniation from surgical drain site following open surgery? In our patient a passive, closed Penrose drain (no. 32) was placed prophylactically in pouch of Douglas. Large meta-analyses [10] have revealed that the indications of prophylactic drains should be minimized in case of uncomplicated surgeries. The drain used had side holes which do not have any influence on drainage but lead to tissue entanglement.

The drain used (Figure 2) had an external diameter of 10 mm. It has been reported that herniation of viscera increases with increase in port size ≥ 10 mm [11]. Where ever possible fascial defects of ≥ 10 mm should be closed. The patient developed cough on postoperative day 3. Recurrent

FIGURE 2: Drainage tube 32 FG.

increase in intra-abdominal pressure caused by coughing or straining, prolonged surgery, poor nutrition, wound infection, obesity, and steroid use are known to cause poor healing and herniation [12]. Also, the large size of puerperal uterus had placed the fallopian tube close to the drain site which may be one of the factors which lead to its herniation.

Wrong technique of insertion and removal can be a causative factor but it was not so in this case. We used asymmetrical method which causes peritoneal stretching for insertion of drain rather than using direct stab incision. While removing, gradual sustained pressure should be used to withdraw the drain.

4. Conclusion

Surgical drains, although used commonly in elective as well as emergency surgeries, are not without complications. Such complications can be avoided by more restricted use of surgical drains.

References

- [1] J. G. Mosley and G. Jantet, "Herniation at the site of an abdominal drain," *British Journal of Clinical Practice*, vol. 32, no. 2, pp. 56–58, 1978.
- [2] O. V. Ozkan, E. Semerci, E. Aslan, S. Ozkan, K. Dolapcioglu, and E. Besirov, "A right sliding indirect inguinal hernia containing paraovarian cyst, fallopian tube, and ovary: a case report," *Archives of Gynecology and Obstetrics*, vol. 279, no. 6, pp. 897–899, 2009.
- [3] G. B. Candiani and M. Candiani, "Posthysterectomy fallopian tube herniation: a report of two cases," *Journal of Reproductive Medicine for the Obstetrician and Gynecologist*, vol. 41, no. 12, pp. 915–920, 1996.
- [4] S. Kulkarni, B. Krijgsman, D. Sharma, and A. V. Kaisary, "Incarcerated small bowel hernia through drain site," *Annals of the Royal College of Surgeons of England*, vol. 86, no. 6, pp. W24–W25, 2004.
- [5] D. C. O'Riordan, L. F. Horgan, and B. R. Davidson, "Drain-site herniation of the appendix," *British Journal of Surgery*, vol. 82, no. 12, p. 1628, 1995.
- [6] F. M. Howard and T. R. Sweeney, "Omental herniation after operative laparoscopy: a case report," *Journal of Reproductive*

Medicine for the Obstetrician and Gynecologist, vol. 39, no. 5, pp. 415–416, 1994.

- [7] B. Vedat, S. Aziz, and K. Çetin, “Evisceration of gallbladder at the site of a Pezzer drain: a case report,” *Cases Journal*, vol. 2, no. 7, article 8601, 2009.
- [8] P. Pianon and M. Lise, “Exteriorization of an ovary: an unusual complication of abdominal drainage,” *British Journal of Surgery*, vol. 79, no. 9, p. 963, 1992.
- [9] H. Tonouchi, Y. Ohmori, M. Kobayashi, and M. Kusunoki, “Trocarr site hernia,” *Archives of Surgery*, vol. 139, no. 11, pp. 1248–1256, 2004.
- [10] K. S. Gurusamy, K. Samraj, P. Mullerat, and B. R. Davidson, “Routine abdominal drainage for uncomplicated laparoscopic cholecystectomy,” *Cochrane Database of Systematic Reviews*, no. 3, article CD006004, 2007.
- [11] A. Dulskas, R. Lunevicius, and J. Stanaitis, “A case report of incisional hernia through a 5 mm lateral port site following laparoscopic cholecystectomy,” *Journal of Minimal Access Surgery*, vol. 7, no. 3, pp. 187–189, 2011.
- [12] A. Loh and P. A. Jones, “Evisceration and other complications of abdominal drains,” *Postgraduate Medical Journal*, vol. 67, no. 789, pp. 687–688, 1991.