Case Report

Spontaneous Pneumomediastinum in Labor

Mohamed Adnane Berdai, Said Benlamkadem, Smael Labib, and Mustapha Harandou

Obstetric and Pediatric Intensive Care Unit, Hassan II University Hospital, Sidi Mohamed Ben Abdellah University, Fez, Morocco

Correspondence should be addressed to Mohamed Adnane Berdai; adnane.berdai@yahoo.fr

Received 23 December 2016; Accepted 31 January 2017; Published 21 February 2017

Academic Editor: Ian MacKenzie

Copyright © 2017 Mohamed Adnane Berdai et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Spontaneous pneumomediastinum and subcutaneous emphysema also known as Hamman's syndrome is a very rare complication of labor that is often related to the valsalva maneuver during the labor. In most case, Hamman's syndrome is a self-limiting condition, rarely complicated unless there are underlying respiratory diseases. Chest X-ray can be a useful early diagnostic technique in severe clinical presentation. We report an uneventful pregnancy in a primigravid parturient, which was complicated in the late second stage of labor by the development of subcutaneous emphysema, pneumomediastinum, and mild pneumothorax. Spontaneous recovery occurred after four days of conservative management. This condition shows the major interest of labor analgesia especially locoregional techniques.

1. Introduction

Labor related pneumomediastinum and subcutaneous emphysema is also known as Hamman's syndrome, it is named after the physician who first described pneumomediastinum in association with subcutaneous emphysema during pregnancy in 1945 [1]. We report the case of an uneventful pregnancy in a primigravid parturient, which was complicated in the late second stage of labor by the development of Hamman's syndrome.

2. Case Report

A 22-year-old nulliparous, with no medical history, presented to our hospital in established labor after a spontaneous rupture of membranes, at 40 weeks of pregnancy. The first stage of labor lasted for 6 hours. During labor, she refused to receive any form of analgesia. At the end of the second stage of labor which had lasted for 2 hours, the patient began to cough; she also complained of shortness of breath and pain over the chest and the neck regions.

Physical examination revealed conscious patient; the blood pressure was 120/90 mmHg, pulse was 96/min, and the oxygen saturation was 98% on 3 L/min of oxygen. The patient had mild dyspnea with breath rate of 24 cycles/minutes.

No edema was present and the patient was afebrile; the cardiopulmonary auscultation was normal.

Palpation showed crepitus underneath the skin of the anterior neck and bilateral subclavian regions consistent with subcutaneous emphysema. The patient had not undergone any radiological examination due to the imminent delivery and the stability of the respiratory function. She received oxygen by facial mask and remained stable. 15 minutes later, the labor ended with a vaginal delivery requiring an episiotomy. The patient delivered a healthy, live infant with Apgar score of 10/10 in the first and fifth minute. One hour later, the crepitations extended to the upper anterior chest wall and to the face including the eyelids without any deterioration in respiratory status. A chest X-ray showed pneumomediastinum with subcutaneous emphysema extending to the soft tissue of the axillary region. Computed tomography (Figures 1 and 2) confirmed pneumomediastinum and subcutaneous emphysema extending to the thoracic, cervical, and abdominal regions. It showed also a minimal pneumothorax. There were no anomalies in the pulmonary parenchyma and thoracic great vessels. Routine blood tests and arterial gas were found to be normal.

The patient had been followed up for 4 days in hospital, without a specific medication. The symptoms have regressed

FIGURE 1: Thoracic computed tomography showing pneumomediastinum, subcutaneous emphysema, and minimal pneumothorax.

FIGURE 2: Cervical computed tomography showing extension of subcutaneous emphysema to the cervical region.

and chest X-ray showed disappearance of pneumomediastinum. The patient had been discharged home. Regular follow-up showed complete recovery without any sequelae.

3. Discussion

The estimated incidences of subcutaneous emphysema and pneumomediastinum in labor and delivery are between 1/2000 and 1/10000 [2]. The pathophysiology involves rupture of marginal alveoli into perivascular tissue planes, with trapping of air into the mediastinum [3]. It is related to the Valsalva maneuver during the labor that increases intrathoracic pressure associated with coughing, vomiting, screaming, or pushing in labor. The increased intrathoracic pressure, combined with decreased vascular calibre, establishes a pressure gradient into the vascular sheath, where air can dissect into the mediastinum and from there into the subcutaneous tissues along fascial planes [4]. Spontaneous pneumomediastinum can be seen in situations other than

Hamman's syndrome: forceful coughing related to asthmatic bronchospasm or infections, physical activity, and vomiting [5].

Despite being unproved, some suggest that Hamman's syndrome is most likely to occur in a long labor [6]. However, there was no evidence to substantiate this. A large analysis of 187 cases of Hamman's syndrome, with and without evidence of pneumothorax, revealed that 95% of women were primiparous and both first and second stages of labor were of normal duration. Average fetal size was also found to be within normal limits [7].

The onset of symptoms of pregnancy-related pneumomediastinum usually develops during labor; however, clinical appearance is often delayed until the postpartum phase [8]. These symptoms have also been reported in the earlier antepartum period, during hyperemesis or even spontaneously at rest [9]. The symptoms include chest pain, dysphagia, dysphonia, dyspnea, cough, palpitations, anxiety, and hemoptysis [10]. A chest X-ray usually reveals the subcutaneous emphysema, and significant pneumomediastinum or pneumopericardium may be visible.

In our case, the patient was primiparous and had regular length of the first and second stages of labor. She delivered with an important pushing effort as she had not any form of analgesia.

Pneumothorax occurs due to a sudden rise in mediastinal pressure with rupture of the parietal mediastinal pleura. It is reported in a third of cases [11]. However, an isolated pneumothorax can occur in patients with predisposing lung disease.

The importance of awareness of this condition, which is usually self-limiting, is to be able to distinguish it from other more life-threatening postpartum conditions such as pulmonary embolism, amniotic fluid embolism, aortic dissection, and myocardial infarction [10].

Hamman's syndrome is a self-limiting condition, rarely complicated, unless there are underlying respiratory diseases. In the absence of pneumothorax that can cause cardiorespiratory collapse, Hamman's syndrome can resolve with bed rest or even with a conservative treatment within 2 weeks [12]. Treatment is supportive and based on administration of oxygen and provision of analgesia. Management includes avoidance of factors likely to exacerbate the condition such as active pushing, the use of nitric oxide and positive pressure ventilation, and attention to the reassurance and comfort of the patient [13]. Regional anaesthesia is preferred for caesarean section, in order to avoid positive pressure ventilation [14].

4. Conclusion

Hamman's syndrome is a rare complication of labor. It is often a benign condition that is treated conservatively and most cases have a self-limiting course. This condition shows the major interest of labor analgesia especially locoregional techniques.

Competing Interests

The authors declare no conflict of interests.

References

- [1] L. Hamman, "Mediastinal emphysema: the frank billings lecture," *Journal of the American Medical Association*, vol. 128, no. 1, pp. 1–6, 1945.
- [2] N. Houari, S. Labib, M. Berdai, and M. Harandou, "Pneumomédiastin et emphysème sous-cutané du post-partum : une nouvelle observation," *Annales Françaises d'Anesthésie et de Réanimation*, vol. 31, no. 9, pp. 728–730, 2012.
- [3] S. E. Weinberger and S. T. Weiss, "Pulmonary diseases," in *Medical Complications During Pregnancy*, G. N. Burrow and T. P. Duffy, Eds., pp. 363–400, WB Saunders, Philadelphia, Pa, USA, 5th edition, 1999.
- [4] M. Marc Bonin, "Hamman's syndrome (spontaneous pneumomediastinum) in a parturient: a case report," *Journal of Obstetrics and Gynaecology Canada*, vol. 28, no. 2, pp. 128–131, 2006.
- [5] E. A. Panacek, A. J. Singer, B. W. Sherman, A. Prescott, and W. F. Rutherford, "Spontaneous pneumomediastinum: clinical and natural history," *Annals of Emergency Medicine*, vol. 21, no. 10, pp. 1222–1227, 1992.
- [6] E. M. Karson, D. Saltzman, and M. R. Davis, "Pneumomediastinum in pregnancy: two case reports and a review of the literature, pathophysiology, and management," *Obstetrics and Gynecology*, vol. 64, no. 3, pp. 39–43, 1984.
- [7] S. R. Reeder, "Subcutaneous emphysema, pneumomediastinum, and pneumothorax in labor and delivery," *American Journal of Obstetrics and Gynecology*, vol. 154, no. 3, pp. 487–489, 1986.
- [8] S. Majer and P. Graber, "Postpartum pneumomediastinum (Hamman's syndrome)," *Canadian Medical Association Journal*, vol. 177, no. 1, p. 32, 2007.
- [9] J. E. Heffner and S. A. Sahn, "Pleural disease in pregnancy," *Clinics in Chest Medicine*, vol. 13, no. 4, pp. 667–678, 1992.
- [10] S. Kandiah, H. Iswariah, and S. Elgey, "Postpartum pneumomediastinum and subcutaneous emphysema: two case reports," *Case Reports in Obstetrics and Gynecology*, vol. 2013, Article ID 735154, 3 pages, 2013.
- [11] M. De Swiet, "Diseases of the respiratory system," in *Medical Disorders in Obstetric Practice*, pp. 1–32, Blackwell Science, Oxford, UK, 3rd edition, 1995.
- [12] S. Kouki and A. A. Fares, "Postpartum spontaneous pneumomediastinum 'Hamman's syndrome,'" *BMJ Case Reports*, vol. 2013, 2013.
- [13] Y. Jayran-Nejad, "Subcutaneous emphysema in labour," *Anaesthesia*, vol. 48, no. 2, pp. 139–140, 1993.
- [14] B. L. Duffy, "Post partum pneumomediastinum," *Anaesthesia and Intensive Care*, vol. 32, no. 1, pp. 117–119, 2004.