Case Report

Ovarian Torsion after Hysterectomy: Case Report and Concise Review of the Reported Cases

Demetrio Larraín , Andrés Casanova, and Iván Rojas

Department of Obstetrics and Gynecology, Clínica Santa María, Santiago, Chile

Correspondence should be addressed to Demetrio Larraín; dlarraind@gmail.com

Received 10 April 2018; Accepted 14 June 2018; Published 4 July 2018

Academic Editor: Maria Grazia Porpora

Copyright © 2018 Demetrio Larraín et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ovarian torsion after hysterectomy is a rare event. The diagnosis of ovarian torsion is challenging because symptoms are nonspecific. We present a case of ovarian torsion 2 years after laparoscopic hysterectomy (LH). Furthermore, we performed a literature review about ovarian torsion after hysterectomy. This case shows that, in cases of acute onset pelvic pain in patients with history of hysterectomy, the adnexal torsion must be kept in mind in the differential diagnosis, especially in those women who had undergone LH.

1. Introduction

Ovarian torsion accounts for 2-3% of all acute gynecological emergencies. It remains a clinically difficult diagnosis as the symptoms are usually nonspecific. Ovarian torsion can occur at any age and also after hysterectomy [1]. Although hysterectomy with ovarian conservation is not a risk factor for torsion [1], it seems to be more frequent after laparoscopic hysterectomy [2]. Therefore, despite it is a rare event, its prevalence could increase in the future with the widespread use of laparoscopic approach. We present a case of ovarian torsion after a laparoscopic hysterectomy and performed a literature review about reported cases.

2. Case Presentation

A 41-year-old woman, gravida 3, para 3, was admitted to our institution with a 12-hour history of acute onset pelvic pain, nausea, and vomiting. She had undergone total laparoscopic hysterectomy 2 years previously. The abdominal exam revealed mild distention and tenderness over the right lower quadrant. Vaginal examination revealed exquisite pain in the right vaginal fornix and the finding of a painful adnexal mass in the rectovaginal pouch of Douglas. Transvaginal ultrasonography showed a 60-mm cystic lesion in the right ovary with moderate ascites. We performed an exploratory

laparoscopy and found a right adnexal torsion (Figure 1) and a right adnexectomy was successfully performed. Since the left ovary was normal a left ovariopexy was also performed.

3. Discussion

Ovarian torsion after hysterectomy is a rare event with a prevalence of 7.91/1000 hysterectomies [2]. Although hysterectomy is not a risk factor for ovarian torsion [1], it has been estimated that approximately 8% of adnexal torsions occur in patients with previous hysterectomy [3, 4]. To date, there are no data on how different hysterectomy techniques may affect the risk of future ovarian torsion. However, several cases of ovarian torsion have been reported after laparoscopic hysterectomy (LH) (Table 1), while to our knowledge, only one case has been published after abdominal approach [6]. This could be explained by the fact that laparoscopic approach has been associated with both fewer postoperative adhesions [7] and less adhesion-related complications [8] when compared to laparotomy, in both gynecologic and pelvic surgery. The latter could be a direct consequence of the lesser peritoneal trauma and less inflammatory response during laparoscopy [9, 10]. Moreover, our technique of LH [11] includes a wide fenestration of the broad ligament, which is left open after surgery. Based on our observations,

TABLE 1: Ovarian torsion after hysterectomy.

Author, year [Reference]	Cases	Time from hysterectomy	Type of hysterectomy	Symptoms
Mashiach, 2004 [2]	7	2.64 years	Laparoscopic	Pelvic pain
Houry, 2001 [3]	7	NA	NA	NA
Lo, 2008 [4]	5	NA	NA	NA
Cieberta, 2016 [5]	1	7 months	Laparoscopic supracervical	Asymptomatic suspicious pelvic mass
Elhjouji, 2015 [6]	1	4 years	Abdominal	Pelvic pain

NA: not available.

FIGURE 1: Laparoscopic view of a twisted right ovary. Note the absence of adhesions, which may have facilitated torsion.

the ovaries remain much more movable after LH when compared to open approach (due to the skeletonization of infundibulopelvic ligament). For that reason, we perform prophylactic oophoropexy after hysterectomy only when the infundibulopelvic ligament has been excessively skeletonized and the ovaries remain too much mobile. However, in agreement with other authors [12, 13], we perform systematic oophoropexy in cases of recurrent torsion, excessive length of utero-ovarian ligament, torsion of a solitary adnexa, or contralateral pexy in case of adnexectomy of the twisted adnexa.

This case shows that, in cases of acute onset pelvic pain in patients with history of hysterectomy, the adnexal torsion must be kept in mind in the differential diagnosis, especially in those women who had undergone LH. In addition, we encourage that, during LH with ovarian conservation, the fenestration of the broad ligament should be performed in the direction of the uterine artery and not towards the infundibulopelvic ligament, in order to keep the ovary more fix to the pelvic side-wall.

Ethical Approval

The retrospective observational nature of the study did not necessitate the local Institutional Ethics Committee approval. Institutional Review Board exemption was obtained.

Consent

Written informed consent was obtained from the patient for publication of the case report and accompanying images.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this article.

References

- [1] V. Asfour, R. Varma, and P. Menon, "Clinical risk factors for ovarian torsion," *Journal of Obstetrics & Gynaecology*, vol. 35, no. 7, pp. 721–725, 2015.
- [2] R. Mashiach, M. Canis, K. Jardon, G. Mage, J.-L. Pouly, and A. Wattiez, "Adnexal torsion after laparoscopic hysterectomy: description of seven cases," *The Journal of the American Association of Gynecologic Laparoscopists*, vol. 11, no. 3, pp. 336–339, 2004.
- [3] D. Houry and J. T. Abbott, "Ovarian torsion: A fifteen-year review," *Annals of Emergency Medicine*, vol. 38, no. 2, pp. 156–159, 2001.
- [4] L.-M. Lo, S.-D. Chang, S.-G. Horng, T.-Y. Yang, C.-L. Lee, and C.-C. Liang, "Laparoscopy versus laparotomy for surgical intervention of ovarian torsion," *Journal of Obstetrics and Gynaecology Research*, vol. 34, no. 6, pp. 1020–1025, 2008.
- [5] M. Cieberta, A. Baran, A. Słabuzewska-Jóźwiak, and G. Jakiel, "Case report of ovarian torsion mimicking ovarian cancer as an uncommon late complication of laparoscopic supracervical hysterectomy," *Przegląd Menopauzalny*, vol. 15, pp. 223–226, 2016.
- [6] A. Elhjouji, O. Zahdi, H. Baba et al., "Adnexal torsion after abdominal hysterectomy: a first observation," *Pan African Medical Journal*, vol. 22, article 9, 2015 (French).
- [7] H. M. Dowson, J. J. Bong, D. P. Lovell, T. R. Worthington, N. D. Karanjia, and T. A. Rockall, "Reduced adhesion formation following laparoscopic versus open colorectal surgery," *British Journal of Surgery*, vol. 95, no. 7, pp. 909–914, 2008.
- [8] M. Al-Sunaidi and T. Tulandi, "Adhesion-related bowel obstruction after hysterectomy for benign conditions," *Obstetrics & Gynecology*, vol. 108, no. 5, pp. 1162–1166, 2006.
- [9] C. N. Gutt, T. Oniu, P. Schemmer, A. Mehrabi, and M. W. Büchler, "Fewer adhesions induced by laparoscopic surgery?" *Surgical Endoscopy*, vol. 18, no. 6, pp. 898–906, 2004.
- [10] D. Roberston, G. Lefebvre, and Clinical Practice Gynaecology Committee, "Adhesion prevention in gynaecological surgery," *Journal of Obstetrics and Gynaecology Canada*, vol. 32, pp. 598–608, 2010.
- [11] N. Bourdel, X. Tran, R. Botchorhisvili, J.-L. Pouly, M. Canis, and G. Mage, "Laparoscopic hysterectomy in 10 steps," *Journal de Chirurgie*, vol. 146, no. 5, pp. 483–491, 2009 (French).
- [12] N. Fuchs, N. Smorgick, Y. Tovbin et al., "Oophoropexy to prevent adnexal torsion: how, when and for whom?" *Journal of Minimally Invasive Gynecology*, vol. 17, no. 2, pp. 205–208, 2010.

- [13] K. Jardon, R. Botchorishvili, B. Rabischong et al., "How I perform.... an ovariopexy after adnexal torsion," *Gynécologie Obstétrique & Fertilité*, vol. 34, no. 6, pp. 529-530, 2006 (French).