Case Report

A Vaginal Angiomyofibroblastoma as a Rare Cause of a Prolapsing Vaginal Mass: A Case Report and Review of the Literature

Harriet Calvert^{1,2}, Supuni Kapurubandara^{3,4,5}, Yogesh Nikam^{3,4,5},
Raghwa Sharma^{5,6,7} and Anita Achan⁶

¹Department of Obstetrics and Gynaecology, The Maitland Hospital, Maitland, NSW, Australia

²Department of Obstetrics and Gynaecology, John Hunter Hospital, Newcastle, NSW, Australia

³Department of Obstetrics and Gynaecology, Westmead Hospital, Sydney, NSW, Australia

⁴Sydney West Advanced Pelvic Surgery (SWAPS), Sydney, NSW, Australia

⁵University of Sydney, Sydney, NSW, Australia

⁶Department of Tissue Pathology and Diagnostic Oncology, Institute of Clinical Pathology and Medical Research, Westmead Hospital, Sydney, NSW, Australia

⁷Western Sydney University, Sydney, NSW, Australia

Correspondence should be addressed to Harriet Calvert; harriet.calvert@hnehealth.nsw.gov.au

Received 27 December 2017; Accepted 21 March 2018; Published 29 April 2018

Academic Editor: Yoshio Yoshida

Copyright © 2018 Harriet Calvert et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction. Angiomyofibroblastoma (AMFB) is a rare, benign, mesenchymal cell tumour which presents as a slow-growing mass. It is most commonly seen in the vulva and is often mistaken for Bartholin's abscess. It is histologically diagnosed by the presence of stromal cells intermingled with small blood vessels. It is morphologically similar to cellular angiofibroma and aggressive angiofibroma, the latter of which is locally invasive and has a possibility of metastasis and a high risk of local recurrence. There is one reported case of an AMFB undergoing sarcomatous transformation. **Case Report.** We report a case of a multiparous, 36-year-old woman with an anterior vaginal mass which was inappropriately treated as a vaginal prolapse prior to definitive surgical management. This is only the second reported case of an AMFB presenting as a prolapsing mass.

1. Introduction

Angiomyofibroblastoma (AMFB) is a rare, benign, mesenchymal tumour that most commonly occurs as a slow-growing mass in the vulva, first described in 1992 [1]. It is often misdiagnosed as Bartholin's gland cyst [1, 2].

This type of solid tumour has also less commonly been described in the vagina and the inguinoscrotal region of men.

It is most prevalent in women in the reproductive age group with a mean age of 45 and varies in size (from 0.5 to 23 cm) but is usually less than 5 cm [1–4].

Histologically, it has a defined border and is characterised by alternating hypo- and hypercellular areas with numerous blood vessels [1, 5].

2. Case Report

We describe a case of a 36-year-old multiparous (G3P2) woman who presented with an acute episode of pelvic pain. She was referred to a general gynaecological clinic after ultrasound findings revealed a 4.1 cm complex left ovarian cyst suggestive of an endometrioma.

She also reported a 2-year history of a bulge that protruded from her vagina and was associated with discomfort and dyspareunia and occasionally required digital reduction especially with tampon use. She had been diagnosed with a vaginal prolapse by a gynaecology clinic at another institution.

FIGURE 1: Sagittal section of MRI showing the 45 mm × 50 mm solid mass in the vesicovaginal septum in relation to the uterus, mass, and vagina (V).

FIGURE 2: Ultrasound scan showing relation of mass to uterus and vagina.

Her past medical history consisted of migraines with aura, exercise induced asthma, and a family history of breast cancer (half-sister). She had never had a PAP smear.

On bimanual examination, a well-delineated solid mass was found on the anterior vaginal wall in the midline, measuring 5 cm by 5 cm. There was no evidence of pelvic organ prolapse with good support of the uterus, posterior wall, and anterior wall above the mass. The cervix was visualised anteriorly and there was no evidence of cervical excitation. A routine PAP smear was performed with difficulty secondary to the vaginal mass.

With respect to investigations, Ca 125 was 29 U/mL giving a low relative malignancy index. A repeat ultrasound scan demonstrated a 2.9 cm left ovarian cyst, suggestive of an endometrioma and a solid mass inferior to the uterus and anterior to the vagina, displacing the bladder (Figure 2).

On Magnetic Resonance Imaging, a 45 mm × 50 mm solid mass in the vesicovaginal septum with a well-defined margin was demonstrated (Figure 1). The mass was displacing the bladder anteriorly and displacing the urethra towards the left of the midline. T2 imaging showed a predominantly hypointense, heterogenous signal with areas of hyperintensity. There was mild enhancement after gadolinium injection. Close to the external urethral orifice, the interface between the mass and the urethra was ill defined. Evidence of a left ovarian endometrioma and endometriosis deposits were seen elsewhere in the pelvis.

FIGURE 3: View of mass on diagnostic laparoscopy; uterus anteverted.

These MRI findings suggested that the mass was either endometriosis with surrounding reactive fibrous and smooth muscle proliferation, neoplasm, or an infection relating to a urethral diverticulum. After a multidisciplinary meeting with a urogynaecologist, the patient underwent an examination under anaesthesia, diagnostic laparoscopy, cystoscopy, excision of endometriosis, and excision of the vaginal mass.

The vaginal mass was removed with laparoscopic assessment via a midline incision on the anterior vaginal wall with lateral dissection around the cystic structure (Figures 3–6). A cystoscopy and urethroscopy suggested no involvement and the cyst was enucleated. Multiple haemostatic sutures were needed with surgical snare to achieve haemostasis and the defect was closed. A repeat cystoscopy and urethroscopy showed no injury.

Histopathological macroscopic assessment of the mass showed pale tan tissue surrounded by a thin capsule and on sectioning a homogeneous whorled tan tissue (Figure 6). Microscopically the low power photomicrographs showed a well-circumscribed border. It comprised collagenised areas of epithelioid to spindle cells with small to thin walled arborizing vessels. Aggregation of cells around vessels was noted and there were no atypical mitoses, necrosis, or atypia (Figures 7 and 8).

The immunohistochemistry showed positive desmin, SMA, CD34, and vimentin. The cells displayed high intensity nuclear positivity for progesterone and oestrogen receptors. These findings were consistent with a diagnosis of angiofibroma.

3. Discussion

AMFB is a very rare benign, mesenchymal tumour with less than 100 cases previously having been reported in the literature. There has been a reported age range of 17–86 with a mean age at presentation of 45 [2, 6, 7].

It commonly presents as a painless, slow-growing, vulval mass and is most commonly diagnosed as Bartholin's cyst or abscess (46%) or a lipoma (15%) [2]. There is only one other reported case of it presenting as a prolapsing vaginal mass [8]. There is often a delay in diagnosis with a mean duration of 29 months between initial symptoms and diagnosis [2, 6].

AMFB is morphologically similar to other invasive mesenchymal cell tumours such as aggressive angiofibroma (AAM) and cellular angiofibroma and they share many

FIGURE 4: Marsupialisation of vaginal mass from anterior vaginal wall.

FIGURE 5: Excision of vaginal mass from anterior vaginal wall.

FIGURE 6: Cut surface of mass.

overlapping immunohistochemical and structural features [9, 10].

It is diagnostically challenging to differentiate between AMFB and AAM but important due to the latter's locally invasive nature, the possibility of metastasis, and the high risk of local recurrence [11, 12]. AMFB can be diagnosed by a higher cellularity, distinct border, plump stromal cells, increased presence of small blood vessels, and a lesser degree of stromal myxoid change [6]. Other differential diagnoses include cellular angiofibroma and vulvovaginal myofibroblastoma. Cellular angiofibromas are uniformly cellular with thick-walled, hyalinised blood vessels without surrounding aggregation of epithelioid or plasmacytoid cells. Adipocytes

are often found in the periphery [10]. Vulvovaginal myofibroblastomas characteristically contain ovoid, spindle, or stellate cells in a variety of architectural patterns. They also do not have the perivascular aggregates seen in AMFB [10]. Both cellular angiofibromas and myofibroblastomas exhibit the loss of RB1 and FOXO1A1 genes due to the deletion of the 13q14 chromosomal region. This typical loss of genetic material is not found in AMFB [13].

Immunohistologically, AMFB tumours have been found to be strongly positive for vimentin, positive for desmin, and to a lesser degree alpha-smooth muscle actin. Staining is rarely useful in differentiating between tumour types [13]. The stromal cells are characteristically positive for oestrogen and progesterone receptors, suggesting a hormonal role in the development of the tumour [14].

There have only been 5 previous reports of MRI findings of an AMFB. All report a mass with well-defined margins and as in our case they have been found to appear as a heterogeneous signal intensity on T2-weighted MRI. All other cases reported fast and persistent enhancement on dynamic gadolinium-enhanced MRI whereas ours showed only mild enhancement [15, 16].

The other studies found a mass with homogeneous intermediate signal intensity on T2 weighted MRI [7, 17, 18].

Ultrasound has been reported to be useful in assessing heterogeneity, vascularity, and delineating infiltration and relation to surrounding structures [7, 19].

It is widely accepted that AMFB can be treated with wide local excision with clear margins. There has only been one case report of a benign local recurrence. This was a pedunculated mass 5 × 3 cm arising from the vaginal vault which was excised with clear margins. Upon follow-up 14 months later 3 small, nodular growths were found close to the site of excision on the anterior and posterior vaginal walls which, when excised, showed the same features as the previous tumour with no transformation [8].

There has also been reported one case of previously diagnosed AMFB undergoing sarcomatous change. A 13 cm vulval mass was resected which showed many accepted features of an AMFB however did show focal sarcomatous change at the resected margin. At 2 years, the mass had recurred at the same site and resection demonstrated a 14 cm mass comprised of only the high-grade sarcomatous component with vascular invasion that was not previously present [20].

Another reported case of a locally invasive recurrence of AMFB at 2 years after resection was due to a misdiagnosed AAM on the original specimen [21]. The local recurrence rate of AAM after clear margin resection has been reported to be up to 47% [22–25].

4. Conclusion

The majority of AMFB occur in the vulva, most commonly presenting as a painless mass.

Vaginal AMFB are rarer and may present later with dyspareunia, awareness of a vaginal mass, or an incidental finding on exam [26–28]. Wide local excision is the recommended treatment, with enough surrounding tissue to enable

FIGURE 7

FIGURE 8: Cross section showing alternating hypercellular areas around blood vessels (H) and hypocellular (O) areas containing slender collagen fibrils with no evidence of necrosis.

the pathologist to differentiate between AMFB and the locally infiltrative AAM.

MRI and US can be useful imaging modalities depending on location of the tumour. Due to the rarity of cases, there are no recommendations on long-term monitoring but due to the reported instances of tumour recurrence and sarcomatous transformation we suggest that follow-up should be considered until at least 2 years postoperatively [8, 20].

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

- [1] C. D. M. Fletcher, W. Y. W. Tsang, C. Fisher, K. C. Lee, and J. K. C. Chan, "Angiomyofibroblastoma of the vulva: A benign neoplasm distinct from aggressive angiofibroma," *The American Journal of Surgical Pathology*, vol. 16, no. 4, pp. 373–382, 1992.
- [2] S. M. Sims, K. Stinson, F. W. McLean, J. D. Davis, and E. J. Wilkinson, "Angiomyofibroblastoma of the vulva: A case report of a pedunculated variant and review of the literature," *Journal of Lower Genital Tract Disease*, vol. 16, no. 2, pp. 149–154, 2012.
- [3] J. Seo, K. Lee, N. Yoon, J. Lee, B. Kim, and D. Bae, "Angiomyofibroblastoma of the vulva," *Obstetrics & Gynecology Science*, vol. 56, no. 5, pp. 349–351, 2013.
- [4] K. Nagai, K. Aadachi, and H. Saito, "Huge pedunculated angiofibroblastoma of the vulva," *International Journal of Clinical Oncology*, vol. 15, no. 2, pp. 201–205, 2010.
- [5] M. R. Nucci and C. D. M. Fletcher, "Vulvovaginal soft tissue tumours: update and review," *Histopathology*, vol. 36, no. 2, pp. 97–108, 2000.
- [6] G. P. Nielsen and R. H. Young, "Mesenchymal tumors and tumor-like lesions of the female genital tract: a selective review with emphasis on recently described entities," *International Journal of Gynecological Pathology*, vol. 20, no. 2, pp. 105–127, 2001.
- [7] B. Wolf, L.-C. Horn, R. Handzel, and J. Eichenkel, "Ultrasound plays a key role in imaging and management of genital angiofibroblastoma: A case report," *Journal of Medical Case Reports*, vol. 9, no. 1, article no. 248, 2015.
- [8] M. M. Saleh, A. H. Yassin, and M. S. Zaklama, "Recurrent angiofibroblastoma of the vagina: a case report," *Eur J Gynaecol Oncol*, vol. 28, no. 4, p. 324, 2007.
- [9] F. Alameda, A. Munné, T. Baró et al., "Vulvar angiofibroma, aggressive angiofibroma, and angiofibroblastoma: An immunohistochemical and ultrastructural study," *Ultrastructural Pathology*, vol. 30, no. 3, pp. 193–205, 2006.
- [10] W. G. McCluggage, "A review and update of morphologically bland vulvovaginal mesenchymal lesions," *Int J Gynecol Pathol*, vol. 24, no. 1, pp. 26–38, 2005.
- [11] R. M. Siassi, T. Papadopoulos, and K. E. Matzel, "Metastasizing aggressive angiofibroma," *The New England Journal of Medicine*, vol. 341, no. 23, article 1772, 1999.
- [12] S. Blandamura, J. Cruz, L. Faure Vergara, I. M. Puerto, and V. Ninfo, "Aggressive angiofibroma: a second case of metastasis with patient's death," *Human Pathology*, vol. 34, no. 10, pp. 1072–1074, 2003.
- [13] G. Magro, A. Righi, R. Caltabiano, L. Casorzo, and M. Michal, "Vulvovaginal angiofibroblastomas: Morphologic, immunohistochemical, and fluorescence in situ hybridization analysis for deletion of 13q14 region," *Human Pathology*, vol. 45, no. 8, pp. 1647–1655, 2014.
- [14] W. G. McCluggage and R. G. White, "Angiofibroblastoma of the vagina," *J Clin Pathol*, pp. 53–803, 2000.
- [15] J. Geng, S. Hu, and F. Wang, "Large paravaginal angiofibroblastoma: Magnetic resonance imaging findings," *Japanese Journal of Radiology*, vol. 29, no. 2, pp. 152–155, 2011.
- [16] J. L. Kyoung, H. M. Jeung, Y. Y. Dae, H. C. Ji, J. L. In, and J. M. Seon, "Angiofibroblastoma arising from the posterior

- perivesical space: A case report with MR findings," *Korean Journal of Radiology*, vol. 9, no. 4, pp. 382–385, 2008.
- [17] H. Kitamura, N. Miyao, Y. Sato, M. Matsukawa, T. Tsukamoto, and T. Sato, "Angiomyofibroblastoma of the female urethra," *International Journal of Urology*, vol. 6, no. 5, pp. 268–270, 1999.
- [18] K. J. Morteale, G. J. Lauwers, P. J. Mergo, and P. R. Ros, "Perineal angiomyofibroblastoma: CT and MR findings with pathologic correlation," *Journal of Computer Assisted Tomography*, vol. 23, no. 5, pp. 687–689, 1999.
- [19] X. Wang, H. Yang, H. Zhang, T. Shi, and W. Ren, "Transvaginal sonographic features of perineal masses in the female lower urogenital tract: A retrospective study of 71 patients," *Ultrasound in Obstetrics & Gynecology*, vol. 43, no. 6, pp. 702–710, 2014.
- [20] G. P. Nielsen, R. H. Young, G. R. Dickersin, and A. E. Rosenberg, "Angiomyofibroblastoma of the vulva with sarcomatous transformation ('angiomyofibrosarcoma')," *The American Journal of Surgical Pathology*, vol. 21, no. 9, pp. 1104–1108, 1997.
- [21] Y.-F. Wang, H.-L. Qian, and H.-M. Jin, "Local recurrent vaginal aggressive angiomyxoma misdiagnosed as cellular angiomyofibroblastoma: A case report," *Experimental and Therapeutic Medicine*, vol. 11, no. 5, pp. 1893–1895, 2016.
- [22] J. F. Fetsch, W. B. Laskin, M. Lefkowitz, L.-G. Kindblom, and J. M. Meis-Kindblom, "Aggressive angiomyxoma: a clinicopathologic study of 29 female patients," *Cancer*, vol. 78, no. 1, pp. 79–90, 1996.
- [23] L. R. Bégin, P. B. Clement, M. E. Kirk, S. Jothy, W. T. Elliott McCaughey, and A. Ferenczy, "Aggressive angiomyxoma of pelvic soft parts: a clinicopathologic study of nine cases," *Human Pathology*, vol. 16, no. 6, pp. 621–628, 1985.
- [24] B. J. Sutton and J. Laudadio, "Aggressive angiomyxoma," *Archives of Pathology & Laboratory Medicine*, vol. 136, no. 2, pp. 217–221, 2012.
- [25] I. M. Chan, E. Hon, S. W. Ngai, T. Y. Ng, and L. C. Wong, "Aggressive angiomyxoma in females: is radical resection the only option?" *Acta Obstetrica et Gynecologica Scandinavica*, vol. 79, no. 3, pp. 216–220, 2000.
- [26] R. Laiyemo, S. Disu, G. Vijaya, and B. Wise, "Post-menopausal vaginal angiomyofibroblastoma: A case report," *Archives of Gynecology and Obstetrics*, vol. 273, no. 2, pp. 129–130, 2005.
- [27] Z. N. Kavak, A. Başgöl, F. Eren, and N. Ceyhan, "Angiomyofibroblastoma of the vulva: A rare but distinct entity," *Acta Obstetrica et Gynecologica Scandinavica*, vol. 79, no. 7, pp. 612–613, 2000.
- [28] K. Banerjee, S. D. Gupta, and S. R. Mathur, "Vaginal angiomyofibroblastoma," *Archives of Gynecology and Obstetrics*, vol. 270, no. 2, pp. 124–125, 2004.