Case Report

Flexor Digitorum Accessorius Longus: Importance of Posterior Ankle Endoscopy

Jorge Pablo Batista,¹ Jorge Javier del Vecchio,² Pau Golanó,³ and Jordi Vega⁴

¹Centro Artroscópico Jorge Batista, 2446 Pueyrredón Avenue, 1st Floor, 1119 Buenos Aires, Argentina

²Favaloro Foundation, 461 Solís Street 1st floor, 1078 Buenos Aires, Argentina

³Human and Embriology Unit, Department of Experimental Pathology and Therapeutics, University of Barcelona, Spain

⁴Hospital Quirón Barcelona, Plaça d'Alfonso Comín, 08023 Barcelona, Spain

Correspondence should be addressed to Jorge Javier del Vecchio; javierdv@mac.com

Received 12 February 2015; Revised 4 April 2015; Accepted 12 April 2015

Academic Editor: Quamar Bismil

Copyright © 2015 Jorge Pablo Batista et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Endoscopy for the posterior region of the ankle through two portals is becoming more widespread for the treatment of a large number of conditions which used to be treated with open surgery years ago. The tendon of the *flexor hallucis longus* (FHL) travels along an osteofibrous tunnel between the posterolateral and posteromedial tubercles of the talus. Chronic inflammation of this tendon may lead to painful stenosing tenosynovitis. The aim of this report is to describe two cases depicting an accessory tendon which is an anatomical variation of the *flexor hallucis longus* in patients with posterior friction syndrome due to posterior ankle impingement and associated with a posteromedial osteochondral lesion of the talus. The anatomical variation (FDAL) described was a finding during an endoscopy of the posterior region of the ankle, and we have spared it by sectioning the superior flexor retinaculum only. The accessory flexor digitorum longus is an anatomical variation and should be taken into account when performing an arthroscopy of the posterior region of the ankle. We recommend this treatment on this type of injury although we admit this does not make a definite conclusion.

1. Introduction

Endoscopy for the posterior region of the ankle through two portals is becoming more widespread for the treatment of a large number of conditions which used to be treated with open surgery years ago [1–5].

No doubt this is mostly due to the significant contribution arthroscopic anatomy has meant for this particular region of the body [6, 7]. At present, it is also used for the treatment of prominent os trigonum, chronic synovitis of the *flexor hallucis longus* (FHL) [8], osteochondral lesions of the posterior region of the talus, loose bodies, subtalar arthrosis, and other less common conditions such as cystic-like bone tumors involving the talus and/or calcaneus and synovial chondromatosis.

The tendon of the *flexor hallucis longus* (FHL) travels along an osteofibrous tunnel between the posterolateral and posteromedial tubercles of the talus. Chronic inflammation

of this tendon may lead to painful stenosing tenosynovitis, typically seen in dancers [8].

Activities including forced foot flexion such as running downhill or playing soccer are a predisposing factor for injuries involving the FHL. Moreover, os trigonum, cysts, *flexor digitorum accessorius longus* (FDAL), and talar dorsal exostosis may lead to tenosynovitis [8–10].

The aim of this report is to describe a clinical case depicting an accessory tendon which is an anatomical variation of the *flexor hallucis longus* in a patient with a posterior friction syndrome due to posterior ankle impingement. The discussion is based on the endoscopic diagnosis of the accessory FDL and the treatment established.

2. Case Report

A 34-year-old male (doing recreational sports) complained of a 6-month history of pain in the posterior region of his left ankle. Standing on tip toes triggered the symptoms.

FIGURE 1: Lateral X-ray. Prominent posterior talar process.

FIGURE 2: Accessory muscle (white arrow).

On physical examination, no pain was observed during either passive or active mobilization or hallux counter resistance (both plantar and dorsal flexion). The ankle hyperplantarflexion test was positive. There was also a mild limitation in movement of the lesser toes with the foot held in dorsiflexion. The lateral X-ray view evidenced the presence of a prominent posterior talar process (Figure 1). The MRI showed a muscular intensity image located close to the posterior tibial neurovascular bundle (Figure 2).

Since conservative treatment administered for 4 months (rest from sports activities, physiotherapy, and medical treatment) failed, and as symptoms remained, a posterior arthroscopy of the ankle was conducted in order to resect the posterior prominent area of the talus and then perform soft tissue debridement.

During the procedure, a large accessory muscle belly was detected, and on dislodging it medially (Figure 3) the tendon of the FHL was identified. The muscle belly was considered an anatomical variation (FDAL) (Figure 4).

The posterior talar process was resected by using both mechanical and motor instruments (Figures 5 and 6), and the

FIGURE 3: Accessory muscle belly (black arrow).

FIGURE 4: FHL and FDAL recognized.

FIGURE 5: Resection of posterior talar process using a chisel.

superior flexor retinaculum was sectioned without involving the accessory muscle and the corresponding tendon (FDAL).

The ankle was bandaged, and active mobilization (flexion-extension) was started immediately after surgery. The patient went back to recreational sports within 3 months. At 19-month follow-up the patient did not have any related symptoms.

FIGURE 6: Complete resection.

3. Discussion

This condition (FDAL) is more common among males although the difference between males and females is not significant. It is unilateral in most cases, although nine bilateral cases have been published [10–14].

According to Bowers et al. [15] the accessory muscle body of the FHL is the second leading cause of anatomical variation occurring in the ankle. The peroneus quartus is the most common, followed by other variations such as the accessory soleus muscle, the medial calcaneal tibial muscle, and the medial fibular-calcaneal muscle.

The accessory muscle body of the FHL may be an anatomical variation identified accidentally during a posterior arthroscopy of the ankle, and it may have several clinical consequences. Two of these entities were described in our series.

Eberle et al. [9] consider that tenosynovitis involving the FHL may be the result of repeated friction of the FHL caused by the accessory muscle body of the tendon inside the tarsal tunnel. On the other hand, Wittmayer and Freed [16] suggest that the presence of an accessory muscle body of the FHL should be highly suspected when assessing the MRI scans of a patient with posterior friction syndrome. This anatomical variation has also been identified as a potential causative agent of FHL syndrome associated with a positive passive hallux dorsiflexion test as reported in this series.

Another point to consider is which approach to use when this anatomical variation is identified. Ogut and Ayhan [17] and Eberle et al. [9] have suggested open surgery and a posteromedial approach to resect the accessory fascicle of the FHL. In our series, we did not think resection was necessary, and the clinical results were similar. The accessory FDL may produce tarsal tunnel syndrome, and this concept has been validated by several authors [12, 13, 16, 18–21].

Some authors like Burks and Deheer [18] and Saar and Bell [20] prefer to resect the accessory FDL by means of open surgery and have obtained a good postoperative clinical outcome.

The accessory flexor digitorum longus is an anatomical variation and should be taken into account when performing

an arthroscopy of the posterior region of the ankle. By identifying this variation muscle structure lesions will be prevented and we will know how to treat them correctly. However, we have not been able to clearly identify the variation in our cases when we assessed the MRI scans after surgery [22].

The anatomical variation (accessory FDL) described was a finding during an endoscopy of the posterior region of the ankle, and we have spared it by sectioning the superior flexor retinaculum only.

The accessory flexor digitorum longus is an anatomical variation and should be taken into account when performing an arthroscopy of the posterior region of the ankle.

We recommend this treatment on this type of injury although we admit this does not make a definite conclusion.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors thank Gabriela Franke and Claudia Tarazona for her assistance in translating this paper. Pau Golanó actively participated in the development and writing of this case report. Unfortunately he died of a stroke during its preparation (July 23, 2014).

References

- [1] T. H. Lui, "Arthroscopy and endoscopy of the foot and ankle: indications for new techniques," *Arthroscopy*, vol. 23, no. 8, pp. 889–902, 2007.
- [2] T. H. Lui, K. B. Chan, and L. K. Chan, "Zone 2 flexor hallucis longus tendoscopy: a cadaveric study," *Foot and Ankle International*, vol. 30, no. 5, pp. 447–451, 2009.
- [3] C. N. van Dijk, P. E. Scholten, and R. Krips, "A 2-portal endoscopic approach for diagnosis and treatment of posterior ankle pathology," *Arthroscopy*, vol. 16, no. 8, pp. 871–876, 2000.
- [4] C. N. van Dijk, "Hindfoot Endoscopy," *Foot and Ankle Clinics*, vol. 11, no. 2, pp. 391–414, 2006.
- [5] C. N. van Dijk, "Hindfoot endoscopy for posterior ankle pain," *Instructional Course Lectures*, vol. 55, pp. 545–554, 2006.
- [6] P. Golanó, P. P. Mariani, M. Rodríguez-Niendenfuhr, P. F. Mariani, and D. Ruano-Gil, "Arthroscopic anatomy of the posterior ankle ligaments," *Arthroscopy*, vol. 18, no. 4, pp. 353–358, 2002.
- [7] P. Golanó, J. Vega, L. Pérez-Carro, and V. Götzens, "Ankle anatomy for the arthroscopist. Part I: the portals," *Foot and Ankle Clinics*, vol. 11, no. 2, pp. 253–273, 2006.
- [8] W. G. Hamilton, "Stenosing tenosynovitis of the flexor hallucis longus tendon and posterior impingement upon the os trigonum in ballet dancers," *Foot and Ankle*, vol. 3, no. 2, pp. 74–80, 1982.
- [9] C. F. Eberle, B. Moran, and T. Gleason, "The accessory flexor digitorum longus as a cause of Flexor Hallucis Syndrome," *Foot and Ankle International*, vol. 23, no. 1, pp. 51–55, 2002.
- [10] H. Nathan, H. Gloobe, and Z. Yosipovitch, "Flexor digitorum accessorius longus," *Clinical Orthopaedics and Related Research*, vol. 113, pp. 158–161, 1975.

- [11] Y. Y. Cheung, Z. S. Rosenberg, E. Colon, and M. Jahss, "MR imaging of flexor digitorum accessorius longus," *Skeletal Radiology*, vol. 28, no. 3, pp. 130–137, 1999.
- [12] M. Kinoshita, R. Okuda, J. Morikawa, and M. Abe, "Tarsal Tunnel Syndrome associated with an accessory muscle," *Foot and Ankle International*, vol. 24, no. 2, pp. 132–136, 2003.
- [13] D. A. Peterson, W. Stinson, and J. R. Lairmore, "The long accessory flexor muscle: an anatomical study," *Foot and Ankle International*, vol. 16, no. 10, pp. 637–640, 1995.
- [14] G. J. Sammarco and S. F. Conti, "Tarsal tunnel syndrome caused by an anomalous muscle," *The Journal of Bone and Joint Surgery—American Volume*, vol. 76, no. 9, pp. 1308–1314, 1994.
- [15] C. A. Bowers, R. W. Mendicino, A. R. Catanzariti, and E. T. Kernick, "The flexor digitorum accessorius longus—a cadaveric study," *Journal of Foot and Ankle Surgery*, vol. 48, no. 2, pp. 111–115, 2009.
- [16] B. C. Wittmayer and L. Freed, "Diagnosis and surgical management of flexor digitorum accessorius longus-induced tarsal tunnel syndrome," *Journal of Foot and Ankle Surgery*, vol. 46, no. 6, pp. 484–487, 2007.
- [17] T. Ogut and E. Ayhan, "Hindfoot endoscopy for accessory flexor digitorum longus and flexor hallucis longus tenosynovitis," *Foot and Ankle Surgery*, vol. 17, no. 1, pp. e7–e9, 2011.
- [18] J. B. Burks and P. A. Deheer, "Tarsal tunnel syndrome secondary to an accessory muscle: a case report," *Journal of Foot and Ankle Surgery*, vol. 40, no. 6, pp. 401–403, 2001.
- [19] D. E. Canter and K. J. Siesel, "Flexor digitorum accessorius longus muscle: an etiology of tarsal tunnel syndrome?" *The Journal of Foot & Ankle Surgery*, vol. 36, no. 3, pp. 226–229, 1997.
- [20] W. E. Saar and J. Bell, "Accessory flexor digitorum longus presenting as tarsal tunnel syndrome: a case report," *Foot & ankle specialist*, vol. 4, no. 6, pp. 379–382, 2011.
- [21] G. J. Sammarco and M. M. Stephens, "Tarsal tunnel syndrome caused by the flexor digitorum accessorius longus. A case report," *The Journal of Bone & Joint Surgery—American Volume*, vol. 72, no. 3, pp. 453–454, 1990.
- [22] W. R. Buschmann, Y. Cheung, and M. H. Jahss, "Magnetic resonance imaging of anomalous leg muscles: accessory soleus, peroneus quartus and the flexor digitorum longus accessorius," *Foot and Ankle*, vol. 12, no. 2, pp. 109–116, 1991.