Case Report

A Lip Lump: An Unexpected Histological Diagnosis of a Lip Schwannoma

Thomas Haigh,¹ John Raad Glore,¹ David Goulesbrough,¹ and Winson Wong²

¹Bradford Teaching Hospitals NHS Foundation Trust, Bradford BD9 6RJ, UK

²Bradford Teaching Hospitals NHS Foundation Trust, University of Leeds, UK

Correspondence should be addressed to Thomas Haigh; t.haigh@hotmail.co.uk

Received 20 September 2016; Revised 1 November 2016; Accepted 17 November 2016; Published 7 February 2017

Academic Editor: Andrea Gallo

Copyright © 2017 Thomas Haigh et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Schwannomas are benign nerve sheath tumours arising from Schwann cells. They comprise 1% of all benign tumours. In the 2016 World Health Organisation Classification of Central Nervous System, they are classified as a tumour of the cranial and paraspinal nerves, Schwannoma 9560/0. A 23-year-old Caucasian lady presented with a seven-month history of a painless right upper lip lump. Examination revealed a small cystic 0.5 cm diameter lesion within the right upper lip. The clinical impression was that of a mucocele. Excision of the lip lesion was performed under local anaesthetic. Histological examination of the excised lesion demonstrated a circumscribed nodule consisting of spindle cells mixed with vascular spaces containing red blood cells and fibrin. Immunohistochemistry for S100 was strongly positive. The findings were consistent with that of a small benign schwannoma. The current consensus is for surgical excision of a conservative nature with no need for margins. If recurrence does occur one needs to consider whether complete enucleation was achieved or whether malignant transformation has occurred.

1. Introduction

Lip lumps can present to both the general practitioner in the community and the specialist in a general ENT clinic following referral. In the case presented the unexpected, rare diagnosis only became clear following excision and histological examination.

2. Case Report

A 23-year-old Caucasian lady presented with a seven-month history of a painless right upper lip lump that she caught whilst eating. She possessed no red flag symptoms and no significant medical history apart from being a smoker with poor dentition.

Examination revealed a small cystic 0.5 cm diameter lesion within the right upper lip. Excision of the lip lesion was performed under local anaesthetic. A piece of tissue measuring $7 \times 3 \times 2$ mm was sent for histological staining and examination.

3. Diagnosis

3.1. Schwannoma. In the 2016 World Health Organisation Classification of Central Nervous System, they are classified as tumour of the cranial and paraspinal nerves, Schwannoma 9560/0 [1].

The diagnosis was made following histological examination of the specimen which demonstrated cytologically bland spindle cells in hypercellular Antoni A and hypocellular Antoni B patterns (Figure 1). Scattered dilated and focally thrombosed blood vessels were visualised which are a feature of these lesions. Staining for S100 immunoperoxidase was strongly positive which is characteristic of schwannoma (Figure 2).

4. Discussion

Schwannomas are benign nerve sheath tumours arising from Schwann cells. They comprise 1% of all benign tumours [2]. The lip is the least likely oral location with only twenty

FIGURE 1: High-power slide of schwannoma: H + E $\times 400$. The tumour comprises cytologically bland spindle cells with no mitotic activity or necrosis. Focal palisading of the nuclei is seen as a characteristic of a schwannoma. This has led to the production of the so-called Verocay bodies.

FIGURE 2: High power slide of schwannoma: S100 (immunoperoxidase) staining $\times 200$. High-power view of the lesion demonstrating the characteristic diffuse positive staining with S100 protein.

documented cases. Despite their rarity, schwannomas should be considered in the differential diagnosis of any lip lump owing to the lip's neural innervation.

In terms of preoperative diagnosis and estimating margins, ultrasound, CT, and MRI can be used but are not considered routine or necessary [3]. In our case no such imaging was performed. Diagnosis is typically made following adequate biopsy. The histological section demonstrated a circumscribed nodule consisting of spindle cells mixed with vascular spaces containing red blood cells and fibrin (Figure 1). No cytological atypia or mitotic activity was noted. Immunohistochemistry for S100 was strongly positive (Figure 2). Findings were consistent with that of a small benign schwannoma. The nodule appeared to be circumscribed and intact suggesting that complete excision had been achieved.

Despite the reassuring histology, lip schwannoma can be histologically misleading and can resemble malignancy [4–6]. However, only three cases of recurrent malignant lip schwannoma have been documented [7–9].

The current consensus is for surgical excision of a conservative nature with no need for margins with the aim being to preserve the nerve of origin [4]. Complete excision should not lead to recurrence. If recurrence does occur one needs

to consider whether complete enucleation was achieved or if malignant transformation is a possibility.

Competing Interests

The authors declare no conflict of interests.

References

- [1] D. N. Louis, H. Ohgaki, O. D. Wiestler, and W. K. Cavenee, "World Health Organization histological classification of tumours of the central nervous system," in *Proceedings of the International Agency for Research on Cancer (IARC '16)*, Lyon Cedex, France, 2016.
- [2] L. F. M. da Silva, B. G. Duarte, B. A. Boiça, H. V. Rocha-Junior, and C. L. Pereira-Stabile, "Intraoral schwannoma: a case report," *Oral and Maxillofacial Surgery*, vol. 17, no. 4, pp. 319–321, 2013.
- [3] J.-I. Asaumi, H. Konouchi, and K. Kishi, "Schwannoma of the upper lip: ultrasound, CT, and MRI findings," *Journal of Oral and Maxillofacial Surgery*, vol. 58, no. 10, pp. 1173–1175, 2000.
- [4] C. C. Humber, M. A. Copete, and F. I. Hohn, "Ancient schwannoma of upper lip: case report with distinct histologic features and review of the literature," *Journal of Oral and Maxillofacial Surgery*, vol. 69, no. 6, pp. e118–e122, 2011.
- [5] L. V. Ackerman and F. H. Taylor, "Neurogenous tumors within the thorax. A clinicopathological evaluation of forty-eight cases," *Cancer*, vol. 4, no. 4, pp. 669–691, 1951.
- [6] Y. C. Chu, Y. M. Yoon, H. S. Man, J. Y. Han, J. M. Kim, and I. S. Park, "Malignant transformation of intrathoracic ancient neurilemmoma in a patient without Von Recklinghausen's disease," *Journal of Korean Medical Science*, vol. 18, no. 2, pp. 295–298, 2003.
- [7] G. Grevers and J. Röcken, "Recurrent malignant schwannoma of the lower lip: report of a case," *Otolaryngology—Head and Neck Surgery*, vol. 113, no. 1, pp. 138–139, 1995.
- [8] Y. Hashhiba, S. Nozaki, K. Yoshizawa, N. Noguchi, K. Nakagawa, and E. Yamamoto, "Recurrent multinodular neurilemmoma of the female upper lip," *International Journal of Oral and Maxillofacial Surgery*, vol. 36, no. 2, pp. 171–173, 2007.
- [9] R. Jurkovic, P. Stanko, S. Galbavy, G. Sieberova, and P. Babal, "Recurrent malignant epitheloid schwannoma of the lower lip," *Bratislava Medical Journal*, vol. 110, no. 2, pp. 116–119, 2009.