Case Report

Liesegang-Like Rings in Lactational Changes in the Breast

Mohd T. Islam, Joyce J. Ou, Katrine Hansen, Rochelle A. Simon, and M. Ruhul Quddus

Department of Pathology, The Warren Alpert Medical School of Brown University and Women & Infants Hospital,
101 Dudley Street Providence, RI 02905, USA

Correspondence should be addressed to M. Ruhul Quddus, mqquddus@wihri.org

Received 20 June 2012; Accepted 13 September 2012

Academic Editors: G. Adonakis and P. Tosi

Copyright © 2012 Mohd T. Islam et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Liesegang-like rings (LR) are periodic structures with equally spaced radial striations formed by a process that involves diffusion, nucleation, flocculation or precipitation, and supersaturation. Being more common *in vitro*, on rare occasions also reported *in vivo* in association with inflammatory or cystic lesions and confused with parasites or calcification on needle aspirates. The current paper documents that LR may be seen in noncystic and noninflammatory changes of the breast.

1. Introduction

Liesegang rings (LRs) are periodic precipitation zones from supersaturated solutions in colloidal systems. The LRs are possibly formed by a process that involves diffusion, nucleation, flocculation or precipitation, and supersaturation. Notable examples include LRs of calcium carbonate in oölitic limestone (in nature), silver chromate in gelatin (*in vitro*), and LRs of glycoprotein in pulmonary corpora amylacea (*in vivo*) [1]. The LRs vary greatly in shape and size, measuring 7–800 microns.

Being more commonly seen *in vitro*, LRs have also rarely been reported *in vivo*—mostly in association with lesions of the kidney and synovium, however, their presence has also been described in the conjunctiva, eyelid, paranasal sinus, peritoneum, and in association with endometriosis [1–3]. Isolated cases of LRs have been reported associated with breast cysts and mammary duct ectasia [4, 5].

These rings usually form in association with cystic, fibrotic, inflamed, or necrotic tissues, and may be mistaken for eggs, larvae, or parasites.

We report here a case of Liesegang rings in lactational change of the breast, noted incidentally in a mastectomy specimen for breast carcinoma.

2. Case Report

A 39-year-old, G0, Hispanic woman presented with a mass in her right breast. The patient denied the use of hormone

replacement therapy, birth control pills, or substance abuse in the past. She had never conceived naturally, had failed *in vitro* fertilization, and had never breast-fed. No family history of breast, ovarian, uterine, or colon carcinoma was present. An axillary ultrasound revealed an enlarged abnormal axillary lymph node. An ultrasound-guided core needle biopsy of the breast mass revealed a moderately differentiated infiltrating duct carcinoma.

The patient opted for a simple mastectomy of the right breast, and right axillary sentinel lymph node sampling was performed. The final pathology revealed a 2.2 cm, grade 2, hormone receptor positive, Her2/neu negative, infiltrating duct carcinoma (Figure 1), with associated grade 2 *in situ* carcinoma and negative margins. One of 5 sentinel lymph nodes showed a 0.5 cm metastatic tumor deposit with extracapsular extension (Figure 1). She received systemic chemotherapy after surgery.

In addition to invasive and *in situ* breast carcinoma, the breast tissue showed proliferative fibrocystic changes as well as lactational changes. In areas of lactational change, focal dilated glands contained intraluminal, acellular structures with equally spaced radial striations; these were identified as Liesegang rings (Figure 2).

3. Discussion and Conclusions

The formation of Liesegang rings is a well-known phenomenon in the field of chemistry, occurring most commonly *in vitro*. It was first described by German biochemist

FIGURE 1: Infiltrating duct carcinoma, Nottingham grade 2 of 3. Inset: Metastatic carcinoma in the axillary lymph node (H&E; original $\times 20$).

FIGURE 2: (a) Liesegang-like rings (LRs) in slightly dilated breast ducts associated with lactational change (H&E; original $\times 10$). (b) Higher magnification of Liesegang-like rings and lactational change (H&E; original $\times 40$).

Ralph E. Liesegang as a process involving the formation of periodic precipitation zones around a central nidus. These precipitation zones are seen microscopically as concentric laminations and are formed via alternating cycles of subsaturation and supersaturation of an insoluble product produced by diffusing reactants within a colloidal matrix or gel [6]. These characteristic Liesegang rings are rarely seen *in vivo*, but when identified are usually found in association with cystic or inflammatory lesions. They may be mistaken for parasites on fine-needle aspiration or surgical specimens of hemorrhagic areas.

The exact composition of Liesegang rings is not fully understood. LR were initially believed to represent parasites, in particular adult forms of the giant kidney worm *Diocotophyma renale* (a large blood-red nematode that infects a variety of fish-eating mammals, especially mink) [1]. The rings may range from 7 to 800 microns in diameter and have uniform, pink-tan, radially striated, double walls. Multiple small rings may also be seen within a larger, dominant ring. Raso et al. [6] noted that LR displayed some morphological differences from *D. renale* when compared with specimens from animals infected naturally or experimentally with the giant kidney worm, as they lack the internal organs of such parasites and the histopathologic changes associated with such infections. Histochemical and immunoperoxidase stains for mucopolysaccharides, amyloid, and keratin are reportedly negative [6]. However, results for calcium, iron,

and hemoglobin vary in the literature [1, 6, 7], suggesting that *in vivo*, local environmental factors may create LR of differing compositions. Special stains and energy-dispersive radiographic analysis or scanning electron microscopy have revealed that some LR also contain silicon and sulfur [1], while ultrastructural analysis has revealed an electron dense core and fine fibrillary rings with a concentric and radial pattern [6, 8].

Liesegang rings are rarely present in biological systems; therefore their presence in cystic and inflamed tissues may be confused with various parasites, algae, calcifications, corpora amylacea, psammoma bodies, and the spheroid type of amyloid. In the current paper, LR were found in association with slightly dilated breast ducts, with the epithelium showing changes similar to those seen in lactation; an incidental finding in a patient with AJCC stage IIB breast carcinoma. The current case documents that LR may also be found in noncystic and noninflammatory conditions in the breast.

References

- [1] S. M. Tuur, A. M. Nelson, and D. W. Gibson, "Liesegang rings in tissue: how to distinguish Liesegang rings from the giant kidney worm, *Diocotophyma renale*," *American Journal of Surgical Pathology*, vol. 11, no. 8, pp. 598–605, 1987.
- [2] N. Sneige, J. G. Batsakis, R. A. Hawkins, and H. P. Doble, "Pseudoparasitic (Liesegang) bodies in paranasal sinus," *Journal of Laryngology and Otology*, vol. 102, no. 8, pp. 730–732, 1988.

- [3] P. B. Clement, R. H. Young, and R. E. Scully, "Liesegang rings in the female genital tract. A report of three cases," *International Journal of Gynecological Pathology*, vol. 8, no. 3, pp. 271–276, 1989.
- [4] R. K. Gupta, A. G. R. McHutchison, and R. Fauck, "Liesegang rings in a needle aspirate from a breast cyst," *Acta Cytologica*, vol. 35, no. 6, pp. 700–702, 1991.
- [5] K. Gavin, N. Banville, D. Gibbons, and C. M. Quinn, "Liesegang rings in inflammatory breast lesions," *Journal of Clinical Pathology*, vol. 58, no. 12, pp. 1343–1344, 2005.
- [6] D. S. Raso, W. B. Greene, J. L. Finley et al., "Morphology and pathogenesis of Liesegang rings in cyst aspirates: report of two cases with ancillary studies," *Diagnostic Pathology*, vol. 19, pp. 116–119, 1998.
- [7] K. L. Pegas, M. L. Edelweiss, E. Cambruzzi et al., "Liesegang rings in xanthogranulomatous pyelonephritis: a case report," *Pathology Research International*, vol. 2010, Article ID 602523, 3 pages, 2010.
- [8] N. Sneige, R. H. Dekmezian, E. G. Silva, J. Cartwright, and A. G. Ayala, "Pseudoparasitic Liesegang structures in perirenal hemorrhagic cysts," *American Journal of Clinical Pathology*, vol. 89, no. 2, pp. 148–153, 1988.