Case Report

Tramadol-Induced Mood Elevation in a Patient with No Previous Psychiatric History

Mugtaba Osman ¹ and Mashaël Mustafa²

¹Armed Forces Centre for Psychiatric Care, Armed Forces Hospital in Taif Region, Taif, Saudi Arabia

²Armed Forces Centre for Health Rehabilitation in Taif, Armed Forces Hospital in Taif Region, Taif, Saudi Arabia

Correspondence should be addressed to Mugtaba Osman; mugtaba.osman@ucdconnect.ie

Received 31 May 2018; Revised 30 July 2018; Accepted 17 September 2018; Published 1 October 2018

Academic Editor: Lut Tamam

Copyright © 2018 Mugtaba Osman and Mashaël Mustafa. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Tramadol is a powerful analgesic medication with antidepressant effects like venlafaxine. Hypomanic features were reported in patients with psychiatric history of mood disturbance when tramadol was prescribed for them. However, it is extremely rare to notice such mood-elevating effect in patients who have no previous psychiatric history. We report on the observation of a distressing mood-elevating effect for tramadol in a patient with no previous psychiatric history. We present the case of a 26-year-old female patient who developed accelerated flow of speech, overactivity, and difficulty in sleeping following intake of tramadol 50 mg dose. These symptoms resolved four hours later and recurred as she retook tramadol. The patient has no history of mood disorder or any other psychiatric comorbidity. Clinicians should exercise caution when prescribing tramadol.

1. Introduction

Tramadol is a well-known powerful centrally acting analgesic medication with established pharmacological effects on opioid systems (mainly μ -opioid receptors agonism) and nonopioid systems (5-Hydroxy Tryptamine and Noradrenaline reuptake inhibition) [1]. Tramadol is, in fact, closely related to chemical structure of all dual action antidepressants, most notably venlafaxine. Thus, tramadol and venlafaxine shared monoaminergic action leading to the tramadol effects on mood [2]. Reports of antidepressant effects for tramadol have trickled in the clinical literature either in combination with other antidepressants [3, 4] or on its own [5–7] in patients with [8] and without need for analgesia [9]. There is currently growing theoretical and clinical concern in terms of drug-drug interaction when tramadol is used concomitantly with serotonergic antidepressants [10–13].

A recent review has clearly concluded that “caution is required” if tramadol is to be prescribed for patients on antidepressants (and vice versa) with clear contraindication for the use of tramadol in patients taking Mono-Amine Oxidase Inhibitors [14]. Of note, tramadol, owing to its powerful blockade of serotonin reuptake, was reported to

cause serotonin syndrome even in the absence of concomitant use of antidepressants [15, 16].

Hypomanic features were reported in patients with psychiatric history of mood disturbance when tramadol was prescribed for them [12, 17–21]. However, it is extremely rare to notice such mood-elevating effect in patients who have no previous psychiatric history [22]. In this report we describe a case of a nonpsychiatric patient who experienced distressing mood symptoms as they start taking tramadol to control her pain.

2. Case Presentation

We present the case of a 26-year-old female patient who developed severe persistent lower back pain secondary to spondylolisthesis in the fifth lumbar vertebra. Tramadol 50 mg dose was prescribed as a nonsurgical measure to achieve satisfactory analgesia. She took one 50 mg tablet and the pain was effectively controlled. However, she almost immediately noticed accelerated flow of speech and was unable to control her desire to talk incessantly. She was also quite overactive and “on the go” for the following four hours following intake of tramadol 50 mg dose. She noticed that she did not want

to “sit still” and continued to walk to-and-fro and engaged in cleaning the house as she experienced increased energy. She was unable to sleep, although she experienced increased irritability with neither extreme happiness nor euphoria. She retook tramadol 50 mg twice 4 days and 7 days later and the same condition (overtalkativeness, overactivity, and distress) reoccurred upon both occasions lasting for exactly 4 hours each time. There was no ataxia, tremors, blurring of vision, or any other neurological signs or symptoms. She did not want to take tramadol anymore.

Notably, she had no previous psychiatric or neurological history of note. She took no psychotropic medications for any physical or psychological reasons. She had no history of illicit substance misuse or dependence.

Apart from spondylolisthesis, she suffered from urticaria and congenital optic disc tilt. She took Chlorzoxazone one tablet on request, ranitidine 150mg daily, and Desloratadine 10mg daily.

3. Discussion

A number of cases of hypomania and mania secondary to tramadol were published over the last few years [12, 17–22]. For instance, Sharma (2016) [17] described a case of female patient who developed hypomanic symptoms after taking tramadol for treatment of fibromyalgia; of note the patient had an established diagnosis of bipolar affective disorder.

Occurrence of mood symptoms after tramadol prescribing constitutes genuine concern to clinicians in terms of prediction of who would get such symptoms and how to initiate effective management for them. One recent study found that prescription of antidepressants has increased following initiation of tramadol therapy [23]. There are no studies, to the best of our knowledge, evaluating prescription of other psychotropic agents, specifically mood stabilizers, in patients who were prescribed tramadol.

What is unique in our case report is the occurrence of hypomanic-like symptoms in a patient with no history of mood disorder or any other psychiatric comorbidity. The overactivity and talkativeness in our patient were almost immediate after the intake of tramadol and they resolved within few hours afterwards. This substantially increases the likelihood of our assertion, where such hypomania-like symptoms were caused by tramadol. All other potential explanations (such as primary subthreshold mood disorder, medical condition-related mood symptoms) were extremely less likely given the history of our patient.

Mood-elevation secondary to tramadol treatment could be regarded as unwanted extension of its antidepressant effect. Many neurological pathways were proposed to have some involvement in the antidepressant effect for tramadol. In addition to serotonergic and noradrenergic receptors, tramadol could exert an antidepressant effect by its ability to bind to D2 and D3 dopaminergic receptors [24, 25] and imidazoline I2 receptors [24, 26]. Many other mechanisms implicated in the antidepressant effects of tramadol were investigated. In a recent experiment Jesse et al. [27] found evidence to suggest that the signalling pathway that encompass L-arginine-nitric oxide-cyclic guanosine monophosphate is

involved in the antidepressant action of tramadol. An earlier experiment found that tramadol caused substantial reduction in the alpha-2 adrenergic receptors in all brain areas, like the effects of mirtazapine and mianserin [28]. Also, other experiments indicated that tramadol causes the alpha-1 adrenergic receptors to increase in density in the brain cortex and results in upregulation of dopaminergic receptors in the nucleus accumbens which has important functions in motivational behaviour [29].

Clinical trials' results indicate that 7% of patients taking tramadol would experience delirium-like Central Nervous System stimulation with a wide range of physical, neurological, and emotional presentations [30]. Emotional presentations recorded are mostly symptoms of hyperactive delirium and include anxiety, sleep difficulties, emotional lability, euphoria, and confusion [31, 32]. However, the precise mechanism of development of such unwanted idiosyncratic adverse reaction to tramadol remains largely unknown. Also, the proportion of patients with no previous psychiatric history who developed such emotional effects, although believed to be quite small, was unclear.

We report on the observation of a distressing mood-elevating effect for tramadol in a patient with no previous psychiatric history. This will, of course, make it prudent for clinicians to exercise caution when prescribing tramadol. Also, we recommend close follow-up after initiation of tramadol treatment, in order to screen for mood symptoms in this cohort of patients.

We also recommend that longitudinal studies be carried out to robustly evaluate the prevalence and predictors of occurrence of mood symptoms in patients using tramadol.

Consent

We thank the patient for providing informed consent for the publication of this case study.

Conflicts of Interest

We declare no conflicts of interest in terms of the publication of this case report.

References

- [1] M. Vazzana, T. Andreani, J. Fanguiero et al., “Tramadol hydrochloride: pharmacokinetics, pharmacodynamics, adverse side effects, co-administration of drugs and new drug delivery systems,” *Biomedicine & Pharmacotherapy*, vol. 70, pp. 234–238, 2015.
- [2] J. S. Markowitz and K. S. Patrick, “Venlafaxine-tramadol similarities,” *Medical Hypotheses*, vol. 51, no. 2, pp. 167–168, 1998.
- [3] D. Szkutnik-Fiedler, K. Kus, M. Balcerkiewicz et al., “Concomitant use of tramadol and venlafaxine - Evaluation of antidepressant-like activity and other behavioral effects in rats,” *Pharmacological Reports*, vol. 64, no. 6, pp. 1350–1358, 2012.
- [4] R. R. Reeves and S. K. Cox, “Similar effects of tramadol and venlafaxine in major depressive disorder,” *Southern Medical Journal*, vol. 101, no. 2, pp. 193–195, 2008.
- [5] N. A. Shapira, M. L. Verduin, and J. D. DeGraw, “Treatment of Refractory Major Depression With Tramadol Monotherapy,”

- Journal of Clinical Psychiatry*, vol. 62, no. 3, pp. 205-206, 2001.
- [6] M. O. Rojas-Corrales, J. Gibert-Rahola, and J. A. Micó, "Tramadol induces antidepressant-type effects in mice," *Life Sciences*, vol. 63, no. 12, pp. PL175-PL180, 1998.
 - [7] A. Rougemont-Bücking, F. Gamma, and J. Panksepp, "Use of tramadol in psychiatric care: A comprehensive review and report of two cases," *Swiss Medical Weekly*, vol. 147, 2017.
 - [8] T. Tetsunaga, T. Tetsunaga, M. Tanaka, and T. Ozaki, "Efficacy of tramadol-acetaminophen tablets in low back pain patients with depression," *Journal of Orthopaedic Science*, vol. 20, no. 2, pp. 281-286, 2015.
 - [9] J. Barber, "Examining the use of tramadol hydrochloride as an antidepressant," *Experimental and Clinical Psychopharmacology*, vol. 19, no. 2, pp. 123-130, 2011.
 - [10] B. J. Mason and K. H. Blackburn, "Possible serotonin syndrome associated with tramadol and sertraline coadministration," *Annals of Pharmacotherapy*, vol. 31, no. 2, pp. 175-177, 1997.
 - [11] D. Mittino, M. Mula, and F. Monaco, "Serotonin syndrome associated with tramadol-sertraline coadministration," *Clinical Neuropharmacology*, vol. 27, no. 3, pp. 150-151, 2004.
 - [12] A. Gonzalez-Pinto, H. Imaz, J. L. P. De Heredia, M. Gutierrez, and J. A. Micó, "Mania and tramadol-fluoxetine combination," *The American Journal of Psychiatry*, vol. 158, no. 6, pp. 964-965, 2001.
 - [13] L. E. Peacock and F. Wright, "Serotonin syndrome secondary to tramadol and citalopram," *Age and Ageing*, vol. 40, no. 4, p. 528, 2011.
 - [14] S. H. Park, R. C. Wackernah, and G. L. Stimmel, "Serotonin syndrome: Is it a reason to avoid the use of tramadol with antidepressants?" *Journal of Pharmacy Practice*, vol. 27, no. 1, pp. 71-78, 2014.
 - [15] J. Takeshita and M. H. Litzinger, "Serotonin syndrome associated with Tramadol," *Primary Care Companion to the Journal of Clinical Psychiatry*, vol. 11, no. 5, p. 273, 2009.
 - [16] A. Tashakori and R. Afshari, "Tramadol overdose as a cause of serotonin syndrome: a case series," *Clinical Toxicology*, vol. 48, no. 4, pp. 337-341, 2010.
 - [17] V. Sharma, "Tramadol-induced hypomania and serotonin syndrome," *The Primary Care Companion for CNS Disorders*, vol. 18, no. 6, 2016.
 - [18] N. Ansermot, O. Chocron, F. Herrera, and C. B. Eap, "Severe manic episode associated with tramadol in a patient with recurrent depressive disorder," *Journal of Clinical Psychopharmacology*, vol. 35, no. 2, pp. 203-204, 2015.
 - [19] K.-J. Chen, M.-L. Lu, and W. W. Shen, "Tramadol-related psychosis in a patient with bipolar I disorder," *Acta Neuropsychiatrica*, vol. 27, no. 2, pp. 126-128, 2015.
 - [20] B. Oronsky and R. Martin, "Intolerance to tramadol may herald bipolar disease," *International Journal of Family Practice*, vol. 6, no. 2, 2007.
 - [21] A. P. John and R. Koloth, "Severe serotonin toxicity and manic switch induced by combined use of tramadol and paroxetine," *Australian & New Zealand Journal of Psychiatry*, vol. 41, no. 2, pp. 192-193, 2007.
 - [22] D. Ceylan, M. Kaçar, and H. Ulas, "Manic episodes associated with tramadol: a case report," *J Clin Psychopharmacol*, vol. 35, no. 1, pp. 111-113, 2015.
 - [23] T. Chen, L. Chen, and R. Knaggs, "Prevalence of antidepressants prescribed to tramadol users in the UK primary care setting - a prescription sequence symmetry analysis," *Value in Health*, vol. 18, no. 7, p. A661, 2015.
 - [24] M. N. Chowta, M. Manjunath, H. N. Gopalakrishna, and P. Gokul, "Evaluation of role of noradrenergic system in the antidepressant activity of tramadol using tail suspension test in Albino mice," *Journal of Pharmacology and Pharmacotherapeutics*, vol. 2, no. 4, pp. 281-282, 2011.
 - [25] M. Filip, K. Wydra, S. Yalcin Inan, M. Dziedzicka-Wasylewska, and E. Przeglasiński, "Opioid and monoamine systems mediate the discriminative stimulus of tramadol in rats," *European Journal of Pharmacology*, vol. 498, no. 1-3, pp. 143-151, 2004.
 - [26] C. R. Jesse, E. A. Wilhelm, C. F. Bortolatto, and C. W. Nogueira, "Evidence for the involvement of the noradrenergic system, dopaminergic and imidazoline receptors in the antidepressant-like effect of tramadol in mice," *Pharmacology Biochemistry & Behavior*, vol. 95, no. 3, pp. 344-350, 2010.
 - [27] C. R. Jesse, C. F. Bortolatto, L. Savegnago, J. B. T. Rocha, and C. W. Nogueira, "Involvement of l-arginine-nitric oxide-cyclic guanosine monophosphate pathway in the antidepressant-like effect of tramadol in the rat forced swimming test," *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, vol. 32, no. 8, pp. 1838-1843, 2008.
 - [28] A. Faron-Górecka, M. Kuśmider, S. Y. Inan, J. Siwanowicz, and M. Dziedzicka-Wasylewska, "Effects of tramadol on α 2-adrenergic receptors in the rat brain," *Brain Research*, vol. 1016, no. 2, pp. 263-267, 2004.
 - [29] A. Faron-Górecka, M. Kuśmider, S. Y. Inan, J. Siwanowicz, T. Piwowarczyk, and M. Dziedzicka-Wasylewska, "Long-term exposure of rats to tramadol alters brain dopamine and α 1-adrenoceptor function that may be related to antidepressant potency," *European Journal of Pharmacology*, vol. 501, no. 1-3, pp. 103-110, 2004.
 - [30] P. P. Gleason, R. F. Frye, and T. O'Toole, "Debilitating reaction following the initial dose of tramadol," *Annals of Pharmacotherapy*, vol. 31, no. 10, pp. 1150-1152, 1997.
 - [31] http://www.druglib.com/druginfo/tramadol/side-effects_adverse-reactions/, accessed on July 30th, 2018.
 - [32] A. Agrawal, S. Diwan, and R. Mahajan, "Severe delirium following single dose of tramadol," *Indian Journal of Medical Sciences*, vol. 63, no. 2, pp. 80-81, 2009.