

Case Report

Subthreshold Autism Spectrum in a Patient with Anorexia Nervosa and Behçet's Syndrome

L. Dell'Osso, B. Carpita , I. M. Cremone, F. Mucci, A. Salerni, D. Marazziti, C. Carmassi, and C. Gesi

Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy

Correspondence should be addressed to B. Carpita; barbara.carpita1986@gmail.com

Received 4 March 2020; Revised 29 May 2020; Accepted 8 June 2020; Published 15 June 2020

Academic Editor: Lut Tamam

Copyright © 2020 L. Dell'Osso et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently, increasing research stressed the presence of subthreshold autistic traits in patients with other psychiatric conditions. In this framework, a significant relationship between anorexia nervosa (AN) and the autism spectrum has been frequently reported, in particular among female samples, to the point that AN has been hypothesized to be a female phenotype of autism spectrum disorder (ASD). On the other hand, among subjects with ASD has been reported a higher prevalence of immune diseases and altered immune functions. While these reports seem to support an association between neurodevelopmental and immune system alterations in ASD, the relationship between the immune system and the broader autism spectrum, including its subthreshold manifestations, remains poorly investigated. In this report, we described the presence of autistic traits in a male inpatient with AN and separation anxiety disorder, who also show a diagnosis of Behçet's syndrome (BS). This case seems to further stress the association between AN and the autism spectrum, which may not be limited to the female gender. Moreover, it further suggests a deeper link between neurodevelopmental and immune system alterations. Implications are discussed in light of the more recent neurobiological and psychopathological hypothesis about the autism spectrum.

1. Introduction

Autism spectrum disorder (ASD) is characterized by a cluster of symptoms (impairment in communication and social interaction, restricted pattern of interests, and repetitive behaviors) associated with a neurodevelopmental alteration, usually manifesting in early childhood [1]. Recently, increasing interest has been paid to investigating the presence of subthreshold autistic traits, providing support to the hypothesis that a subthreshold autism spectrum may represent a dimension spanning across several mental disorders [1]. The interest in investigating autistic traits lies in the fact that a growing number of studies are stressing their role as a vulnerability factor towards the development of different kinds of psychiatric disorders, as well as suicidal thoughts and behaviors [1–9]. Moreover, milder forms of ASD, without intellectual impairment and with less evident social difficulties, may often remain underdiagnosed, coming to clinical attention only during adult life, when patients develop other

disorders in comorbidity [1, 2]. In this framework, a significant relationship of anorexia nervosa (AN) with both ASD and autistic traits has been reported in the literature [10, 11]. Although ASD has been considered a disorder with a strikingly higher prevalence among males, recently several authors questioned this data [12, 13]. Evidence from literature seems to point out that gender differences in ASD prevalence may be related to a male bias in defining ASD criteria [10–13]. ASD is likely to be more prevalent than previously expected among females, although in this group, it would manifest with different kinds of symptoms [10–13]. Firstly, females with ASD frequently show a higher ability to cope with social difficulties by the use of camouflaging strategies, such as imitating others' behaviours, and, as a consequence, milder forms of ASD in females would remain more frequently undiagnosed [14]. Moreover, females with ASD often show a different pattern of restricted interests, such as spending time with animals, fictions, and also focusing on food and diet [10–13, 15]. Since the '80s, the association of

ASD or subthreshold autistic traits with AN has gained increasing interest in the scientific literature. After the first studies which reported a higher presence of AN among female relatives of ASD patients, several researches were conducted on the prevalence of autistic-like features among females with AN, also with a longitudinal design, confirming the presence of significant overlap between the autism spectrum and AN [10–13]. In light of these findings, some authors also hypothesized that AN should be considered as a specific presentation of ASD, which would be more frequent among females [10–13]. However, most of the studies about AN and the autism spectrum (ASD or subthreshold autistic traits) were conducted in samples composed almost exclusively by females, and no study specifically focused on the relationship between ASD and AN among males [12, 13]. In addition, few studies investigated gender differences in clinical presentations of AN, mostly reporting only a higher tendency towards a chronic course among females, while no significant difference was found for other clinical features [16]. The presence of a lower focus on thinness and a higher focus on other features (such as large shoulders and narrow waist) among males was not confirmed by all the studies [17].

On the other hand, among subjects with ASD has been reported a higher prevalence of immune diseases, as well as findings of altered immune functions [18, 19]. Some studies highlighted also, among mothers of subjects with ASD, the presence of autoantibodies directed against fetal brain tissue [20, 21]. While these reports seem to support an association between neurodevelopment alterations and immune system dysfunctions in ASD, the relationship between the immune system and the broader autism spectrum, including its subthreshold manifestations, has been poorly investigated [19]. In this case presentation, we aim to report and discuss the presence of autistic traits in a male patient with AN and Behçet's syndrome (BS).

2. Case Presentation

Mr. XY is a 33-year-old man who was referred to the psychiatric service by his rheumatologist for a severe weight loss associated with complete food refusal. He was admitted to the psychiatric inpatient unit, and detailed information about the case history was collected.

Mr. XY was first diagnosed with a BS in 2003, over twelve years before the first psychiatric admission. BS is a disease featuring a chronic vasculitis with a multisystemic involvement of both arteries and veins [22]. Orogenital and ocular lesions are common in BS, but it may show a wide range of clinical manifestations, including involvement of the central nervous system (CNS) [22, 23].

BS is supposed to have a strong genetic vulnerability and, although its pathogenesis is still not clear, it was associated with an alteration of the immune and inflammatory response [22, 23]. Soon after, Mr. XY underwent a genetic examination, showing a HLA-B51 carrier status (an allele associated with the presence of BS) [22]; moreover, a brain magnetic resonance imaging (MRI) highlighted a diffuse signal intensity alteration of the right cerebellar hemisphere. He was first

prescribed glucocorticoids (often employed in the treatment of the acute involvement of the CNS in BS) and colchicine, an alkaloid with anti-inflammatory and immunomodulatory properties used with beneficial effects in different kinds of diseases, from BS from gout [23, 24]. It was also prescribed cyclosporine, an immunosuppressive agent that interferes with the transcription of several cytokines [25]; however, cyclosporine intake was discontinued by Mr. XY in a few months because of gastrointestinal side effects. Since two years from the first diagnosis, he had started to show signs of neuromuscular damage (including tremors, muscle weakness with recurrent falls, dyspnea with reduced exercise tolerance, and evidence of restrictive ventilatory defect with no diffusion impairment). During 2009, he reported recurrent headaches, and as a consequence, a new brain MRI was performed, showing a cavernous angioma located in the right cerebellar hemisphere. Two years later, a further MRI showed a second cavernous angioma in the right ventromedial prefrontal cortex and a venous dysplasia in the right cerebellar hemisphere. A detailed psychiatric history was also obtained. Mr. XY was the first of four children. No family history of mental disorders was reported. He recounted himself as a child and adolescent with pronounced anxiety traits: when he was between four and nine years old, he was quite worried of being separated from his mother, showing a lack of independence in several contexts, such as sleeping, socializing, and attending school. During adolescence and early adulthood, he continued to be overly worried about his parents' sake and to suffer from anxiety symptoms, avoiding social contexts. He never enjoyed practicing sports and he remembers himself as having always been thin and weak. After high school, he started to work as a religion teacher. He had only one stable relationship, lasting about four years, with a woman nine years older than him; he had no children. He described himself as a quiet person, very introspective, and ruminative. Besides his job, he devoted himself to political interests, taking part in the council of the small town where he lived. He never smoked or used illicit drugs. About ten months before the current admission, Mr. XY was shocked by the sudden death of a member of the town council, 30 years older than him, whom the patient remembers as a close friend. Shortly after, he started to be very focused on diet, restricting food intake and selecting only low-calorie food. However, spontaneously, in a few weeks, he regained a normal eating behavior and sufficient social and work adjustment. Despite that, he continued to show intense rumination over his friend's death and started again to fear that some untoward events could happen to his parents. One month before the first psychiatric admission, Mr. XY showed another episode of food restriction, following the break up of his relationship with his girlfriend. He suddenly developed depressive mood, feelings of hopelessness, inner tension, outbursts of anger, and rumination with difficulties to fall asleep. Despite having several talks with his ex-girlfriend, he felt he could not understand the reasons why she broke up with him and started a calorie-restricted diet, feeling that taking control of his body shape could help him to boost his self-esteem and hopefully to bring back his ex-girlfriend. When he saw his rheumatologist, who had treated

him for more than 10 years, he had lost about 8 kg in a month and his BMI was around 18.4, such that he was referred to the psychiatric team. He was admitted voluntarily to the psychiatric inpatient unit. On admission, Mr. XY appeared very skinny, moderately unkempt, and slow at movement. His eyes moved little, staring with a plane face at the clinician's eyes. His speech was slow and circumstantial; answers were given in monosyllables, with a monotone voice. He was strongly worried with his body weight and with food calorie content, completely refusing to eat, such that nasogastric feeding was needed.

Mr. XY was diagnosed as having a partial form of post-traumatic stress disorder (PTSD), a separation anxiety disorder, and AN. However, during his hospital stay, a peculiar social functioning was progressively observed. Mr. XY continued to show a staring gaze, approaching people in an awkward way, without showing social awareness, as if he disregarded other people's feelings. Even when talking about emotionally charged situations, he used a matter-of-fact tone. Despite stating that he was desperate, he appeared calm, being always complacent towards anyone. When asked about his feelings, he found it very hard to express them, rationalizing and intellectualizing his difficulties.

Taking into account these traits, which suggested autistic-like impairments in social communication and in socioemotional reciprocity, a deeper evaluation was conducted. First, Mr. XY's mother was asked to answer some questions, revealing that Mr. XY had been referred to a psychologist during elementary school for anxiety symptoms, but also because he used to be easily distracted by sounds and other stimuli during classes. Moreover, she reported that his classmates often picked on him for his social awkwardness and his overinvolvement in mechanical gadgets.

In addition, Mr. XY agreed to complete the short (14-Item) version of the Ritvo Autism and Asperger Diagnostic Scale-Revised (RAADS-R) [26], a self-report screening questionnaire for ASD, retrieving a score of 11 (cut-off score 14); he fulfilled also the self-report Autism Spectrum Quotient (AQ) [27] and the Adult Autism Subthreshold Spectrum (AdAS Spectrum) [28], both providing a dimensional measure of autistic traits, reporting a score of 22 and 51, respectively. He also gave written informed consent for the publication of his case.

When discharging from the hospital, the initial diagnoses of partial PTSD, separation anxiety disorder, and AN were confirmed. However, the psychiatric team agreed that a sub-threshold autism spectrum was also present and could have shaped the clinical picture of Mr. XY.

3. Discussion

In light of the most recent literature, the autistic-like traits found in Mr. XY allow a more accurate description of the patient's phenotype, and in addition, they may also offer better insight on several aspects of his clinical presentation.

First of all, subjects with ASD are known to show significant difficulties in adjusting to life events, eventually showing an increased vulnerability towards the development of suicidal behaviors in the context of an adjustment disorder

[29]. Moreover, autistic traits have been reported to increase the likelihood of developing posttraumatic symptoms, leading to hypothesize that some autistic dimensions might raise the risk of a broad range of trauma and stress-related disorders [5, 6, 30, 31].

On the other hand, despite no studies have so far investigated the relationship between separation anxiety and autistic traits, subjects with ASD often experience symptoms of separation anxiety [32] that are in turn tightly interwoven with pathological reactions to grief [33]. According to these observations, it might be hypothesized that not only full-blown autism but also high levels of autistic traits would be associated with significant separation sensitivity: this latter may have contributed to trigger the complex reaction to separation and loss exhibited by Mr. XY.

Furthermore, some remarks can be made about the relationship between Mr. XY's autistic traits and AN. The existing literature, mainly based on samples of females with AN, consistently shows that AN is associated with certain pre-morbid traits, such as perfectionism, inhibition, anxiety, and rigidity [34, 35], while the scant literature on AN among males reports similar traits in anorexic males as well [16, 17]. More recently, specific attention has been devoted to the rigid eating habits showed by women with AN, which resemble the narrow interests and repetitive behavior of autistic people, and to the deficits in theory of mind and social intelligence that seem to be common in both AN and ASD [36–38]. As stated above, AN has been hypothesized to be a specific ASD phenotype (which would be more frequently, but possibly not exclusively, associated with female gender), characterized by restricted interest and repetitive behaviors focused on food and diet [12, 13]. On this basis, autistic dimensions have been investigated mostly among AN females, who showed higher levels of autistic traits compared to healthy control subjects [10, 11]. However, while female AN has already been linked to high levels of autistic traits, no study specifically focused on investigating the relationship between AN and ASD, or subthreshold autistic traits, in male samples. The reported case of an adult male with AN and autistic-like features may lead to hypothesize that autism spectrum and AN would be associated also among males and that atypical presentations of autistic traits would not be limited to the female gender. Mr. XY, in fact, reported a RAADS-14 score slightly under the threshold for a possible diagnosis of ASD. Accordingly, he did not endorse all criteria needed for a clinical diagnosis of ASD according to DSM-5 guidelines. However, he reported a score of 22 on the AQ, an instrument developed to quantify autistic traits in both clinical and nonclinical samples. Interestingly, Mr. XY's score of 22 is quite close to the mean score of about 21 found by Baron-Cohen and colleagues among AN female patients [39] and far above mean scores reported among healthy controls [40, 41]. Although a significant disproportion between males and females in AN prevalence is usually reported, gender has been actually reported to have little effect on the clinical features of AN, with the only exception of a more chronic course among females [16]. As previously reported in female samples, the case of Mr. XY shows the presence of autistic traits in a male inpatient with AN, confirming that

similarities in the clinical presentation of AN between males and females might outweigh the differences, being linked to the autism spectrum in both sexes. However, given the unique nature of this case in literature, and the rarity of AN diagnosis among males, findings from this case are of limited generalizability, and further studies are requested to clarify the relationship between autism spectrum and AN in light of gender differences: deepening the knowledge in this field may lead to a better understanding of eventual gender biases in both autism spectrum and eating disorders.

In this case, it is noteworthy that Mr. XY shows not only AN but also a higher vulnerability towards developing psychiatric symptoms after adjustment to separation and grief. Moreover, he also showed social phobic traits, another element that has been associated with ASD and female gender [41, 42]. In particular, social anxiety (which features, as ASD, an impairment in the social brain) [43, 44] has been hypothesized to be higher in subjects with autistic traits or milder forms of ASD, who would be more aware of their social difficulties compared to patients with more severe forms of the disorder [41].

This clinical picture should be globally considered in light of the most recent hypotheses in the field of psychopathology, which suggest the possible presence of a neurodevelopmental alteration at the basis of different kinds of psychiatric conditions [45–47]. According to this conceptualization, while the specific severity and timing of the neurodevelopmental alteration and its interaction with the environment may determine different psychopathological trajectories, a common feature of this “neurodevelopmental continuum” would be a higher vulnerability towards traumatic or stressful events during lifetime [45–47]. This perspective may shed more light on the altered, often autistic-like, premorbid functioning reported in psychiatric patients suffering from different kinds of disorders (AN, mood disorders, and schizophrenia) [13, 48–50] as is the case of Mr. XY. Finally, the occurrence of autistic traits in a subject suffering from BS is of specific interest. In both ASD and BS, involvement of the immune system has been hypothesized. For example, dysregulation of immune function and neuroinflammation, as well as the presence of maternal autoantibodies directed against fetal brain tissue, have been involved in the pathophysiology of ASD [19, 21]. Recently, some authors highlighted the potential role of maternal antibodies against stress-induced phosphoprotein 1 (STIP1) [51], a ligand of prion protein found in the developing brain that has been shown to mediate neurogenesis in cultured hippocampal neurons [52, 53]. Intriguingly, previous data also described the presence of autoantibodies to STIP1 in the blood and CSF of patients with neuro-Behçet’s disease [54]. A precocious involvement of the central nervous system, usually occurring in 4 to 49% of BS sufferers after more than 10 years of disease [55], clearly emerges also from Mr. XY’s clinical history, allowing speculating that both Mr. XY’s autistic subthreshold spectrum and neuro-Behçet symptoms might share a common, autoimmune physiopathology. Globally, this case adds to previous literature that stressed a deep intertwining between the immune system and neurode-

velopmental alterations. The possible link between ASD and autoimmunity has been a recurrent topic since 1970, when authors highlighted the presence of immune aberrations and altered immune responses in children with ASD, including a higher prevalence of allergies [56]. In the last decades, increasing literature highlighted several kinds of immune system alterations, such as altered levels of cytokines, in patients with ASD [57]. The authors suggested the presence of shared genetic underpinnings between ASD and autoimmune diseases, which would interact with early environmental factors, in particular during intrauterine life, in shaping neuroimmune patterns [19, 21]. On the other hand, also AN has been linked with the presence of immune system alterations: while some authors stressed the presence of a higher vulnerability of patients with autoimmune diseases towards the development of AN, pointing out a possible role of the immune system in the pathogenesis of eating disorders [58], others highlighted the presence of immune system dysregulation in patients with AN, hypothesizing that it may be caused by the altered nutritional status [59]. In this framework, recent researches on the role of gut microbiota in both immune and psychiatric disorders have further broadened the perspectives [19]. Microbiota composition has been reported to be altered in both ASD and AN, and several authors hypothesized a role of the so-called microbiota-gut-brain axis in the pathophysiology of these disorders [19, 60]. It is noteworthy that the first investigations on the link between microbiota and ASD were conducted after the frequent observations of gastrointestinal symptoms and/or disordered eating habits in ASD children, further stressing the link between ASD and AN [19]. Microbiota is also known for its crucial role in modulating the immune response, including cytokine levels, exerting proinflammatory or anti-inflammatory activities depending on its compositions [19]. Many authors hypothesized a possible biunivocal influence between microbiota and the CNS through the immune system [19, 61]. The case of Mr. XY seems in line with these hypotheses, showing the presence, at the same time, of autistic symptoms, AN, and BS, a disease linked to altered immune response. However, this case is, at the same time, quite unique in the scientific literature, where the research on ASD and AN mostly focused on females. Moreover, while, as stated above, both autism spectrum and AN were reported to be associated with immune system alterations, scant literature focused on the concomitant occurrence of these three conditions. A recent review further remarked the link between ASD, AN, specific immune alterations, and microbiota profiles, suggesting that future researches in the field of immunopsychiatry should focus on mucosal immunity [61]. Within an integrative framework between central and peripheral systems, more clinical research is needed to clarify the link between neurodevelopment, immune system, and eating disorders.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

- [1] L. Dell'Osso, D. Muti, B. Carpita et al., "The Adult Autism Sub-threshold Spectrum (AdAS) model: a neurodevelopmental approach to mental disorders," *Journal of Psychopathology*, vol. 24, pp. 118–124, 2018.
- [2] K. Takara and T. Kondo, "Comorbid atypical autistic traits as a potential risk factor for suicide attempts among adult depressed patients: a case-control study," *Annals of General Psychiatry*, vol. 13, no. 1, p. 33, 2014.
- [3] L. Dell'Osso, C. Conversano, M. Corsi et al., "Polysubstance and behavioral addictions in a patient with bipolar disorder: role of lifetime subthreshold autism spectrum," *Case Reports in Psychiatry*, vol. 2018, Article ID 1547975, 5 pages, 2018.
- [4] L. Dell'Osso, I. M. Cremona, B. Carpita et al., "Correlates of autistic traits among patients with borderline personality disorder," *Comprehensive Psychiatry*, vol. 83, pp. 7–11, 2018.
- [5] B. Carpita, D. Muti, A. Muscarella et al., "Sex differences in the relationship between PTSD spectrum symptoms and autistic traits in a sample of university students," *Clinical Practice and Epidemiology in Mental Health*, vol. 15, no. 1, pp. 110–119, 2019.
- [6] L. Dell'Osso, B. Carpita, I. M. Cremona et al., "The mediating effect of trauma and stressor related symptoms and ruminations on the relationship between autistic traits and mood spectrum," *Psychiatry Research*, vol. 279, pp. 123–129, 2019.
- [7] B. Carpita, C. Carmassi, S. Calderoni et al., "The broad autism phenotype in real-life: clinical and functional correlates of autism spectrum symptoms and rumination among parents of patients with autism spectrum disorder," *CNS Spectrums*, pp. 1–9, 2019.
- [8] L. Dell'Osso, B. Carpita, C. A. Bertelloni et al., "Subthreshold autism spectrum in bipolar disorder: prevalence and clinical correlates," *Psychiatry Research*, vol. 281, p. 112605, 2019.
- [9] L. Dell'Osso, B. Carpita, D. Muti et al., "Mood symptoms and suicidality across the autism spectrum," *Comprehensive Psychiatry*, vol. 91, pp. 34–38, 2019.
- [10] H. Westwood, I. Eisler, W. Mandy, J. Leppanen, J. Treasure, and K. Tchanturia, "Using the autism-spectrum quotient to measure autistic traits in anorexia nervosa: a systematic review and meta-analysis," *Journal of Autism and Developmental Disorders*, vol. 46, no. 3, pp. 964–977, 2016.
- [11] H. Westwood and K. Tchanturia, "Autism spectrum disorder in anorexia nervosa: an updated literature review," *Current Psychiatry Reports*, vol. 19, no. 7, p. 41, 2017.
- [12] L. Dell'Osso, B. Carpita, C. Gesi et al., "Subthreshold autism spectrum disorder in patients with eating disorders," *Comprehensive Psychiatry*, vol. 81, pp. 66–72, 2018.
- [13] L. Dell'Osso, M. Abelli, B. Carpita et al., "Historical evolution of the concept of anorexia nervosa and relationships with orthorexia nervosa, autism, and obsessive-compulsive spectrum," *Neuropsychiatric Disease and Treatment*, vol. 12, pp. 1651–1660, 2016.
- [14] L. Hull, W. Mandy, M. C. Lai et al., "Development and validation of the Camouflaging Autistic Traits Questionnaire (CAT-Q)," *Journal of Autism and Developmental Disorders*, vol. 49, no. 3, pp. 819–833, 2019.
- [15] M. C. Lai, M. V. Lombardo, B. Auyeung, B. Chakrabarti, and S. Baron-Cohen, "Sex/gender differences and autism: setting the scene for future research," *Journal of the American Academy of Child and Adolescent Psychiatry*, vol. 54, no. 1, pp. 11–24, 2015.
- [16] M. Strober, R. Freeman, C. Lampert, J. Diamond, C. Teplinsky, and M. DeAntonio, "Are there gender differences in core symptoms, temperament, and short-term prospective outcome in anorexia nervosa?," *The International Journal of Eating Disorders*, vol. 39, no. 7, pp. 570–575, 2006.
- [17] A. M. Muise, D. G. Stein, and G. Arbes, "Eating disorders in adolescent boys: a review of the adolescent and young adult literature," *The Journal of Adolescent Health*, vol. 33, no. 6, pp. 427–435, 2003.
- [18] P. Goines and J. van de Water, "The immune system's role in the biology of autism," *Current Opinion in Neurology*, vol. 23, no. 2, pp. 111–117, 2010.
- [19] B. Carpita, D. Marazziti, L. Palego, G. Giannaccini, L. Betti, and L. Dell'Osso, "Microbiota, immune system and autism spectrum disorders. An integrative model towards novel treatment options," *Current Medicinal Chemistry*, vol. 26, 2019.
- [20] D. Braunschweig and J. Van De Water, "Maternal autoantibodies in autism," *Archives of Neurology*, vol. 69, no. 6, pp. 693–699, 2012.
- [21] B. Carpita, D. Muti, and L. Dell'Osso, "Oxidative stress, maternal diabetes, and autism spectrum disorders," *Oxidative Medicine and Cellular Longevity*, vol. 2018, Article ID 3717215, 9 pages, 2018.
- [22] M. de Menthon, M. P. Lavalley, C. Maldini, L. Guillevin, and A. Mahr, "HLA-B51/B5 and the risk of Behçet's disease: a systematic review and meta-analysis of case-control genetic association studies," *Arthritis and Rheumatism*, vol. 61, no. 10, pp. 1287–1296, 2009.
- [23] S. Saip, G. Akman-Demir, and A. Siva, "Neuro-Behçet syndrome," *Neurologic Aspects of Systemic Disease Part III*, vol. 121, pp. 1703–1723, 2014.
- [24] A. Y. Gasparyan, L. Ayzvazyan, M. Yessirkepov, and G. D. Kitas, "Colchicine as an anti-inflammatory and cardioprotective agent," *Expert Opinion on Drug Metabolism & Toxicology*, vol. 11, no. 11, pp. 1781–1794, 2015.
- [25] D. Faulds, K. L. Goa, and P. Benfield, "Cyclosporin," *Drugs*, vol. 45, no. 6, pp. 953–1040, 1993.
- [26] J. M. Erksso, L. M. Andersen, and S. Bejerot, "RAADS-14 screen: validity of a screening tool for autism spectrum disorder in an adult psychiatric population," *Molecular Autism*, vol. 4, no. 1, p. 49, 2013.
- [27] S. Baron-Cohen, S. Wheelwright, R. Skinner, J. Martin, and E. Clubley, "The autism-spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians," *Journal of Autism and Developmental Disorders*, vol. 31, no. 1, pp. 5–17, 2001.
- [28] L. Dell'Osso, C. Gesi, E. Massimetti et al., "Adult Autism Sub-threshold Spectrum (AdAS Spectrum): validation of a questionnaire investigating subthreshold autism spectrum," *Comprehensive Psychiatry*, vol. 73, pp. 61–83, 2017.
- [29] K. Kato, K. Mikami, F. Akama et al., "Clinical features of suicide attempts in adults with autism spectrum disorders," *General Hospital Psychiatry*, vol. 35, no. 1, pp. 50–53, 2013.
- [30] A. L. Roberts, K. C. Koenen, K. Lyall, E. B. Robinson, and M. G. Weisskopf, "Association of autistic traits in adulthood with childhood abuse, interpersonal victimization, and post-traumatic stress," *Child Abuse & Neglect*, vol. 45, pp. 135–142, 2015.
- [31] L. Dell'Osso, D. Muti, P. Lorenzi, A. Della Vecchia, C. Carmassi, and B. Carpita, "Autistic traits and rumination as vulnerability factors towards post-traumatic stress

- symptoms: shaping psychopathological trajectories,” *Journal of Psychopathology*, vol. 26, pp. 12–20, 2020.
- [32] A. Gillott and P. J. Standen, “Levels of anxiety and sources of stress in adults with autism,” *Journal of Intellectual Disabilities*, vol. 11, pp. 359–370, 2016.
- [33] C. Gesi, C. Carmassi, K. M. Shear et al., “Adult separation anxiety disorder in complicated grief: an exploratory study on frequency and correlates,” *Comprehensive Psychiatry*, vol. 72, pp. 6–12, 2017.
- [34] M. Strober, R. Freeman, and W. Morrell, “The long-term course of severe anorexia nervosa in adolescents: survival analysis of recovery, relapse, and outcome predictors over 10–15 years in a prospective study,” *The International Journal of Eating Disorders*, vol. 22, no. 4, pp. 339–360, 1997.
- [35] K. L. Klump, C. M. Bulik, C. Pollice et al., “Temperament and character in women with anorexia nervosa,” *The Journal of Nervous and Mental Disease*, vol. 188, no. 9, pp. 559–567, 2000.
- [36] K. Tchanturia, F. Happé, J. Godley, J. Treasure, N. Bara-Carril, and U. Schmidt, ““Theory of mind” in anorexia nervosa,” *European Eating Disorders Review*, vol. 12, no. 6, pp. 361–366, 2004.
- [37] K. Tchanturia, H. Davies, M. Roberts et al., “Poor cognitive flexibility in eating disorders: examining the evidence using the Wisconsin card sorting task,” *PLoS One*, vol. 7, no. 1, article e28331, 2012.
- [38] D. Hambrook, G. Brown, and K. Tchanturia, “Emotional intelligence in anorexia nervosa: is anxiety a missing piece of the puzzle?,” *Psychiatry Research*, vol. 200, no. 1, pp. 12–19, 2012.
- [39] S. Baron-Cohen, T. Jaffa, S. Davies, B. Auyeung, C. Allison, and S. Wheelwright, “Do girls with anorexia nervosa have elevated autistic traits?,” *Molecular Autism*, vol. 4, no. 1, p. 24, 2013.
- [40] S. Wheelwright, B. Auyeung, C. Allison, and S. Baron-Cohen, “Defining the broader, medium and narrow autism phenotype among parents using the autism spectrum quotient (AQ),” *Molecular Autism*, vol. 1, no. 1, p. 10, 2010.
- [41] L. E. Jobe and S. W. White, “Loneliness, social relationships, and a broader autism phenotype in college students,” *Personality and Individual Differences*, vol. 42, no. 8, pp. 1479–1489, 2007.
- [42] L. Dell’Osso, M. Abelli, S. Pini et al., “The influence of gender on social anxiety spectrum symptoms in a sample of university students,” *Rivista di Psichiatria*, vol. 50, no. 6, pp. 295–301, 2015.
- [43] D. Marazziti, M. Abelli, S. Baroni, B. Carpita, A. Piccinni, and L. Dell’Osso, “Recent findings on the pathophysiology of social anxiety disorder,” *Clinical Neuropsychiatry*, vol. 11, pp. 91–100, 2014.
- [44] D. Marazziti, M. Abelli, S. Baroni, B. Carpita, C. E. Ramacciotti, and L. Dell’Osso, “Neurobiological correlates of social anxiety disorder: an update,” *CNS Spectrums*, vol. 20, no. 2, pp. 100–111, 2015.
- [45] M. J. Owen and M. C. O’Donovan, “Schizophrenia and the neurodevelopmental continuum: evidence from genomics,” *World Psychiatry*, vol. 16, no. 3, pp. 227–235, 2017.
- [46] L. Dell’Osso, P. Lorenzi, and B. Carpita, “Autistic traits and illness trajectories,” *Clinical Practice and Epidemiology in Mental Health*, vol. 15, no. 1, pp. 94–98, 2019.
- [47] L. Dell’Osso, P. Lorenzi, and B. Carpita, “The neurodevelopmental continuum towards a neurodevelopmental gradient hypothesis,” *Journal of Psychopathology*, vol. 25, pp. 179–182, 2019.
- [48] J. Tiihonen, J. Haukka, M. Henriksson et al., “Premorbid intellectual functioning in bipolar disorder and schizophrenia: results from a cohort study of male conscripts,” *The American Journal of Psychiatry*, vol. 162, no. 10, pp. 1904–1910, 2005.
- [49] P. Rocca, S. Galderisi, A. Rossi et al., “Disorganization and real-world functioning in schizophrenia: results from the multicenter study of the Italian Network for Research on Psychoses,” *Schizophrenia Research*, vol. 201, pp. 105–112, 2018.
- [50] P. Bucci, S. Galderisi, A. Mucci et al., “Premorbid academic and social functioning in patients with schizophrenia and its associations with negative symptoms and cognition,” *Acta Psychiatrica Scandinavica*, vol. 138, no. 3, pp. 253–266, 2018.
- [51] D. Braunschweig, P. Krakowiak, P. Duncanson et al., “Autism-specific maternal autoantibodies recognize critical proteins in developing brain,” *Translational Psychiatry*, vol. 3, no. 7, article e277, 2013.
- [52] M. H. Lopes, G. N. Hajj, A. G. Muras et al., “Interaction of cellular prion and stress-inducible protein 1 promotes neurogenesis and neuroprotection by distinct signaling pathways,” *The Journal of Neuroscience*, vol. 25, no. 49, pp. 11330–11339, 2005.
- [53] G. N. M. Hajj, T. G. Santos, Z. S. P. Cook, and V. R. Martins, “Developmental expression of prion protein and its ligands stress-inducible protein 1 and vitronectin,” *The Journal of Comparative Neurology*, vol. 517, no. 3, pp. 371–384, 2009.
- [54] B. Vural, E. Uğurel, E. Tüzün et al., “Anti-neuronal and stress-induced-phosphoprotein 1 antibodies in neuro-Behçet’s disease,” *Journal of Neuroimmunology*, vol. 239, no. 1–2, pp. 91–97, 2011.
- [55] D. Kidd, A. Steuer, A. M. Denman, and P. Rudge, “Neurological complications in Behçet’s syndrome,” *Brain*, vol. 122, no. 11, pp. 2183–2194, 1999.
- [56] E. G. Stubbs and M. L. Crawford, “Depressed lymphocyte responsiveness in autistic children,” *Journal of Autism and Childhood Schizophrenia*, vol. 7, no. 1, pp. 49–55, 1977.
- [57] A. Masi, N. Glozier, R. Dale, and A. J. Guastella, “The immune system, cytokines, and biomarkers in autism spectrum disorder,” *Neuroscience Bulletin*, vol. 33, no. 2, pp. 194–204, 2017.
- [58] S. Zerwas, J. T. Larsen, L. Petersen et al., “Eating disorders, autoimmune, and autoinflammatory disease,” *Pediatrics*, vol. 140, no. 6, article e20162089, 2017.
- [59] A. Marcos, “The immune system in eating disorders: an overview,” *Nutrition*, vol. 13, no. 10, pp. 853–862, 1997.
- [60] R. Roubalová, P. Procházková, H. Papežová, K. Smitka, M. Bilej, and H. Tlaskalová-Hogenová, “Anorexia nervosa: gut microbiota-immune-brain interactions,” *Clinical Nutrition*, vol. 39, no. 3, pp. 676–684, 2020.
- [61] T. Gabriel, S. Paul, A. Berger, and C. Massoubre, “Anorexia nervosa and autism spectrum disorders: future hopes linked to mucosal immunity,” *Neuroimmunomodulation*, vol. 26, no. 6, pp. 265–275, 2020.