Case Report

Bilateral Congenital Agenesis of the Long Head of the Biceps Tendon: The Beginning

Francisco Rego Costa, Cátia Esteves, and Lina Melão

Department of Radiology, São João Hospital, Alameda Professor Hernâni Monteiro, 4200-319 Porto, Portugal

Correspondence should be addressed to Francisco Rego Costa; franciscoregocosta@gmail.com

Received 25 November 2015; Revised 30 December 2015; Accepted 13 January 2016

Academic Editor: Toshihiro Akisue

Copyright © 2016 Francisco Rego Costa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The biceps brachii muscle is prone to variants but absence of the long head of the biceps (LHB) tendon is an exceptionally rare anomaly. This report concerns the fourth case of bilateral congenital absence of the LHB tendon and presents the ultrasonography (US) and magnetic resonance (MR) findings. Our case has the peculiarity of being the first in which bilateral LHB tendon agenesis is not associated with rotator cuff or labral tears.

1. Introduction

Absence of the long head of the biceps (LHB) tendon is a very unusual anomaly. We report the fourth case of bilateral congenital absence of the LHB tendon in the English literature. Ultrasonography (US) and magnetic resonance (MR) findings are exposed.

2. Case Report

A 29-year-old male, Muay Thai amateur fighter, presented with right anterior shoulder pain at rest that exacerbated with overhead activities. The pain was moderate for months but worsened in the last few weeks, specially after sports activity. He had no symptoms in the left shoulder.

Although the patient is a sportsman, there was no history of a significant lesion/traumatic event involving the upper limbs, including shoulder dislocation. He also denied orthopedic surgery to both shoulders.

The physical exam was almost unremarkable. The upper-cut and Speed's tests were negative and the "Popeye" sign was absent. Digital pressure over the bicipital groove did not elicit pain. Although the rotatory stress test and the lateral decubitus load and shift test were negative, the anterior apprehension sign was doubtful. Also there was some limited

range of motion when the patient tried to elevate the left arm with the forearm in supination. Overall, there was no clear evidence of biceps pathology or anterior instability.

On the ultrasonography of the painful and contralateral shoulder we were not able to demonstrate the LHB tendon in its groove (Figure 1), suggesting the presence of a tear or agenesis. There was no evidence of rotator cuff tears, bursitis, or effusion.

Magnetic resonance imaging of both shoulders revealed bilateral absence of the LHB tendon and shallow intertubercular sulci (Figures 2 and 3). There was no evidence of rotator cuff tears on the right side or labral tears bilaterally (Figures 4, 5, and 6). As the patient had no symptoms on the left side, only axial plane MR images were obtained, with the main purpose of proving bilateral agenesis. On these images there was no evidence of rotator cuff abnormalities (Figure 7).

3. Discussion

The biceps brachii muscle is one of the most diverse in the human body; however, congenital absence of the long head of the biceps tendon is a rare anomaly [1–7]. Frequently there are associated findings such as hypoplastic intertubercular groove [1–3] and shoulder instability [4–7]. Unilateral absence is associated with other skeletal and

FIGURE 1: Ultrasound (US) images of both shoulders. (a) US parasagittal image of the left shoulder rotator interval. The long head of the biceps tendon is not identifiable in its normal position (white arrow). (b) US parasagittal image of the right shoulder rotator interval. The long head of the biceps tendon is not discernible in its normal position (white arrow). HH = humeral head; SS = supraspinatus tendon; SC = subscapularis tendon; Del = deltoid muscle.

FIGURE 2: Axial DP FS SE MR sequential images of the right shoulder (TR = 2540; TE = 11.18; NEX = 2; EC = 1) demonstrating a shallow bicipital groove (black arrows) and absence of the long head of the biceps tendon. There is a small amount of fluid in the subacromial/subdeltoid bursa (white arrow), without pathologic significance.

nonskeletal congenital anomalies in 57% of cases [1, 4–6], including at least one case associated with VATER complex (vertebral defects, anal atresia, tracheoesophageal fistula with esophageal atresia, and radial and renal anomalies) [4]. Bilateral absence is extremely uncommon with only three prior reports in the English literature [1–3]. Solely one of these

three cases presented with an additional skeletal abnormality (radial ray) [1].

In our case, the patient was healthy and physically active and no other congenital anomalies were evident. As in other reported cases, bilateral agenesis of the LHB tendon is most commonly diagnosed in a young adult patient complaining

FIGURE 3: Axial DP FS SE MR sequential images of the left shoulder (TR = 2540; TE = 11.18; NEX = 2; EC = 1) showing a shallow bicipital groove (black arrows) and absence of the long head of the biceps tendon.

FIGURE 4: Axial DP FS SE MR images of the right (a) and left (b) shoulder (TR = 475; TE = 13.43; NEX = 1; EC = 1) showing normal labrum morphology bilaterally, without evidence of tears (white arrows). In (a) there is a cyst beneath the lateral humeral head margin, without pathologic significance (asterisk).

of nonspecific shoulder pain [1–3]. Evidence of shallow intertubercular groove is the “rule,” and our case is not an exception. The finding of a shallow groove is very helpful in the differential diagnosis, favoring agenesis over the much more common diagnosis of a biceps tear.

Our case is the first to describe bilateral absence of the LHB tendon not associated with rotator cuff or labral tears. No other significant abnormality was discernible in both shoulders. These findings support that agenesis of the LHB tendon can, by itself, be a source of symptoms and functional

FIGURE 5: Sagittal T2 FS SE MR images of the right shoulder (TR = 2600; TE = 63.9; NEX = 3; EC = 1) (a) and (b) demonstrating the rotator cuff tendons (white arrows) with normal thickness and hypointensity, without tears.

FIGURE 6: Coronal T2 FS SE MR images of the right shoulder (TR = 3500; TE = 63.2; NEX = 2; EC = 1) (a) and (b) showing the normal supraspinatus tendon long axis (white arrows), without tears.

FIGURE 7: Axial DP FS SE MR images of the left shoulder (TR = 475; TE = 13.4; NEX = 2; EC = 1). (a) The supraspinatus tendon is visible, without any discontinuity (white arrow). (b) The subscapularis tendon (black arrow) and the infraspinatus tendon (white arrow) show normal thickness and hypointensity, without tears.

deficit. Based on other cases described, which emphasized the role of the LHB tendon as a shoulder stabilizer [3, 8, 9], we believe that this case can represent the earliest stage of the shoulder disease process generated by the absence of the LHB tendon.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] C. Maldjian, C. Borrero, R. Adam, and D. Vyas, "Nature abhors a vacuum: bilateral prominent rotator cable in bilateral congenital absence of the long head of the biceps tendon," *Skeletal Radiology*, vol. 43, no. 1, pp. 75–78, 2014.
- [2] M. C. Koplas, C. S. Winalski, W. H. Ulmer Jr., and M. Recht, "Bilateral congenital absence of the long head of the biceps tendon," *Skeletal Radiology*, vol. 38, no. 7, pp. 715–719, 2009.
- [3] K. M. Kuhn, J. Carney, D. Solomon, and M. Provencher, "Bilateral absence of the long head of the biceps tendon," *Military Medicine*, vol. 174, no. 5, pp. 548–550, 2009.
- [4] E. L. Smith, E. G. Matzkin, D. H. Kim, J. K. Harpstrite, and D. M. Kan, "Congenital absence of the long head of the biceps brachii tendon as a Vater association," *The American Journal of Orthopedics*, vol. 31, no. 8, pp. 452–454, 2002.
- [5] J. C. Franco, T. P. Knapp, and B. R. Mandelbaum, "Congenital absence of the long head of the biceps tendon. A case report," *The Journal of Bone and Joint Surgery—American Volume*, vol. 87, no. 7, pp. 1584–1586, 2005.
- [6] S. A. Sayeed, J. P. Shah, M. S. Collins, and D. L. Dahm, "Absence of the long head of the biceps tendon associated with glenoid dysplasia and posterior labral tear," *Clinical Anatomy*, vol. 21, no. 7, pp. 728–732, 2008.
- [7] D. A. Glueck, S. D. Mair, and D. L. Johnson, "Shoulder instability with absence of the long head of the biceps tendon," *Arthroscopy*, vol. 19, no. 7, pp. 787–789, 2003.
- [8] C. S. Neer II, "Impingement lesions," *Clinical Orthopaedics and Related Research*, vol. 173, pp. 70–77, 1983.
- [9] E. Etoi, D. K. Kuechle, S. R. Newman, B. F. Morrey, and K. N. An, "Stabilizing function of the biceps in stable and unstable shoulders," *The Journal of Bone & Joint Surgery—British Volume*, vol. 75, no. 4, pp. 546–550, 1993.