Case Report

Capsule Endoscopy: A Cause of Late Small Bowel Obstruction and Perforation

Anders Peter Skovsen, Jakob Burcharth, and Stefan Kobbelgaard Burgdorf

Department of Surgery, Herlev Hospital, University of Copenhagen, Herlev Ringvej 75, 2730 Herlev, Denmark

Correspondence should be addressed to Anders Peter Skovsen; skovsen@dadlnet.dk

Received 20 October 2013; Accepted 13 November 2013

Academic Editors: T. Colak, B. Kirshtein, and G. La Greca

Copyright © 2013 Anders Peter Skovsen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Case Report. A 71-year-old man was admitted to the department of gastroenterology with diffuse abdominal pain. Through the previous 12 months, the patient had experienced episodes of vomiting and watery diarrhea of increasing intensity as well as weight loss. The patient was evaluated with ultrasound, MRI, and subsequently a capsule endoscopy. Six months later, the patient presented, and an abdominal CT-scan showed mechanical small bowel obstruction with suspicion of metallic foreign body and perforation. Laparotomy showed perforation, stenosis, and foreign body, approximately 5 cm from the ileocecal valve. A right hemicolectomy and distal ileectomy (60 cm) with an ileostomy were performed. On further inspection of resection, a capsule endoscope was found impacted in a stenosis. The ileostomy was later reversed without complications. **Conclusion.** It is important to be aware of the possibility of capsule retention, especially in patients with known or suspected Crohn's disease, due to the propensity of Crohn's disease to form stenosis of the bowel. In cases where a stenosis is suspected, it is warranted to perform a patency capsule swallow before subjecting the patient to a capsule endoscopy.

1. Introduction

Capsule endoscopy (CE) has been used since the year 2000, and over 600,000 CE have been performed until now [1, 2]. CE is useful in diagnosing small bowel conditions such as occult bleeding and possible inflammatory bowel disease (IBD) and is generally a secure and safe examination. However, complications can occur. Capsule retention is among the most serious complications and is defined as the capsule not having passed through the bowel within two weeks of ingestion [3]. Retention of the capsule is often seen by structural small bowel abnormalities such as ulcer, tumours, and narrow anastomoses [4]. So far no cases of capsule retention in normal small bowel have been reported. Only three case reports including a total of five patients have reported capsule retention leading to mechanical small bowel obstruction.

The aim of this case report is to present an atypical case of capsule retention leading to mechanical obstruction and small bowel perforation six months after ingestion of the capsule.

2. Case Report

A 71-year-old man was admitted to the department of gastroenterology with diffuse abdominal pain. Through the previous 12 months, the patient had experienced episodes of vomiting and watery diarrhea of increasing intensity as well as weight loss. Both abdominal ultrasound and MRI had shown signs of terminal ileitis. Despite also having undergone a full colonoscopy and gastroscopy without confirmation of inflammatory bowel disease, the patient was suspected of having Crohn's disease. Six months prior to admission, the patient had swallowed a capsule endoscope, and the image feed was deemed normal.

On presentation, an abdominal CT-scan was performed (Figure 1), showing mechanical small bowel obstruction with suspicion of metallic foreign body and perforation. Lab results showed C-reactive protein (CRP) of 204 mg/L.

The patient was taken to theater for a laparotomy. In the terminal ileum, approximately 5 cm from the ileocecal valve, a stenosis and a foreign body were found. Adhesions, chronic inflammation, and fibrinous cover characterized the area.

FIGURE 1: CT-abdomen showing small bowel obstruction, radio-dense foreign body (capsule endoscope), and intraperitoneal air.

FIGURE 2: Photograph of bowel resection, cutup, showing capsule endoscope in situ.

A small perforation with localized fecal peritonitis was discovered near the ileocecal valve. A right hemicolectomy and distal ileectomy (60 cm) with an ileostomy were performed. On further inspection of resection, a capsule endoscope was found impacted in a stenosis (Figure 2). The stenosis was histologically verified as Crohn's disease. The ileostomy was later reversed without complications.

3. Discussion

Small bowel obstruction following retention of a capsule endoscope is very rare. We present a case where impaction of the retained capsule resulted in small bowel obstruction more than six months after ingestion.

Crohn's disease can affect the entire digestive tract. No gold standard exists in the diagnosis of the disease, and definitive diagnosis is often obtained by a combination of clinical, biochemical, radiological, endoscopic, and immunological findings. Capsule endoscopy is superior in assessing small bowel mucosal lesions compared to all other modalities, especially in the proximal and mid-small bowel [5]. Capsule retention is defined as having a capsule remaining in the digestive tract for a minimum of 2 weeks. Only surgical or endoscopic treatment has proven viable in removing a

retained capsule. There are no data for successful medical treatment [6].

In a systematic review involving 22,840 procedures, 184 capsules were reported to be retained in 104 prospective studies and in 46 retrospective studies [3]. The pooled retention rate was 1.4%, which also has been reported by others [7]. Retention rates for confirmed Crohn's disease are reported as 5–13% [6].

It is important to be aware of the possibility of capsule retention, especially in patients with known or suspected Crohn's disease, due to the propensity of Crohn's disease to form stenosis of the bowel. In cases where a stenosis is suspected, it may be warranted to perform a patency capsule swallow before subjecting the patient to a capsule endoscopy.

Disclosure

This paper conforms to ICMJE criteria for authorship and contributorship.

Conflict of Interests

None of the authors of this paper has a financial or personal relationship with other people or organisations that could inappropriately influence or bias the content of this paper.

References

- [1] M. Waterman and R. Eliakim, "Capsule enteroscopy of the small intestine," *Abdominal Imaging*, vol. 34, no. 4, pp. 452–458, 2009.
- [2] G. Iddan, G. Meron, A. Glukhovsky, and P. Swain, "Wireless capsule endoscopy," *Nature*, vol. 405, no. 6785, pp. 417–418, 2000.
- [3] D. Cave, P. Legnani, R. de Franchis, and B. S. Lewis, "ICCE consensus for capsule retention," *Endoscopy*, vol. 37, no. 10, pp. 1065–1067, 2005.
- [4] O. S. Lin, J. J. Brandabur, D. B. Schembre, M.-S. Soon, and R. A. Kozarek, "Acute symptomatic small bowel obstruction due to capsule impaction," *Gastrointestinal Endoscopy*, vol. 65, no. 4, pp. 725–728, 2007.
- [5] S. L. Triester, J. A. Leighton, G. I. Leontiadis et al., "A meta-analysis of the yield of capsule endoscopy compared to other diagnostic modalities in patients with non-stricturing small bowel Crohn's disease," *The American Journal of Gastroenterology*, vol. 101, no. 5, pp. 954–964, 2006.
- [6] Z. Liao, R. Gao, C. Xu, and Z.-S. Li, "Indications and detection, completion, and retention rates of small-bowel capsule endoscopy: a systematic review," *Gastrointestinal Endoscopy*, vol. 71, no. 2, pp. 280–286, 2010.
- [7] F. Li, S. R. Gurudu, G. de Petris et al., "Retention of the capsule endoscope: a single-center experience of 1000 capsule endoscopy procedures," *Gastrointestinal Endoscopy*, vol. 68, no. 1, pp. 174–180, 2008.