Case Report

Infected Abdominal Aortic Aneurysm with *Helicobacter cinaedi*

Kazuhiro Nishida,¹ Takamasa Iwasawa,² Atsushi Tamura,³ and Alan T. Lefor⁴

¹Department of Surgery, Yokosuka General Hospital, Uvama, Kanagawa 238-8567, Japan

²Department of Cardiovascular Internal Medicine, Yokosuka General Hospital, Uvama, Kanagawa 238-8567, Japan

³Department of Cardiovascular Surgery, Yokosuka General Hospital, Uvama, Kanagawa 238-8567, Japan

⁴Department of Surgery, Jichi Medical University, Tochigi, Japan

Correspondence should be addressed to Kazuhiro Nishida; pandadebanda4@gmail.com

Received 13 November 2015; Revised 25 November 2015; Accepted 28 December 2015

Academic Editor: Mario Ganau

Copyright © 2016 Kazuhiro Nishida et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Helicobacter cinaedi is a rare human pathogen which has various clinical manifestations such as cellulitis, bacteremia, arthritis, meningitis, and infectious endocarditis. We report an abdominal aortic aneurysm infected with *Helicobacter cinaedi*, treated successfully with surgical repair and long-term antimicrobial therapy.

1. Introduction

Helicobacter cinaedi (*H. cinaedi*) was first reported in 1985 from the rectal swab of a homosexual man with proctocolitis [1]. This organism is usually isolated from immunocompromised patients, such as HIV infected patients, but recent data shows that immunocompetent hosts can also be infected [2, 3]. A multicenter analysis in Japan found that *H. cinaedi* was isolated from 0.22% of positive blood cultures [4]. This organism has various clinical manifestations, including cellulitis, bacteremia, arthritis, infectious endocarditis, prosthetic graft infection, and infected aneurysms [3–7]. We describe a patient with *H. cinaedi* infected abdominal aortic aneurysm.

2. Case Presentation

A previously healthy 64-year-old man presented with progressive low back pain over two days. His temperature was 36.4°C and there was no abdominal or low back tenderness on physical examination. There was no humoral immunity deficit and HIV antigen/antibody test was negative. A contrast enhanced computed tomography scan showed an infrarenal aortic aneurysm with an eccentric wall that extended to the left common iliac artery, measuring 55 mm in diameter. The following day he had a fever and an infected

abdominal aortic aneurysm was suspected. Three sets of blood cultures were obtained and meropenem and vancomycin begun. Due to sepsis induced respiratory failure and acute kidney injury, he was transferred to the intensive care unit and treated with mechanical ventilation and hemodialysis. One set of blood culture turned positive and *H. cinaedi* was identified by 16S rRNA sequence analysis with 99% similarity and amplification of *gyr B* gene which is specific to *H. cinaedi*. Since the aneurysm had rapidly enlarged to 70 mm, surgical repair with a prosthetic graft was performed on hospital day 27. Pathological analysis of specimen revealed atherosclerosis and infiltration of neutrophils which is consistent with infected aneurysm. He did well postoperatively. After a total of eight weeks of antibiotic treatment, he was discharged on postoperative day 34.

3. Discussion

H. cinaedi is a Gram negative spiral rod which colonizes the gastrointestinal tract of various animals such as hamsters and rhesus monkeys [8]. Contact with animals is thought to be a risk factor for infection, which our patient denied. Diagnosis can be difficult because it rarely grows in traditional culture media. Identification of the organism is based on 16S ribosomal RNA sequence analysis [4, 5]. Optimal duration of therapy is still unknown. Uçkay et al. recommend

prolonged antibiotic therapy because of the potential for recurrence [5]. In the present patient, the isolate was sensitive to penicillins, cephalosporins, and minocycline but resistant to macrolides, fluoroquinolones, and vancomycin. Postoperative minocycline for two weeks and cefazolin for four weeks were administered. Dubois et al. reported 44 cases of infected abdominal aortic aneurysms. *In situ* reconstruction was more often performed than extra-anatomic reconstruction (37 versus six), and in-hospital mortality was low (18.9% versus 50%), but three patients had recurrent infections [9]. Kakuta et al. reported three patients with *H. cinaedi* infected abdominal aortic aneurysms treated with *in situ* reconstruction, and all patients survived [7]. In this patient, *in situ* reconstruction was performed, without evidence of recurrence after two-year follow-up.

4. Conclusion

A patient with a *H. cinaedi* infected abdominal aortic aneurysm was treated successfully with surgical repair and long-term antimicrobial therapy.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] P. A. Totten, C. L. Fennell, F. C. Tenover et al., “*Campylobacter cinaedi* (sp. nov.) and *Campylobacter fennelliae* (sp. nov.): two new *Campylobacter* species associated with enteric disease in homosexual men,” *Journal of Infectious Diseases*, vol. 151, no. 1, pp. 131–139, 1985.
- [2] S. Lasry, J. Simon, A. Marais, J. Pouchot, P. Vinceneux, and Y. Boussougant, “*Helicobacter cinaedi* septic arthritis and bacteremia in an immunocompetent patient,” *Clinical Infectious Diseases*, vol. 31, no. 1, pp. 201–202, 2000.
- [3] Y. Uwamino, K. Muranaka, R. Hase, Y. Otsuka, and N. Hosokawa, “Clinical features of community-acquired *Helicobacter cinaedi* bacteremia,” *Helicobacter*, 2015.
- [4] T. Matsumoto, M. Goto, H. Murakami et al., “Multicenter study to evaluate bloodstream infection by *Helicobacter cinaedi* in Japan,” *Journal of Clinical Microbiology*, vol. 45, no. 9, pp. 2853–2857, 2007.
- [5] I. Uçkay, J. Garbino, P.-Y. Dietrich, B. Ninet, P. Rohner, and V. Jacomo, “Recurrent bacteremia with *Helicobacter cinaedi*: case report and review of the literature,” *BMC Infectious Diseases*, vol. 6, article 86, 2006.
- [6] Y. Suematsu, S. Morizumi, K. Okamura, and M. Kawata, “A rare case of axillobifemoral bypass graft infection caused by *Helicobacter cinaedi*,” *Journal of Vascular Surgery*, vol. 61, no. 1, pp. 231–233, 2014.
- [7] R. Kakuta, H. Yano, H. Kanamori et al., “*Helicobacter cinaedi* infection of abdominal aortic aneurysm, Japan,” *Emerging Infectious Diseases*, vol. 20, no. 11, pp. 1942–1945, 2014.
- [8] K. R. Fernandez, L. M. Hansen, P. Vandamme, B. L. Beaman, and J. V. Solnick, “Captive rhesus monkeys (*Macaca mulatta*) are commonly infected with *Helicobacter cinaedi*,” *Journal of Clinical Microbiology*, vol. 40, no. 6, pp. 1908–1912, 2002.
- [9] M. Dubois, K. Daenens, S. Houthoofd, W. E. Peetermans, and I. Fourneau, “Treatment of mycotic aneurysms with involvement of the abdominal aorta: single-centre experience in 44 consecutive cases,” *European Journal of Vascular and Endovascular Surgery*, vol. 40, no. 4, pp. 450–456, 2010.