NONLINEAR VOLTERRA DIFFERENCE EQUATIONS IN SPACE l^p

MICHAEL I. GIL' AND RIGOBERTO MEDINA

Received 4 December 2003

We consider a class of vector nonlinear discrete-time Volterra equations in space l^p and derive estimates for the norms of solutions. These estimates give us explicit stability conditions, which allow us to avoid finding Lyapunov functionals.

1. Introduction and statement of the main result

Volterra difference equations arise in the mathematical modeling of some real phenomena, and also in numerical schemes for solving differential and integral equations (cf. [7, 8] and the references therein).

One of the basic methods in the theory of stability and boundedness of Volterra difference equations is the direct Lyapunov method (see [1, 3, 4] and the references therein). But finding the Lyapunov functionals for Volterra difference equations is a difficult mathematical problem.

In this paper, we derive estimates for the c_0 - and l^p -norms of solutions for a class of vector Volterra difference equations. These estimates give us explicit stability conditions. To establish the solution estimates, we will interpret the Volterra equations with matrix kernels as operator equations in appropriate spaces. Such an approach for Volterra difference equations has been used by Kolmanovskii and Myshkis [7], Kolmanovskii et al. [8], Kwapisz [9], Medina [10, 11], and Gil' and Medina [6]. Under some restriction, our results generalize the main results from [6, 8, 11].

Let $\mathbb{C}^n$ be an n -dimensional complex Euclidean space with the Euclidean norm $\|\cdot\|_{\mathbb{C}^n}$. For a positive $r \leq \infty$, put

$$\omega_r = \{h \in \mathbb{C}^n : \|\cdot\|_{\mathbb{C}^n} \leq r\}. \quad (1.1)$$

As usual, $c_0 = c_0(\mathbb{C}^n)$ is the Banach space of sequences of vectors from $\mathbb{C}^n$ equipped with the norm

$$\|h\|_{c_0} = \sup_k \|h_k\|_{\mathbb{C}^n} \quad (h = (h_k)_{k=1}^\infty \in c_0, h_k \in \mathbb{C}^n, k = 1, 2, \dots) \quad (1.2)$$

and $l^p = l^p(\mathbb{C}^n)$ ($1 < p < \infty$) is the Banach space of sequences of vectors from $\mathbb{C}^n$ equipped with the norm

$$\|h\|_{l^p} = \left[\sum_{k=1}^{\infty} \|h_k\|_{\mathbb{C}^n}^p \right]^{1/p}, \quad (h = (h_k)_{k=1}^{\infty} \in l^p, h_k \in \mathbb{C}^n, k = 1, 2, \dots). \quad (1.3)$$

Let

$$a_{jk}(h_1, \dots, h_{j-1}) \quad (h_1, \dots, h_{j-1} \in \omega_r; k < j, j = 1, 2, \dots) \quad (1.4)$$

be $n \times n$ matrices dependent on $j - 1$ arguments. Consider the equation

$$x_j = G_j(x_1, \dots, x_{j-1}) + \sum_{k=1}^{j-1} a_{jk}(x_1, \dots, x_{j-1})x_k \quad (j = 1, 2, \dots), \quad (1.5)$$

where the mappings $G_j : \mathbb{C}^{(j-1)n} \rightarrow \mathbb{C}^n$ have the properties

$$f_j := \sup_{h_1, \dots, h_{j-1} \in \omega_r} \|G_j(h_1, \dots, h_{j-1})\|_{\mathbb{C}^n} < \infty \quad (j = 1, 2, \dots). \quad (1.6)$$

Moreover, $G_1 \in \mathbb{C}^n$ is given and

$$f := \{f_1, f_2, \dots\} \in l^p(\mathbb{C}). \quad (1.7)$$

In addition, it is assumed that

$$v_{jk} = \sup_{h_1, \dots, h_{j-1}} \|a_{jk}\|_{\mathbb{C}^n} < \infty \quad (k < j, j = 1, 2, \dots), \quad (1.8)$$

$$N_p(V) := \left[\sum_{j=1}^{\infty} \left(\sum_{k=1}^{j-1} v_{jk}^{p'} \right)^{p/p'} \right]^{1/p} < \infty \quad (1.9)$$

with

$$\frac{1}{p'} + \frac{1}{p} = 1. \quad (1.10)$$

To formulate the result, denote

$$\begin{aligned} m_p(V) &= \sum_{k=0}^{\infty} \frac{N_p^k(V)}{\sqrt[p]{k!}}, \\ Q_p(V) &= \sup_{j=1,2,\dots} \left[\sum_{k=1}^{j-1} v_{jk}^{p'} \right]^{1/p'}. \end{aligned} \quad (1.11)$$

Now we are in a position to formulate the main result of the paper.

THEOREM 1.1. *Let conditions (1.7) and (1.9) hold. Then a solution $x = (x_1, x_2, \dots)$ of (1.5) satisfies the inequalities*

$$\begin{aligned} \|x\|_{l^p} &\leq m_p(V) \|f\|_{l^p}, \\ \|x\|_{c_0} &\leq \|f\|_{c_0} + Q_p(V) m_p(V) \|f\|_{l^p}, \end{aligned} \quad (1.12)$$

provided

$$\|f\|_{c_0} + m_p(V) Q_p(V) \|f\|_{l^p} < r. \quad (1.13)$$

Note that due to the Hölder inequality,

$$m_p(V) = \sum_{k=0}^{\infty} \frac{a^k N_p^k(V)}{a^k \sqrt[p]{k!}} \leq \left[\sum_{k=0}^{\infty} a^{kp'} \right]^{1/p'} \left[\sum_{k=0}^{\infty} \frac{N_p^{pk}(V)}{a^{kp} k!} \right]^{1/p} \quad (1.14)$$

for any positive $a < 1$. So

$$m_p(V) \leq (1 - a^{p'})^{-1/p'} \exp \left[\frac{N_p^p(V)}{pa^p} \right]. \quad (1.15)$$

In particular, taking

$$a = \sqrt[p]{\frac{1}{p}}, \quad (1.16)$$

we have

$$m_p(V) \leq b_p \exp [N_p^p(V)], \quad (1.17)$$

where

$$b_p = \left(1 - \frac{1}{p^{p'/p}} \right)^{-1/p'}. \quad (1.18)$$

2. Proof of Theorem 1.1

First, assume that $r = \infty$. Then conditions (1.7) and (1.9) imply

$$\|x_j\|_{\mathbb{C}^n} \leq f_j + \sum_{k=1}^{j-1} v_{jk} \|x_k\|_{\mathbb{C}^n} \quad (j = 1, 2, \dots). \quad (2.1)$$

Define on $l^p = l^p(\mathbb{R})$ the operator V by

$$[Vh]_j = \sum_{k=1}^{j-1} h_k. \quad (2.2)$$

Here, $[h]_j$ means the j th coordinate of the element $h \in l^p(\mathbb{R})$. The operator V is a quasinilpotent one. So, due to the well-known lemma from the book by Dalec'kiĭ and Kreĭn (see [2, Lemma 3.2.1]) (the comparison principle),

$$\|x_j\|_{\mathbb{C}^n} \leq y_j, \quad (2.3)$$

where y_j is a solution of the equation

$$y_j = f_j + \sum_{k=1}^{j-1} v_{jk} y_k \quad (j = 1, 2, \dots). \quad (2.4)$$

Rewrite this equation as

$$y = f + Vy. \quad (2.5)$$

LEMMA 2.1. *Let conditions (1.7) and (1.9) hold. Then a solution y of (2.5) satisfies the inequality*

$$\|y\|_{l^p} \leq m_p(V) \|f\|_{l^p}. \quad (2.6)$$

Proof. Rewrite (2.5) as

$$y = (I - V)^{-1} f. \quad (2.7)$$

By [5, Lemma 4.3],

$$\|V^k\|_{l^p} \leq \frac{N_p^k(V)}{\sqrt[p]{k!}}. \quad (2.8)$$

Since

$$(I - V)^{-1} = \sum_{k=0}^{\infty} V^k, \quad (2.9)$$

we have

$$\|(I - V)^{-1}\|_{l^p} \leq m_p(V), \quad (2.10)$$

concluding the proof. $\square$

LEMMA 2.2. *Let conditions (1.7) and (1.9) hold. Then a solution y of (2.5) satisfies the inequality*

$$\|y\|_{c_0} \leq \|f\|_{c_0} + m_p(V)\|f\|_{l^p}. \quad (2.11)$$

Proof. From (2.5) it follows that

$$\|y\|_{c_0} \leq \|f\|_{c_0} + \|Vy\|_{c_0}. \quad (2.12)$$

But due to Hölder's inequality

$$\|Vy\|_{c_0} \leq \sup_{j=1,2,\dots} \left[\sum_{k=1}^{j-1} v_{jk}^{p'} \right]^{1/p'} \|y\|_{l^p} = Q_p(V)\|y\|_{l^p}, \quad (2.13)$$

now (2.12) and Lemma 2.1 yield

$$\|y\|_{c_0} \leq \|f\|_{c_0} + Q_p(V)\|y\|_{l^p} \leq \|f\|_{c_0} + Q_p(V)m_p(V)\|f\|_{l^p} \quad (2.14)$$

as claimed. $\square$

Proof of Theorem 1.1. If $r = \infty$, then the required result follows from Lemmas 2.1 and 2.2. Let now $r < \infty$. By a simple application of the Urysohn lemma and Lemma 2.2, we get the required result. $\square$

Acknowledgment

This research was supported by Kamea fund of Israel and by Fondecyt (Chile) under Grant no. 1.030.460.

References

- [1] M. R. Crisci, V. B. Kolmanovskii, E. Russo, and A. Vecchio, *Stability of continuous and discrete Volterra integro-differential equations by Liapunov approach*, J. Integral Equations Appl. **7** (1995), no. 4, 393–411.
- [2] J. L. Dalec'kii and M. G. Krein, *Stability of Solutions of Differential Equations in Banach Space*, American Mathematical Society, Rhode Island, 1974.
- [3] S. N. Elaydi, *An Introduction to Difference Equations*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 1996.
- [4] S. N. Elaydi and S. Murakami, *Asymptotic stability versus exponential stability in linear Volterra difference equations of convolution type*, J. Differ. Equations Appl. **2** (1996), no. 4, 401–410.
- [5] M. I. Gil', *Invertibility and spectrum of Hille-Tamarkin matrices*, Math. Nachr. **244** (2002), 78–88.
- [6] M. I. Gil' and R. Medina, *Boundedness of solutions of matrix nonlinear Volterra difference equations*, Discrete Dyn. Nat. Soc. **7** (2002), no. 1, 19–22.

- [7] V. B. Kolmanovskii and A. D. Myshkis, *Stability in the first approximation of some Volterra difference equations*, J. Differ. Equations Appl. **3** (1998), no. 5-6, 563–569.
- [8] V. B. Kolmanovskii, A. D. Myshkis, and J.-P. Richard, *Estimate of solutions for some Volterra difference equations*, Nonlinear Anal. Ser. A: Theory Methods **40** (2000), no. 1–8, 345–363.
- [9] M. Kwapisz, *On l^p solutions of discrete Volterra equations*, Aequationes Math. **43** (1992), no. 2-3, 191–197.
- [10] R. Medina, *Solvability of discrete Volterra equations in weighted spaces*, Dynam. Systems Appl. **5** (1996), no. 3, 407–421.
- [11] ———, *Stability results for nonlinear difference equations*, Nonlinear Stud. **6** (1999), no. 1, 73–83.

Michael I. Gil': Department of Mathematics, Ben Gurion University of the Negev, P.O. Box 653, Beer-Sheva 84105, Israel

E-mail address: gilmi@cs.bgu.ac.il

Rigoberto Medina: Departamento de Ciencias Exactas, Universidad de Los Lagos, Casilla 933, Osorno, Chile

E-mail address: rmedina@ulagos.cl