Research Article

Dynamics of a Class of Higher Order Difference Equations

Bratislav D. Iričanin

Received 16 October 2007; Revised 11 November 2007; Accepted 29 November 2007

We prove that all positive solutions of the autonomous difference equation $x_n = \alpha x_{n-k} / (1 + x_{n-k} + f(x_{n-1}, \dots, x_{n-m}))$, $n \in \mathbb{N}_0$, where $k, m \in \mathbb{N}$, and f is a continuous function satisfying the condition $\beta \min\{u_1, \dots, u_m\} \leq f(u_1, \dots, u_m) \leq \beta \max\{u_1, \dots, u_m\}$ for some $\beta \in (0, 1)$, converge to the positive equilibrium $\bar{x} = (\alpha - 1)/(\beta + 1)$ if $\alpha > 1$.

Copyright © 2007 Bratislav D. Iričanin. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper, we investigate the global stability of positive solutions of the following autonomous difference equation:

$$x_n = \frac{\alpha x_{n-k}}{1 + x_{n-k} + f(x_{n-1}, \dots, x_{n-m})}, \quad n \in \mathbb{N}_0 := \{0, 1, 2, \dots\}, \quad (1.1)$$

where $k, m \in \mathbb{N}$, and f is a continuous function satisfying the condition

$$\beta \min\{u_1, \dots, u_m\} \leq f(u_1, \dots, u_m) \leq \beta \max\{u_1, \dots, u_m\} \quad (1.2)$$

for some $\beta \in (0, 1)$ (the case $\beta = 0$ is not of some interest since in the case equation it turned as Riccati' one).

Note that in a view of relations (1.2), $\bar{x} = (\alpha - 1)/(\beta + 1)$ is a unique positive equilibrium of (1.1), if $\alpha > 1$.

Further, note that the behaviour of positive solutions of (1.1) for the case $\alpha \in (0, 1)$ is quite simple. Namely, in this case, we have $x_n \leq \alpha x_{n-k}$, so that the sequences $(x_{l[k+r]})_{l \in \mathbb{N}}$, $r \in \{0, 1, \dots, k-1\}$ converge to zero, and consequently, the sequence x_n does. The case

2 Discrete Dynamics in Nature and Society

$\alpha = 1$ is slightly complicated. In this case, the sequences $(x_{kl+r})_{l \in \mathbb{N}}$, $r \in \{0, 1, \dots, k-1\}$ are still convergent, as positive and nonincreasing. If we replace n in (1.1) by kl , $l \in \mathbb{N}$, and then let $l \rightarrow \infty$, we obtain

$$\phi_0 = \frac{\phi_0}{1 + \phi_0 + f(\phi_1, \dots, \phi_m)}, \quad (1.3)$$

where $\phi_i := \lim_{l \rightarrow \infty} x_{kl+i}$, $i \in \{0, 1, \dots, k-1\}$. Without loss of generality, we may assume that $\phi_0 \neq 0$. From (1.3), we have that $\phi_0 + f(\phi_1, \dots, \phi_m) = 0$, which implies $\phi_0 = 0$, a contradiction. Hence, every positive solution of (1.1) converges to zero, also in this case.

Equation (1.1) for the case $\alpha \in (0, 1]$ is a particular case of the difference equation

$$x_n = g(x_{n-1}, \dots, x_{n-s}), \quad (1.4)$$

where the function g satisfies the condition

$$g(u_1, \dots, u_s) \leq \max\{u_1, \dots, u_s\}. \quad (1.5)$$

Equation (1.4), whose function g satisfies condition (1.5) or the following condition:

$$\lim_{x \rightarrow \infty} \frac{g(x, \dots, x)}{x} = 1, \quad (1.6)$$

has been extensively studied by many authors (see, e.g., [9, 14–21, 25]).

In the proof of the result, we use the method of so-called “frame” sequences, that is, a discrete analog of the method of frame curves, commonly used in the theory of differential equations. This method and closely related methods have been used in the literature for many times; see, for example, [21, 1–5, 7, 10, 11, 22–24] and the related references therein. Our motivation stems from [10–12]. Recently, there has been a great interest in studying nonlinear difference equations and systems, in particular those which model some real-life situations in population biology and ecology (see, e.g., [18, 20, 21, 25, 10, 6, 8, 13] and the references cited therein).

2. The global stability of (1.1)

We prove the main result of this paper in this section. Before this, we need a lemma.

LEMMA 2.1. *Assume that $\alpha > 1$, $\beta \in (0, 1)$, $\varepsilon \in (0, (\alpha - 1)(1 - \beta)/(1 + \beta))$, and that $(m_n)_{n \in \mathbb{N}}$ and $(M_n)_{n \in \mathbb{N}}$ are sequences defined as follows:*

$$m_n = \alpha - 1 - \beta M_{n-1} - \frac{\varepsilon}{2^{n-1}}, \quad M_n = \alpha - 1 - \beta m_n + \frac{\varepsilon}{2^{n-1}}, \quad (2.1)$$

for $n \geq 2$, with initial values

$$m_1 = (\alpha - 1) - \beta(\alpha - 1 + \varepsilon) - \varepsilon, \quad M_1 = \alpha - 1 + \varepsilon. \quad (2.2)$$

Then

$$\lim_{n \rightarrow \infty} m_n = \lim_{n \rightarrow \infty} M_n = \frac{\alpha - 1}{1 + \beta}. \quad (2.3)$$

Proof. From (2.1) we obtain the following linear first-order difference equation:

$$M_n = \beta^2 M_{n-1} + (\alpha - 1)(1 - \beta) + (2\beta + 1) \frac{\varepsilon}{2^{n-1}}, \quad n \geq 2, \quad (2.4)$$

whose general solution is

$$M_n = \beta^{2n-2} M_1 + (\alpha - 1)(1 - \beta) \frac{\beta^{2n-2} - 1}{\beta^2 - 1} + \frac{(2\beta + 1)\varepsilon}{2^{n-1}} \sum_{j=0}^{n-2} (2\beta^2)^j. \quad (2.5)$$

Letting $n \rightarrow \infty$ in (2.5), using the assumption $\beta \in (0, 1)$ and Stoltz theorem, it follows that $\lim_{n \rightarrow \infty} M_n = (\alpha - 1)/(1 + \beta)$.

From this and (2.1), it easily follows that $\lim_{n \rightarrow \infty} m_n = (\alpha - 1)/(1 + \beta)$ too, as claimed. $\square$

Now, we are able to formulate and to prove our main result.

THEOREM 2.2. *Assume that $\alpha > 1$, and f is a continuous function satisfying condition (1.2) for some $\beta \in (0, 1)$. Then, every positive solution of (1.1) converges to the positive equilibrium $\bar{x} = (\alpha - 1)/(\beta + 1)$.*

Proof. From (1.1), we have that

$$x_n = \frac{\alpha x_{n-k}}{1 + x_{n-k} + f(x_{n-1}, \dots, x_{n-m})} \leq \frac{\alpha x_{n-k}}{1 + x_{n-k}}, \quad n \in \mathbb{N}. \quad (2.6)$$

Assume that u_n is a solution of the following difference equation:

$$u_n = \frac{\alpha u_{n-k}}{1 + u_{n-k}}, \quad (2.7)$$

with initial values $u_0 = x_0, \dots, u_{-k} = x_{-k}$. It is clear that (2.7) can be reduced into k -independent Riccati equations of the form $z_n = \alpha z_{n-1}/(1 + z_{n-1})$. It is well known that for $\alpha > 1$, there is finite limit $\lim_{n \rightarrow \infty} z_n$ (which is equal to $\alpha - 1$). From this and since in the light of the monotonicity of the function $f(x) = \alpha x/(1 + x)$, we have that $x_n \leq u_n$ for $n \geq -k$. By letting $n \rightarrow \infty$, it follows that

$$S = \limsup_{n \rightarrow \infty} x_n \leq \alpha - 1 = \lim_{n \rightarrow \infty} u_n. \quad (2.8)$$

From (2.8), we have that for every $\varepsilon \in (0, (\alpha - 1)(1 - \beta)/(1 + \beta))$,

$$x_n \leq \alpha - 1 + \varepsilon, \quad (2.9)$$

for $n \geq n_0$. From (1.1), condition (1.2), and relation (2.9), it follows that

$$\frac{\alpha x_{n-k}}{1 + x_{n-k} + \beta(\alpha - 1 + \varepsilon)} \leq \frac{\alpha x_{n-k}}{1 + x_{n-k} + f(x_{n-1}, \dots, x_{n-m})} = x_n \quad (2.10)$$

for every $n \geq n_0 + m$. Assume that (y_n) is a solution of the following difference equation:

$$y_n = \frac{\alpha y_{n-k}}{1 + y_{n-k} + \beta(\alpha - 1 + \varepsilon)} \quad (2.11)$$

4 Discrete Dynamics in Nature and Society

with initial values $y_{n_0} = x_{n_0}, \dots, y_{n_0+k-1} = x_{n_0+k-1}$. Then, since the function $g(x) = \alpha x / (1 + \beta(\alpha - 1 + \varepsilon) + x)$ is increasing on the interval $(0, \infty)$, it is easy to see by the induction that $y_n \leq x_n$ for $n \geq n_0$, and that

$$\lim_{n \rightarrow \infty} y_n = (\alpha - 1) - \beta(\alpha - 1 + \varepsilon). \quad (2.12)$$

Hence, we obtain that

$$0 < (\alpha - 1) - \beta(\alpha - 1 + \varepsilon) \leq \liminf_{n \rightarrow \infty} x_n = I. \quad (2.13)$$

In this way, we formed two frame sequences (y_n) and (u_n) such that $y_n \leq x_n \leq u_n$ for $n \geq n_0 + m$.

Now, let $\varepsilon \in (0, (\alpha - 1)(1 - \beta)/(1 + \beta))$ and sequences $(m_n)_{n \in \mathbb{N}}$ and $(M_n)_{n \in \mathbb{N}}$ be defined by (2.1) with (2.2).

Then we have

$$0 < m_1 \leq I \leq S \leq M_1. \quad (2.14)$$

On the other hand, similar to (2.6)–(2.13), for each $t \in \mathbb{N} \setminus \{1\}$ fixed, we can form the sequences $(y_n^{(t)})$ and $(u_n^{(t)})$ defined by

$$u_n^{(t)} = \frac{\alpha u_{n-k}^{(t)}}{1 + u_{n-k}^{(t)} + \beta m_{t-1}}, \quad y_n^{(t)} = \frac{\alpha y_{n-k}^{(t)}}{1 + y_{n-k}^{(t)} + \beta M_t}, \quad (2.15)$$

and easily show that

$$\begin{aligned} \lim_{n \rightarrow \infty} u_n^{(t)} &= \alpha - 1 - \beta m_{t-1}, & \lim_{n \rightarrow \infty} y_n^{(t)} &= \alpha - 1 - \beta M_t, \\ \alpha - 1 - \beta M_t - \frac{\varepsilon}{2^{t-1}} &< y_n^{(t)} \leq x_n \leq u_n^{(t)} < \alpha - 1 - \beta m_{t-1} + \frac{\varepsilon}{2^{t-1}}, & n \geq n_t. \end{aligned} \quad (2.16)$$

From this and Lemma 2.1, it follows that

$$m_t \leq I \leq S \leq M_t \quad (2.17)$$

for every $t \in \mathbb{N}$. Letting $t \rightarrow \infty$ in relations (2.17), the result follows. $\square$

By Theorem 2.2 and the change of variables $x_n = y_n/c$, we obtain the following corollary.

COROLLARY 2.3. *Assume that $k, m \in \mathbb{N}$, α_j , $j \in \{1, \dots, m\}$, are nonnegative numbers such that $\sum_{j=1}^m \alpha_j = 1$, $\alpha > 1$, $c > 0$, and $\beta \in (0, c)$. Then, every positive solution of the difference equation*

$$x_n = \frac{\alpha x_{n-k}}{1 + c x_{n-k} + \beta \sum_{j=1}^m \alpha_j x_{n-j}}, \quad n \in \mathbb{N}_0, \quad (2.18)$$

converges to the positive equilibrium $\bar{x} = (\alpha - 1)/(\beta + c)$.

In the following example, we show that the function f need not be a linear one.

Example 2.4. Let

$$f(u_1, \dots, u_m) = \sqrt[a]{\frac{\sum_{j=1}^m u_j^a}{m}}, \quad (2.19)$$

where $a > 0$; then this function satisfies conditions of Theorem 2.2. Hence, every positive solution of the difference equation

$$x_n = \frac{\alpha x_{n-k}}{1 + x_{n-k} + \beta \sqrt[a]{(x_{n-1}^a + x_{n-2}^a + \dots + x_{n-m}^a)/m}} \quad (2.20)$$

converges to the positive equilibrium $\bar{x} = (\alpha - 1)/(\beta + 1)$.

Acknowledgments

The author would like to thank the referee for many valuable comments which improved the presentation of the paper. The research was partly supported by the Serbian Ministry of Science, through the Mathematical Institute of SASA, Project no. 144013.

References

- [1] K. S. Berenhaut, J. D. Foley, and S. Stević, "Quantitative bounds for the recursive sequence $y_{n+1} = A + y_n/y_{n-k}$," *Applied Mathematics Letters*, vol. 19, no. 9, pp. 983–989, 2006.
- [2] L. Berg, "On the asymptotics of nonlinear difference equations," *Zeitschrift für Analysis und ihre Anwendungen*, vol. 21, no. 4, pp. 1061–1074, 2002.
- [3] L. Berg, "Inclusion theorems for non-linear difference equations with applications," *Journal of Difference Equations and Applications*, vol. 10, no. 4, pp. 399–408, 2004.
- [4] L. Berg, "Corrections to: "Inclusion theorems for non-linear difference equations with applications,"" *Journal of Difference Equations and Applications*, vol. 11, no. 2, pp. 181–182, 2005.
- [5] L. Berg and L. von Wolfersdorf, "On a class of generalized autoconvolution equations of the third kind," *Zeitschrift für Analysis und ihre Anwendungen*, vol. 24, no. 2, pp. 217–250, 2005.
- [6] R. J. Beverton and S. J. Holt, "On the dynamics of exploited fish populations," *Fisheries Investigations*, vol. 19, pp. 1–53, 1957.
- [7] R. DeVault, G. Ladas, and S. W. Schultz, "On the recursive sequence $x_{n+1} = A/x_n + 1/x_{n-1}$," *Proceedings of the American Mathematical Society*, vol. 126, no. 11, pp. 3257–3261, 1998.
- [8] M. E. Fisher and B. S. Goh, "Stability results for delayed-recruitment models in population dynamics," *Journal of Mathematical Biology*, vol. 19, no. 1, pp. 147–156, 1984.
- [9] G. L. Karakostas and S. Stević, "Slowly varying solutions of the difference equation $x_{n+1} = f(x_n, \dots, x_{n-k+1}) + g(n, x_n, \dots, x_{n-k+1})$," *Journal of Difference Equations and Applications*, vol. 10, no. 3, pp. 249–255, 2004.
- [10] V. L. Kocić and G. Ladas, *Global Asymptotic Behaviour of Nonlinear Difference Equations of Higher Order with Applications*, Kluwer Academic, Dordrecht, The Netherlands, 1993.
- [11] P. Liu and X. Cui, "Hyperbolic logistic difference equation with infinitely many delays," *Mathematics and Computers in Simulation*, vol. 52, no. 3-4, pp. 231–250, 2000.
- [12] R. M. Nigmatulin, "Global stability of a discrete population dynamics model with two delays," *Automation and Remote Control*, vol. 66, no. 12, pp. 1964–1971, 2005.
- [13] E. C. Pielou, *An Introduction to Mathematical Ecology*, Wiley-Interscience, London, UK, 1969.
- [14] S. Stević, "A note on bounded sequences satisfying linear inequalities," *Indian Journal of Mathematics*, vol. 43, no. 2, pp. 223–230, 2001.
- [15] S. Stević, "A generalization of the Copson's theorem concerning sequences which satisfy a linear inequality," *Indian Journal of Mathematics*, vol. 43, no. 3, pp. 277–282, 2001.

- [16] S. Stević, "A global convergence results with applications to periodic solutions," *Indian Journal of Pure and Applied Mathematics*, vol. 33, no. 1, pp. 45–53, 2002.
- [17] S. Stević, "A global convergence result," *Indian Journal of Mathematics*, vol. 44, no. 3, pp. 361–368, 2002.
- [18] S. Stević, "Asymptotic behavior of a sequence defined by iteration with applications," *Colloquium Mathematicum*, vol. 93, no. 2, pp. 267–276, 2002.
- [19] S. Stević, "On the recursive sequence $x_{n+1} = x_n - 1/g(x_n)$," *Taiwanese Journal of Mathematics*, vol. 6, no. 3, pp. 405–414, 2002.
- [20] S. Stević, "Asymptotic behavior of a nonlinear difference equation," *Indian Journal of Pure and Applied Mathematics*, vol. 34, no. 12, pp. 1681–1687, 2003.
- [21] S. Stević, "Asymptotic behavior of a class of nonlinear difference equations," *Discrete Dynamics in Nature and Society*, vol. 2006, Article ID 47156, 10 pages, 2006.
- [22] S. Stević, "Global stability and asymptotics of some classes of rational difference equations," *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 60–68, 2006.
- [23] S. Stević, "On monotone solutions of some classes of difference equations," *Discrete Dynamics in Nature and Society*, vol. 2006, Article ID 53890, 9 pages, 2006.
- [24] S. Stević, "On positive solutions of a $(k+1)$ th order difference equation," *Applied Mathematics Letters*, vol. 19, no. 5, pp. 427–431, 2006.
- [25] S. Stević, "Asymptotics of some classes of higher-order difference equations," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 56813, 20 pages, 2007.

Bratislav D. Iričanin: Faculty of Electrical Engineering, University of Belgrade,
 Bulevar Kralja Aleksandra 73, 11000 Beograd, Serbia
 Email address: iricanin@galeb.etf.bg.ac.yu