Hindawi Publishing Corporation
Discrete Dynamics in Nature and Society
Volume 2008, Article ID 193872, 7 pages
doi:10.1155/2008/193872

*Research Article*

# A New Part-Metric-Related Inequality Chain and an Application

**Xiaofan Yang,[1,2] Fangkuan Sun,[2] and Yuan Yan Tang[2,3]**

[1] *School of Computer and Information, Chongqing Jiaotong University, Chongqing 400074, China*
[2] *College of Computer Science, Chongqing University, Chongqing 400044, China*
[3] *Department of Computer Science, Hong Kong Baptist University, Kowloon, Hong Kong*

Correspondence should be addressed to Xiaofan Yang, xf_yang@163.com

Received 28 September 2007; Accepted 6 November 2007

Recommended by Stevo Stevic

Part-metric-related (PMR) inequality chains are elegant and are useful in the study of difference equations. In this paper, we establish a new PMR inequality chain, which is then applied to show the global asymptotic stability of a class of rational difference equations.



## 1. Introduction

A part-metric related (PMR) inequality chain is a chain of inequalities of the form

$$\min_{1\le i\le n}\left\{a_i, \frac{1}{a_i}\right\} \le f(a_1,\ldots,a_n) \le \max_{1\le i\le n}\left\{a_i, \frac{1}{a_i}\right\}, \tag{1.1}$$

which is closely related to the well-known part metric [1] and has important applications in the study of difference equations [2–13]. Below are three previously known PMR inequality chains:

$$\min_{1\le i\le 4}\left\{a_i, \frac{1}{a_i}\right\} \le \frac{a_1+a_2+a_3a_4}{a_1a_2+a_3+a_4} \le \max_{1\le i\le 4}\left\{a_i, \frac{1}{a_i}\right\} \quad \text{(see [5])}, \tag{1.2}$$

$$\min_{1\le i\le k}\left\{a_i, \frac{1}{a_i}\right\} \le \frac{a_1+\cdots+a_{k-2}+a_{k-1}a_k}{a_1a_2+a_3+\cdots+a_k} \le \max_{1\le i\le k}\left\{a_i, \frac{1}{a_i}\right\} \quad \text{(see [11])}, \tag{1.3}$$

$$\min_{1\le i\le 5}\left\{a_i, \frac{1}{a_i}\right\} \le \frac{(1+w)a_1a_2a_3+a_4+a_5}{a_1a_2+a_1a_3+a_2a_3+wa_4a_5} \le \max_{1\le i\le 5}\left\{a_i, \frac{1}{a_i}\right\}, \quad 1\le w\le 2 \quad \text{(see [13])}. \tag{1.4}$$

In this article, we establish the following PMR inequality chain:

$$\min_{1\le i\le 2p-1}\left\{a_i, \frac{1}{a_i}\right\} \le h_w(a_1,\dots,a_{2p-1}) \le \max_{1\le i\le 2p-1}\left\{a_i, \frac{1}{a_i}\right\}, \tag{1.5}$$

where $h_w$ will be defined in the next section, $p-2 \le w \le p-1$. When $p=3$, chain (1.5) reduces to chain (1.4). On this basis, we prove that the difference equation

$$x_n = h_w(x_{n-2p+1},\dots,x_{n-1}), \quad n=1,2,\dots, \tag{1.6}$$

with positive initial conditions admits a globally asymptotically stable equilibrium $c=1$.

## 2. Main results

This section establishes the main results of this paper. For a function $f(a_1,\dots,a_{2p-1})$, let

$$f(a_1,\dots,a_{2p-1})|_{i_1\sim i_r} = f(a_1,\dots,a_{2p-1})|_{a_{i_j}=m,1\le j\le r}. \tag{2.1}$$

**Lemma 2.1.** *Let $a_1,\dots,a_n,b_1,\dots,b_n > 0$. Then $\min_{1\le i\le n}\{a_i/b_i\} \le (a_1+\cdots+a_n)/(b_1+\cdots+b_n) \le \max_{1\le i\le n}\{a_i/b_i\}$. One equality in the chain holds if and only if $a_1/b_1=\cdots=a_n/b_n$.*

For $p\ge 3$ and $w>0$, define a function $h_w:(\mathfrak{R}_+)^{2p-1}\to\mathfrak{R}_+$ as follows:

$$h_w(a_1,\dots,a_{2p-1}) = \frac{(1+w)\prod_{i=1}^{p}a_i + \prod_{i=p+1}^{2p-1}a_i \times \sum_{i=p+1}^{2p-1}(1/a_i)}{\prod_{i=1}^{p}a_i \times \sum_{i=1}^{p}(1/a_i) + w\prod_{i=p+1}^{2p-1}a_i}. \tag{2.2}$$

Below are two examples of this function:

$$\begin{aligned} h_w(a_1,\dots,a_5) &= \frac{(1+w)a_1a_2a_3 + a_4 + a_5}{a_1a_2 + a_1a_3 + a_2a_3 + wa_4a_5}, \\ h_w(a_1,\dots,a_7) &= \frac{(1+w)a_1a_2a_3a_4 + a_5a_6 + a_5a_7 + a_6a_7}{a_1a_2a_3 + a_1a_2a_4 + a_1a_3a_4 + a_2a_3a_4 + wa_5a_6a_7}. \end{aligned} \tag{2.3}$$

For brevity, let $h_w = h_w(a_1,\dots,a_{2p-1})$. Note that, for each $a_r$, $h_w$ is linear fractional in $a_r$. As a consequence, $h_w$ is monotone in $a_r$. Through simple calculations, we get the following two lemmas.

**Lemma 2.2.** *Let $p \geq 3$, $a_1, \ldots, a_{2p-1} > 0$, $m = \min_{1\leq i\leq 2p-1}\{a_i\}$, $1 \leq r \leq p$.*

(1) *If $h_{p-2}$ is increasing in $a_r$, then $h_{p-2} \leq (p-1)/\sum_{i=1,i\neq r}^{p}(1/a_i)$. The equality holds if and only if $h_{p-2}$ is constant in $a_r$.*

(2) *If $h_{p-2}$ is strictly decreasing in $a_r$, then $h_{p-2} \leq h_{p-2}|_{a_r=m}$. The equality holds if and only if $a_r = m$.*

**Lemma 2.3.** *Let $p \geq 3$, $a_1, \ldots, a_{2p-1} > 0$, $m = \min_{1\leq i\leq 2p-1}\{a_i\}$, $p+1 \leq r \leq 2p-1$.*

(1) *If $h_{p-2}$ is increasing in $a_r$, then $h_{p-2} \leq \sum_{i=p+1,i\neq r}^{2p-1}(1/a_i)/(p-2)$. The equality holds if and only if $h_{p-2}$ is constant in $a_r$.*

(2) *If $h_{p-2}$ is strictly decreasing in $a_r$, then $h_{p-2} \leq h_{p-2}|_{a_r=m}$. The equality holds if and only if $a_r = m$.*

**Theorem 2.4.** *Let $p \geq 3$, $a_1, \ldots, a_{2p-1} > 0$. Then $\min_{1\leq i\leq 2p-1}\{a_i, 1/a_i\} \leq h_{p-2} \leq \max_{1\leq i\leq 2p-1}\{a_i, 1/a_i\}$. One of the two equalities holds if and only if $a_1 = \cdots = a_{2p-1} = 1$.*

*Proof.* Let $m = \min_{1\leq i\leq 2p-1}\{a_i\}$, $M = \max_{1\leq i\leq 2p-1}\{a_i\}$.

We prove only $h_{p-2} \leq \max\{M, 1/m\}$ because $\min\{M, 1/m\} \leq h_{p-2}$ can be proved similarly. We proceed by distinguishing two possible cases.

*Case 1.* There is a permutation $i_1, \ldots, i_{2p-1}$ of $1, 2, \ldots, 2p-1$ such that for each $1 \leq k \leq 2p-1$, either $a_{i_k} = m$ or $h_{p-2}|_{i_1\sim i_{k-1}}$ is strictly decreasing in $a_{i_k}$. Then

$$h_{p-2} \leq h_{p-2}|_{i_1} \leq \cdots \leq h_{p-2}|_{i_1\sim i_{2p-1}} = \frac{1}{2}\left(m + \frac{1}{m}\right) \leq \max\left\{m, \frac{1}{m}\right\} \leq \max\left\{M, \frac{1}{m}\right\}. \tag{2.4}$$

*Case 2.* There is a partial permutation $i_1, \ldots, i_r$ of $1, 2, \ldots, 2p-1$ $(1 \leq r \leq 2p-2)$ such that (a) for each $1 \leq k \leq r$, either $a_{i_k} = m$ or $h_{p-2}|_{i_1\sim i_{k-1}}$ is strictly decreasing in $a_{i_k}$, and (b) for each $t \in \{1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$, $a_{i_t} \neq m$ and $h_{p-2}|_{i_1\sim i_r}$ is increasing in $a_t$. Then

$$m < M, \qquad h_{p-2} \leq h_{p-2}|_{i_1} \leq h_{p-2}|_{i_1\sim i_2} \leq \cdots \leq h_{p-2}|_{i_1\sim i_r}. \tag{2.5}$$

Since $r \leq 2p-2$, there is $t \in \{1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$. If $t \in \{1, \ldots, p\} - \{i_1, \ldots, i_r\}$, it follows from (2.5) and Lemma 2.2 that

$$h_{p-2} \leq h_{p-2}|_{i_1\sim i_r} \leq \frac{(p-1)}{\sum_{i=1,i\neq t}^{p}(1/a_i)|_{i_1\sim i_r}} \leq \max_{1\leq i\leq p, i\neq t}\{a_i\}|_{i_1\sim i_r} \leq M \leq \max\left\{M, \frac{1}{m}\right\}. \tag{2.6}$$

Whereas if $t \in \{p+1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$, it follows from (2.5) and Lemma 2.3 that

$$h_{p-2} \leq h_{p-2}|_{i_1\sim i_r} \leq \frac{\sum_{i=p+1,i\neq t}^{2p-1}(1/a_i)}{(p-2)|_{i_1\sim i_r}} \leq \max_{p+1\leq i\leq 2p-1, i\neq t}\left\{\frac{1}{a_i}\right\}|_{i_1\sim i_r} \leq \frac{1}{m} \leq \max\left\{M, \frac{1}{m}\right\}. \tag{2.7}$$

Hence, $h_{p-2} \leq \max\{M, 1/m\}$ is proven.

Second, we prove that $a_1 = \cdots = a_{2p-1} = 1$ if $h_{p-2} = \max\{M, 1/m\}$. The claim of "$a_1 = \cdots = a_{2p-1} = 1$ if $h_{p-2} = \min\{M, 1/m\}$" can be treated similarly. To this end, we need to prove the following. □

*Claim 1.* If $h_{p-2} = \max\{M, 1/m\}$, then there is a permutation $i_1, \ldots, i_{2p-1}$ of $1, \ldots, 2p-1$ such that for each $1 \leq k \leq 2p-1$, either $a_{i_k} = m$ or $h_{p-2}|_{i_1\sim i_{k-1}}$ is strictly decreasing in $a_{i_k}$.

*Proof of Claim 1.* On the contrary, assume that Claim 1 is not true. Then there is a partial permutation $i_1, \ldots, i_r$ of $1, 2, \ldots, 2p-1$ $(1 \le r \le 2p-2)$ such that (a) for each $1 \le k \le r$, either $a_{i_k} = m$ or $h_{p-2}|_{i_1 \sim i_{k-1}}$ is strictly decreasing in $a_{i_k}$, and (b) for each $t \in \{1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$, $a_{i_t} \neq m$ and $h_{p-2}|_{i_1 \sim i_r}$ is increasing in $a_t$. One of the following two cases must occur.

*Case 1.* There is $t \in \{1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$ such that $h_{p-2}|_{i_1 \sim i_r}$ is strictly increasing in $a_t$. If $t \in \{1, \ldots, p\} - \{i_1, \ldots, i_r\}$, it follows by (2.5), (2.6), and Lemma 2.2 that

$$h_{p-2} \le h_{p-2}|_{i_1 \sim i_r} < \frac{(p-1)}{\sum_{i=1, i \neq t}^{p} (1/a_i)|_{i_1 \sim i_r}} \le \max_{1 \le i \le p, i \neq t} \{a_i\}|_{i_1 \sim i_r} \le \max\left\{M, \frac{1}{m}\right\}. \tag{2.8}$$

A contradiction occurs. Whereas if $t \in \{p+1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$, it follows by (2.5), (2.7), and Lemma 2.3 that

$$h_{p-2} \le h_{p-2}|_{i_1 \sim i_r} < \frac{\sum_{i=p+1, i \neq t}^{2p-1} (1/a_i)}{(p-2)|_{i_1 \sim i_r}} \le \max_{p+1 \le i \le 2p-1, i \neq t} \left\{\frac{1}{a_i}\right\}|_{i_1 \sim i_r} \le \max\left\{M, \frac{1}{m}\right\}. \tag{2.9}$$

Again a contradiction occurs.

*Case 2.* For each $t \in \{1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$, $h_{p-2}|_{i_1 \sim i_r}$ is constant in $a_t$.

First, let us show that $\{1, \ldots, p\} \subseteq \{i_1, \ldots, i_r\}$. Otherwise, there is $t \in \{1, \ldots, p\} - \{i_1, \ldots, i_r\}$. By Lemma 2.2, we have

$$h_{p-2}|_{i_1 \sim i_r} = \frac{(p-1)}{\sum_{i=1, i \neq t}^{p} (1/a_i)|_{i_1 \sim i_r}}. \tag{2.10}$$

If there is $s \in \{1, \ldots, p\} - \{i_1, \ldots, i_r, t\}$, it follows from (2.10) that $h_{p-2}|_{i_1 \sim i_r}$ is strictly increasing in $a_s$, a contradiction occurs. So, $\{1, \ldots, p\} - \{i_1, \ldots, i_r\} = \{t\}$ and thus

$$\max\left\{M, \frac{1}{m}\right\} = h_{p-2} \le h_{p-2}|_{i_1 \sim i_r} = h_{p-2}(a_1, \ldots, a_{2p-1})|_{a_{i_1} = \cdots = a_{i_r} = m} = m < M, \tag{2.11}$$

from which a contradiction follows. So, $\{1, \ldots, p\} \subseteq \{i_1, \ldots, i_r\}$. □

According to the previous argument, there is $t \in \{p+1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\}$. By Lemma 2.3, we get

$$h_{p-2}|_{i_1 \sim i_r} = \frac{\sum_{i=p+1, i \neq t}^{2p-1} (1/a_i)}{(p-2)|_{i_1 \sim i_r}}. \tag{2.12}$$

If there is $s \in \{p+1, \ldots, 2p-1\} - \{i_1, \ldots, i_r, t\}$, it follows from (2.12) that $h_{p-2}|_{i_1 \sim i_r}$ is strictly decreasing in $a_s$, a contradiction. So, $\{p+1, \ldots, 2p-1\} - \{i_1, \ldots, i_r\} = \{t\}$ and thus

$$a_1 = \cdots = a_{t-1} = a_{t+1} = \cdots = a_{2p-1} = m. \tag{2.13}$$

By (2.13) and (2.2), we get

$$h_{p-2} = h_{p-2}|_{i_1 \sim i_r} = \frac{(p-1)m^3 + m + (p-2)a_t}{pm^2 + (p-2)ma_t}. \tag{2.14}$$

Since $h_{p-2}|_{i_1\sim i_r}$ is constant in $a_t$, and $(\mathrm{d}/\mathrm{d}a_t)h_{p-2}|_{i_1\sim i_r} = ((p-1)(p-2)m^2(1-m^2))/[pm^2+(p-2)ma_t]^2$, we derive $m=1$. From (2.12) and (2.13), we get $h_{p-2}|_{i_1\sim i_r} = 1/m$. Since $h_{p-2} = \max\{M, 1/m\}$, all equalities in chains (2.5) and (2.7) hold. These plus $m=1$ yield $h_{p-2}|_{i_1\sim i_r} = 1/m = 1 \geq M$, from which we derive $M = m = 1$. So, $a_t = 1 = m$. This is a contradiction. Claim 1 is proved.

By Claim 1 and $h_{p-2} = \max\{M, 1/m\}$, all equalities in (2.4) must hold. This plus Lemma 2.2 yields $a_1 = \cdots = a_{2p-1} = m$ and $h_{p-2}(m,\ldots,m) = (m+1/m)/2 = m$. This implies $a_1 = \cdots = a_{2p-1} = 1$.

**Theorem 2.5.** *Let $p \geq 3$, $a_1,\ldots,a_{2p-1} > 0$. Then, $\min_{1\leq i\leq 2p-1}\{a_i, 1/a_i\} \leq h_{p-1} \leq \max_{1\leq i\leq 2p-1}\{a_i, 1/a_i\}$. One of the two equalities holds if and only if $a_1 = \cdots = a_p = 1/a_{p+1} = \cdots = 1/a_{2p-1}$.*

*Proof.* By Lemma 2.1 and (2.2), we get

$$\begin{aligned} h_{p-1} &\leq \max\left\{a_1,\ldots,a_p,\frac{1}{a_{p+1}},\ldots,\frac{1}{a_{2p-1}}\right\} \leq \max_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}, \\ h_{p-1} &\geq \min\left\{a_1,\ldots,a_p,\frac{1}{a_{p+1}},\ldots,\frac{1}{a_{2p-1}}\right\} \geq \min_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}. \end{aligned} \tag{2.15}$$

The second claim follows immediately from Lemma 2.1.

We are ready to present the main result of this paper. □

**Theorem 2.6.** *Let $p \geq 3$, $p-2 \leq w \leq p-1$, $a_1,\ldots,a_{2p-1} > 0$. Let*

$$a_k = h_w(a_{k-2p+1},\ldots,a_{k-1}), \quad k = 2p, 2p+1, \ldots. \tag{2.16}$$

*Then $\min_{1\leq i\leq 2p-1}\{a_i, 1/a_i\} \leq a_k \leq \max_{1\leq i\leq 2p-1}\{a_i, 1/a_i\}$, $k = 2p, 2p+1, \ldots$. If $k \geq 2p+1$, then one of the two equalities holds if and only if $a_1 = \cdots = a_{2p-1} = 1$.*

*Proof.* Regard $h_w$ as a linear fractional function in $w$, which is monotone in $w$. By Theorems 2.4 and 2.5, we obtain

$$\begin{aligned} a_{2p} &\geq \min\{h_{p-2}(a_1,\ldots,a_{2p-1}), h_{p-1}(a_1,\ldots,a_{2p-1})\} \geq \min_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}, \\ a_{2p} &\leq \max\{h_{p-2}(a_1,\ldots,a_{2p-1}), h_{p-1}(a_1,\ldots,a_{2p-1})\} \leq \max_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}, \\ a_{2p+1} &\geq \min\{h_{p-2}(a_2,\ldots,a_{2p}), h_{p-1}(a_2,\ldots,a_{2p})\} \geq \min_{2\leq i\leq 2p}\left\{a_i,\frac{1}{a_i}\right\} \geq \min_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}, \\ a_{2p+1} &\leq \max\{h_{p-2}(a_2,\ldots,a_{2p}), h_{p-1}(a_2,\ldots,a_{2p})\} \leq \max_{2\leq i\leq 2p}\left\{a_i,\frac{1}{a_i}\right\} \leq \max_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}. \end{aligned} \tag{2.17}$$

Working inductively, we conclude that for $k = 2p, 2p+1, \ldots$,

$$a_k \geq \min\{h_{p-2}(a_{k-2p+1},\ldots,a_{k-1}), h_{p-1}(a_{k-2p+1},\ldots,a_{k-1})\} \geq \min_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}, \tag{2.18}$$

$$a_k \leq \max\{h_{p-2}(a_{k-2p+1},\ldots,a_{k-1}), h_{p-1}(a_{k-2p+1},\ldots,a_{k-1})\} \leq \max_{1\leq i\leq 2p-1}\left\{a_i,\frac{1}{a_i}\right\}. \tag{2.19}$$

□

*Claim 2.* If $a_{2p+1} = \max_{1\le i\le 2p-1}\{a_i, 1/a_i\}$, then $a_1 = \cdots = a_{2p-1} = 1$.

*Proof of Claim 2.* By (2.19), we get

$$a_{2p+1} = \max\left\{h_{p-2}(a_2,\ldots,a_{2p}), h_{p-1}(a_2,\ldots,a_{2p})\right\} = \max_{1\le i\le 2p-1}\left\{a_i, \frac{1}{a_i}\right\}. \tag{2.20}$$

Here, we encounter two possible cases.

*Case 1.* $a_{2p+1} = h_{p-2}(a_2,\ldots,a_{2p}) = \max_{1\le i\le 2p-1}\{a_i, 1/a_i\}$. By Theorem 2.4, we get $a_2 = \cdots = a_{2p} = 1$ and, hence, $a_{2p+1} = 1$. Then $1 = a_{2p+1} = \max_{1\le i\le 2p-1}\{a_i, 1/a_i\} = \max\{a_1, 1/a_1\}$, implying $a_1 = 1$.

*Case 2.* $a_{2p+1} = h_{p-1}(a_2,\ldots,a_{2p}) = \max_{1\le i\le 2p-1}\{a_i, 1/a_i\}$. By Theorem 2.5, we get

$$a_2 = \cdots = a_{p+1} = \frac{1}{a_{p+2}} = \cdots = \frac{1}{a_{2p}}, \tag{2.21}$$

and consequently,

$$a_{2p+1} = h_{p-1}(a_2,\ldots,a_{2p}) = a_2. \tag{2.22}$$

Then,

$$\begin{aligned}\max_{1\le i\le 2p-1}\left\{a_i, \frac{1}{a_i}\right\} = a_{2p+1} = \frac{1}{a_{2p}} &\le \frac{1}{\min\left\{h_{p-2}(a_1,\ldots,a_{2p-1}), h_{p-1}(a_1,\ldots,a_{2p-1})\right\}} \\ &\le \max_{1\le i\le 2p-1}\left\{a_i, \frac{1}{a_i}\right\}.\end{aligned} \tag{2.23}$$

Hence, all equalities in this chain hold. In particular, we have

$$\min\left\{h_{p-2}(a_1,\ldots,a_{2p-1}), h_{p-1}(a_1,\ldots,a_{2p-1})\right\} = \min_{1\le i\le 2p-1}\left\{a_i, \frac{1}{a_i}\right\}. \tag{2.24}$$

If $h_{p-2}(a_1,\ldots,a_{2p-1}) = \min_{1\le i\le 2p-1}\{a_i, 1/a_i\}$, it follows from Theorem 2.4 that $a_1 = \cdots = a_{2p-1} = 1$. Now, assume that $h_{p-1}(a_1,\ldots,a_{2p-1}) = \min_{1\le i\le 2p-1}\{a_i, 1/a_i\}$. By Theorem 2.5, we get

$$a_1 = \cdots = a_p = \frac{1}{a_{p+1}} = \cdots = \frac{1}{a_{2p-1}}. \tag{2.25}$$

Equations (2.21) and (2.25) imply that $a_1 = \cdots = a_{2p-1} = 1$. Claim 2 is proven. □

By Claim 2 and working inductively, we get that if $a_k = \max_{1\le i\le 2p-1}\{a_i, 1/a_i\}$ for some $k \ge 2p+1$, then $a_1 = \cdots = a_{2p-1} = 1$.

Similarly, we can show that $a_1 = \cdots = a_{2p-1} = 1$ if $a_k = \min_{1\le i\le 2p-1}\{a_i, 1/a_i\}$ holds for some $k \ge 2p+1$.

As an application of Theorem 2.6, we have the following theorem.

**Theorem 2.7.** *Let $p \ge 3$, $p-2 \le w \le p-1$. The difference equation*

$$x_n = h_w(x_{n-2p+1},\ldots,x_{n-1}), \quad n = 1, 2, \ldots, \tag{2.26}$$

*with positive initial conditions admits the globally asymptotically stable equilibrium $c = 1$.*

The proof of this theorem is similar to those in [11, 13], and hence is omitted.

## Acknowledgments

The authors are grateful to the anonymous referees for their valuable comments and suggestions. This work is supported by Natural Science Foundation of China (10771227), Program for New Century Excellent Talent of Educational Ministry of China (NCET-05-0759), Doctorate Foundation of Educational Ministry of China (20050611001), and Natural Science Foundation of Chongqing CSTC (2006BB2231).

## References

[1] A. C. Thompson, "On certain contraction mappings in a partially ordered vector space," *Proceedings of the American Mathematical Society*, vol. 14, no. 3, pp. 438–443, 1963.

[2] A. M. Amleh, N. Kruse, and G. Ladas, "On a class of difference equations with strong negative feedback," *Journal of Difference Equations and Applications*, vol. 5, no. 6, pp. 497–515, 1999.

[3] K. S. Berenhaut, J. D. Foley, and S. Stević, "The global attractivity of the rational difference equation $y_n = y_{n-k} + y_{n-m}/1 + y_{n-k}y_{n-m}$," *Applied Mathematics Letters*, vol. 20, no. 1, pp. 54–58, 2007.

[4] K. S. Berenhaut and S. Stević, "The global attractivity of a higher order rational difference equation," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 940–944, 2007.

[5] N. Kruse and T. Nesemann, "Global asymptotic stability in some discrete dynamical systems," *Journal of Mathematical Analysis and Applications*, vol. 235, no. 1, pp. 151–158, 1999.

[6] T. Nesemann, "Positive nonlinear difference equations: some results and applications," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 47, no. 7, pp. 4707–4717, 2001.

[7] G. Papaschinopoulos and C. J. Schinas, "Global asymptotic stability and oscillation of a family of difference equations," *Journal of Mathematical Analysis and Applications*, vol. 294, no. 2, pp. 614–620, 2004.

[8] S. Stević, "Global stability and asymptotics of some classes of rational difference equations," *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 60–68, 2006.

[9] S. Stević, "Existence of nontrivial solutions of a rational difference equation," *Applied Mathematics Letters*, vol. 20, no. 1, pp. 28–31, 2007.

[10] T. Sun and H. Xi, "Global asymptotic stability of a family of difference equations," *Journal of Mathematical Analysis and Applications*, vol. 309, no. 2, pp. 724–728, 2005.

[11] X. Yang, "Global asymptotic stability in a class of generalized Putnam equations," *Journal of Mathematical Analysis and Applications*, vol. 322, no. 2, pp. 693–698, 2006.

[12] X. Yang, D. J. Evans, and G. M. Megson, "Global asymptotic stability in a class of Putnam-type equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 64, no. 1, pp. 42–50, 2006.

[13] X. Yang, M. Yang, and H. Liu, "A part-metric-related inequality chain and application to the stability analysis of difference equation," *Journal of Inequalities and Applications*, vol. 2007, Article ID 19618, 9 pages, 2007.