Research Article

Permanence and Global Attractivity of Discrete Predator-Prey System with Hassell-Varley Type Functional Response

Runxin Wu and Lin Li

Department of Mathematics and Physics, Fujian University of Technology, Fuzhou, Fujian 350108, China

Correspondence should be addressed to Runxin Wu, runxinwu@163.com

Received 25 February 2009; Accepted 4 April 2009

Recommended by Leonid Berezhansky

By constructing a suitable Lyapunov function and using the comparison theorem of difference equation, sufficient conditions which ensure the permanence and global attractivity of the discrete predator-prey system with Hassell-Varley type functional response are obtained. Example together with its numerical simulation shows that the main results are verifiable.

Copyright © 2009 R. Wu and L. Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The dynamic relationship between predators and their prey has long been and will continue to be one of the dominant themes in both ecology and mathematical ecology due to its universal existence and importance [1]. The most popular predator-prey model is the one with Holling type II functional response [2]:

$$\begin{aligned}\frac{dx}{dt} &= ax\left(1 - \frac{x}{k}\right) - \frac{cxy}{m+x}, \\ \frac{dy}{dt} &= y\left(-d + \frac{fx}{m+x}\right), \\ x(0) &> 0, \quad y(0) > 0,\end{aligned}\tag{1.1}$$

where x , y denote the density of prey and predator species at time t , respectively. The constants a , k , c , m , f , d are all positive constants that stand for prey intrinsic growth rate,

carrying capacity of prey species, capturing rate, half saturation constant, maximal predator growth rate, predator death rate, respectively.

Standard Lotka-Volterra type models, on which a large body of existing predator-prey theory is built, assume that the per capita rate of predation depends on the prey numbers only. There is growing explicit biological and physiological evidence [3–8] that in many situations, especially when predators have to search and share or compete for food, a more suitable general predator-prey model should be based on the “ratio-dependent” theory.

Arditi and Ginzburg [9] proposed the following predator-prey model with ratio-dependent type functional response:

$$\begin{aligned}\frac{dx}{dt} &= ax \left(1 - \frac{x}{k}\right) - \frac{cxy}{my + x}, \\ \frac{dy}{dt} &= y \left(-d + \frac{fx}{my + x}\right), \\ x(0) &> 0, \quad y(0) > 0.\end{aligned}\tag{1.2}$$

It was known that the functional response can depend on predator density in other ways. One of the more widely known ones is due to Hassell and Varley [10]. A general predator-prey model with Hassell-Varley type functional response may take the following form:

$$\begin{aligned}\frac{dx}{dt} &= x \left(a - \frac{x}{k}\right) - \frac{cxy}{my^r + x}, \\ \frac{dy}{dt} &= y \left(-d + \frac{fx}{my^r + x}\right), \quad r \in (0, 1), \\ x(0) &> 0, \quad y(0) > 0.\end{aligned}\tag{1.3}$$

This model is appropriate for interactions, where predators form groups and have applications in biological control. System (1.3) can display richer and more plausible dynamics. In a typical predator-prey interaction where predators do not form groups, one can assume that $r = 1$, producing the so-called ratio-dependent predator-prey dynamics [11]. For terrestrial predators that form a fixed number of tight groups, it is often reasonable to assume that $r = 1/2$. For aquatic predators that form a fixed number of tight groups, $r = 1/3$ may be more appropriate. Recently, Hsu [11] presents a systematic analysis on the above system.

On the other hand, when the size of the population is rarely small or the population has nonoverlapping generation, the discrete time models are more appropriate than the continuous ones [12–24]. This motivated us to propose and study the discrete analogous of predator-prey system (1.3):

$$\begin{aligned}x(k+1) &= x(k) \exp \left\{ a(k) - b(k)x(k) - \frac{c(k)y(k)}{m(k)y^r(k) + x(k)} \right\}, \\ y(k+1) &= y(k) \exp \left\{ -d(k) + \frac{f(k)x(k)}{m(k)y^r(k) + x(k)} \right\},\end{aligned}\tag{1.4}$$

where $r \in (0, 1)$; $\{a(k)\}$, $\{b(k)\}$, $\{c(k)\}$, $\{d(k)\}$, $\{m(k)\}$, $\{f(k)\}$ are all bounded nonnegative sequences. For the rest of the paper, we use the following notations: for any bounded sequence $\{g(k)\}$, set

$$g^u = \sup_{k \in N} g(k), \quad g^l = \inf_{k \in N} g(k). \quad (1.5)$$

By the biological meaning, we will focus our discussion on the positive solution of system of (1.3). Thus, we require that

$$x(0) > 0, \quad y(0) > 0. \quad (1.6)$$

2. Permanence

In order to establish the persistent result for system (1.4), we make some preparations.

Definition 2.1. System (1.4) said to be permanent if there exist positive constants m and M , which are independent of the solution of system (1.4), such that for any positive solution $\{x(k), y(k)\}$ of system (1.4) satisfies

$$m \leq \liminf_{k \rightarrow +\infty} \{x(k), y(k)\} \leq \limsup_{k \rightarrow +\infty} \{x(k), y(k)\} \leq M. \quad (2.1)$$

Lemma 2.2 (see [23]). Assume that $\{x(k)\}$ satisfies $x(k) > 0$ and

$$x(k+1) \leq x(k) \exp\{a(k) - b(k)x(k)\} \quad (2.2)$$

for $k \in N$, where $a(k)$ and $b(k)$ are all nonnegative sequences bounded above and below by positive constants. Then

$$\limsup_{k \rightarrow +\infty} x(k) \leq \frac{1}{b^l} \exp(a^u - 1). \quad (2.3)$$

Lemma 2.3 (see [23]). Assume that $\{x(k)\}$ satisfies

$$x(k+1) \geq x(k) \exp\{a(k) - b(k)x(k)\}, \quad k \geq N_0, \quad (2.4)$$

$\limsup_{k \rightarrow +\infty} x(k) \leq x^*$ and $x(N_0) > 0$, where $a(k)$ and $b(k)$ are all nonnegative sequences bounded above and below by positive constants and $N_0 \in N$. Then

$$\liminf_{k \rightarrow +\infty} x(k) \geq \frac{a^l \exp\{a^l - b^u x^*\}}{b^u}. \quad (2.5)$$

Theorem 2.4. *Assume that*

$$a^l - \frac{c^u M_2^{1-r}}{m^l} > 0, \quad (H_1)$$

$$f^l > d^u \quad (H_2)$$

hold, then system (1.4) is permanent, that is, for any positive solution $\{x(k), y(k)\}$ of system (1.4), one has

$$\begin{aligned} m_1 &\leq \liminf_{k \rightarrow +\infty} x(k) \leq \limsup_{k \rightarrow +\infty} x(k) \leq M_1, \\ m_2 &\leq \liminf_{k \rightarrow +\infty} y(k) \leq \limsup_{k \rightarrow +\infty} y(k) \leq M_2, \end{aligned} \quad (2.6)$$

where

$$\begin{aligned} m_1 &= \frac{a^l - (c^u M_2^{1-r}/m^l)}{b^u} \exp \left\{ a^l - \frac{c^u M_2^{1-r}}{m^l} - b^u M_1 \right\}, \\ m_2 &= \min \left\{ \left\{ \frac{(f^l - d^u)m_1}{m^u d^u} \right\}^{1/r}, \left\{ \frac{(f^l - d^u)m_1}{m^u d^u} \right\}^{1/r} \exp \left\{ -d^u + \frac{f^l m_1}{m^u M_2^r + m_1} \right\} \right\}, \end{aligned} \quad (2.7)$$

$$M_1 = \frac{1}{b^l} \exp(a^u - 1),$$

$$M_2 = \left\{ \frac{f^u M_1}{m^l d^l} \right\}^{1/r} \exp \{ -d^l + f^u \}.$$

Proof. We divided the proof into four claims.

Claim 1. From the first equation of (1.4), we have

$$x(k+1) \leq x(k) \exp \{ a(k) - b(k)x(k) \}. \quad (2.8)$$

By Lemma 2.2, we have

$$\limsup_{k \rightarrow +\infty} x(k) \leq \frac{1}{b^l} \exp(a^u - 1) \stackrel{\text{def}}{=} M_1. \quad (2.9)$$

Above inequality shows that for any $\varepsilon > 0$, there exists a $k_1 > 0$, such that

$$x(k+1) \leq M_1 + \varepsilon, \quad \forall k \geq k_1. \quad (2.10)$$

Claim 2. We divide it into two cases to prove that

$$\limsup_{k \rightarrow +\infty} y(k) \leq M_2. \quad (2.11)$$

Case (i)

There exists an $l_0 \geq k_1$, such that $y(l_0 + 1) \geq y(l_0)$. Then by the second equation of system (1.4), we have

$$-d(l_0) + \frac{f(l_0)x(l_0)}{m(l_0)y^r(l_0) + x(l_0)} \geq 0. \quad (2.12)$$

Hence,

$$-d(l_0) + \frac{f(l_0)x(l_0)}{m(l_0)y^r(l_0)} \geq 0, \quad (2.13)$$

therefore,

$$y^r(l_0) \leq \frac{(l_0)x(l_0)}{m(l_0)d(l_0)} \leq \frac{f^u(M_1 + \varepsilon)}{m^l d^l}, \quad (2.14)$$

and so,

$$y(l_0) \leq \left\{ \frac{f^u(M_1 + \varepsilon)}{m^l d^l} \right\}^{1/r}. \quad (2.15)$$

It follows that

$$\begin{aligned} y(l_0 + 1) &= y(l_0) \exp \left\{ -d(l_0) + \frac{f(l_0)x(l_0)}{m(l_0)y^r(l_0) + x(l_0)} \right\} \\ &\leq \left\{ \frac{f^u(M_1 + \varepsilon)}{m^l d^l} \right\}^{1/r} \exp \left\{ -d^l + f^u \right\} \stackrel{\text{def}}{=} M_{2\varepsilon}. \end{aligned} \quad (2.16)$$

We claim that

$$y(k) \leq M_{2\varepsilon} \quad \forall k \geq l_0. \quad (2.17)$$

By a way of contradiction, assume that there exists a $p_0 > l_0$ such that $y(p_0) > M_{2\varepsilon}$. Then $p_0 \geq l_0 + 2$. Let $y(\tilde{p}_0) \geq l_0 + 2$ be the smallest integer such that $y(\tilde{p}_0) \geq M_{2\varepsilon}$. Then $y(\tilde{p}_0) > y(\tilde{p}_0 - 1)$. The above argument produces that $y(\tilde{p}_0) \leq M_{2\varepsilon}$, a contradiction. This prove the claim.

Case (ii)

We assume that $y(k+1) < y(k)$ for all $K \geq K_1$. Since $y(k)$ is nonincreasing and has a lower bound 0, we know that $\lim_{k \rightarrow +\infty} y(k)$ exists, denoted by $\bar{y}$, then

$$\lim_{k \rightarrow +\infty} y(k) = \bar{y}. \quad (2.18)$$

We claim that

$$\bar{y} \leq \left\{ \frac{f^u(M_1 + \varepsilon)}{m^l d^l} \right\}^{1/r}. \quad (2.19)$$

By a way of contradiction, assume that $\bar{y} > \{f^u(M_1 + \varepsilon)/m^l d^l\}^{1/r}$. Taking limit in the second equation in system (1.4) gives

$$\lim_{k \rightarrow +\infty} \left\{ -d(k) + \frac{f(k)x(k)}{m(k)y^r(k) + x(k)} \right\} = 0, \quad (2.20)$$

which is a contradiction since for $K > K_1$

$$-d(k) + \frac{f(k)x(k)}{m(k)y^r(k) + x(k)} \leq -d^l + \frac{f^u(M_1 + \varepsilon)}{m^l \bar{y}^r} < 0. \quad (2.21)$$

This prove the claim, then we have

$$\limsup_{k \rightarrow +\infty} y(k) = \lim_{k \rightarrow +\infty} y(k) = \bar{y} \leq \left\{ \frac{f^u(M_1 + \varepsilon)}{m^l d^l} \right\}^{1/r}. \quad (2.22)$$

Combining Cases (i) and (ii), we see that

$$\limsup_{k \rightarrow +\infty} y(k) \leq M_{2\varepsilon}. \quad (2.23)$$

Let $\varepsilon \rightarrow 0$, we have

$$\limsup_{k \rightarrow +\infty} y(k) \leq \left\{ \frac{f^u M_1}{m^l d^l} \right\}^{1/r} \exp\{-d^l + f^u\} = M_2. \quad (2.24)$$

Claim 3 ($\liminf_{k \rightarrow \infty} x(k) \geq m_1$). Conditions (H_1) imply that for enough small positive constant ε , we have

$$a^l - \frac{c^u(M_2 + \varepsilon)^{1-r}}{m^l} > 0. \quad (2.25)$$

For above ε , it follows from Claims 1 and 2 that there exists a k_2 such that for all $k > k_2$

$$x(k) \leq M_1 + \varepsilon, \quad y(k) \leq M_2 + \varepsilon. \quad (2.26)$$

From the first equation of (1.4), we have

$$x(k+1) \geq x(k) \exp \left\{ a^l - \frac{c^u(M_2 + \varepsilon)^{1-r}}{m^l} - b^u x(k) \right\}. \quad (2.27)$$

By applying Lemma 2.3 to above inequality, we have

$$\liminf_{k \rightarrow +\infty} x(k) \geq \frac{a^l - (c^u(M_2 + \varepsilon)^{1-r}/m^l)}{b^u} \exp \left\{ a^l - \frac{c^u(M_2 + \varepsilon)^{1-r}}{m^l} - b^u(M_1 + \varepsilon) \right\}. \quad (2.28)$$

Setting $\varepsilon \rightarrow 0$ in (2.28) leads to

$$\liminf_{k \rightarrow +\infty} x(k) \geq \frac{a^l - (c^u M_2^{1-r}/m^l)}{b^u} \exp \left\{ a^l - \frac{c^u M_2^{1-r}}{m^l} - b^u M_1 \right\} \stackrel{\text{def}}{=} m_1. \quad (2.29)$$

This ends the proof of Claim 3.

Claim 4. For any small positive constant $\varepsilon < m_1/2$, from Claims 1–3, it follows that there exists a $k_3 > k_2$ such that for all $k > k_3$

$$x(k) \geq m_1 - \varepsilon, \quad x(k) \leq M_1 + \varepsilon, \quad y(k) \leq M_2 + \varepsilon. \quad (2.30)$$

We present two cases to prove that

$$\liminf_{k \rightarrow +\infty} y(k) \geq m_2 \quad (2.31)$$

Case (i)

There exists an $n_0 \geq k_3$ such that $y(n_0 + 1) \leq y(n_0)$, then

$$-d(n_0) + \frac{f(n_0)x(n_0)}{m(n_0)y^r(n_0) + x(n_0)} \leq 0. \quad (2.32)$$

Hence

$$y(n_0) \geq \left\{ \frac{(f^l - d^u)(m_1 - \varepsilon)}{m^u d^u} \right\}^{1/r} \stackrel{\text{def}}{=} c_{1\varepsilon}, \quad (2.33)$$

and so,

$$y(n_0 + 1) \geq \left\{ \frac{(f^l - d^u)(m_1 - \varepsilon)}{m^u d^u} \right\}^{1/r} \exp \left\{ -d^u + \frac{f^l(m_1 - \varepsilon)}{m^u(M_2 + \varepsilon)^r + (m_1 - \varepsilon)} \right\} \stackrel{\text{def}}{=} c_{2\varepsilon}. \quad (2.34)$$

Set

$$m_{2\varepsilon} = \min\{c_{1\varepsilon}, c_{2\varepsilon}\}. \quad (2.35)$$

We claim that $y(k) \geq m_{2\varepsilon}$ for $k \geq n_0$. By a way of contradiction, assume that there exists a $q_0 \geq n_0$, such that $y(q_0) < m_{2\varepsilon}$. Then $q_0 \geq n_0 + 2$. Let $\tilde{q}_0 \geq n_0 + 2$ be the smallest integer such that $y(\tilde{q}_0) < m_{2\varepsilon}$. Then $y(\tilde{q}_0) < y(\tilde{q}_0 - 1)$, which implies that $y(q_0) \leq m_{2\varepsilon}$, a contradiction, this proves the claim.

Case (ii)

We assume that $y(k + 1) > y(k)$ for all $k > k_3$. According to (2.30), $\lim_{k \rightarrow +\infty} y(k)$ exists, denoted by $\underline{y}$, then

$$\lim_{k \rightarrow +\infty} y(k) = \underline{y}. \quad (2.36)$$

We claim that

$$\underline{y} \geq m_{2\varepsilon}. \quad (2.37)$$

By the way of contradiction, assume that $\underline{y} < m_{2\varepsilon}$. Taking limit in the second equation in system (1.4) gives

$$\lim_{k \rightarrow +\infty} \left\{ -d(k) + \frac{f(k)x(k)}{m(k)y^r(k) + x(k)} \right\} = 0, \quad (2.38)$$

which is a contradiction since for $k > k_3$,

$$-d(k) + \frac{f(k)x(k)}{m(k)y^r(k) + x(k)} \geq -d^u + \frac{f^l(m_1 - \varepsilon)}{m^u y^r + (m_1 - \varepsilon)} > 0. \quad (2.39)$$

The above analysis show that

$$\liminf_{k \rightarrow +\infty} y(k) \geq m_{2\varepsilon}. \quad (2.40)$$

Letting $\varepsilon \rightarrow 0$, we have

$$\liminf_{k \rightarrow +\infty} y(k) \geq m_2, \quad (2.41)$$

where

$$m_2 = \min \left\{ \left\{ \frac{(f^l - d^u)m_1}{m^u d^u} \right\}^{1/r}, \left\{ \frac{(f^l - d^u)m_1}{m^u d^u} \right\}^{1/r} \exp \left\{ -d^u + \frac{f^l m_1}{m^u M_2^r + m_1} \right\} \right\}. \quad (2.42)$$

According to Claims 1–4, we can easily find that the result of Theorem 2.4 holds. $\square$

3. Global Attractivity

Theorem 3.1. Assume that (H_1) and (H_2) hold. Assume further that there exist positive constants α, β , and δ such that

$$\alpha \min \left\{ b^l, \frac{2}{M_1} - b^u \right\} - \alpha \frac{c^u M_2^{1-(r/2)}}{4m^l m_2} - \beta \frac{f^u M_1^{1/2}}{4m_1 m_2^{r/2}} > \delta, \quad (H_3)$$

$$\beta \min \left\{ \frac{f^l m^l m_1 r}{(m^u M_2^r + M_1)^2 M_2^{1-r}}, \frac{2}{M_2} - \frac{f^u M_1^{1/2} r}{4m_2 m_1^{1/2}} \right\} - \alpha \frac{c^u M_1^{1/2}}{4m^l m_2^r m_1^{1/2}} - \alpha \frac{c^u M_2^r (1-r)}{4m_1 m_2^r} > \delta. \quad (H_4)$$

Then system (1.4) with initial condition (1.6) is globally attractive, that is, for any two positive solutions $(x_1(k), y_1(k))$ and $(x_2(k), y_2(k))$ of system (1.4), one has

$$\lim_{k \rightarrow +\infty} |x_1(k) - x_2(k)| = 0, \quad \lim_{k \rightarrow +\infty} |y_1(k) - y_2(k)| = 0. \quad (3.1)$$

Proof. From conditions (H_3) and (H_4) , there exists an enough small positive constant $\varepsilon < \min\{m_1/2, m_2/2\}$ such that

$$\begin{aligned} & \alpha \min \left\{ b^l, \frac{2}{M_1 + \varepsilon} - b^u \right\} - \alpha \frac{c^u (M_2 + \varepsilon)^{1-(r/2)}}{4m^l (m_2 - \varepsilon)} - \beta \frac{f^u (M_1 + \varepsilon)^{1/2}}{4(m_1 - \varepsilon)(m_2 - \varepsilon)^{r/2}} > \delta, \\ & \beta \min \left\{ \frac{f^l m^l (m_1 - \varepsilon) r}{[m^u (M_2 + \varepsilon)^r + (M_1 + \varepsilon)]^2 (M_2 + \varepsilon)^{1-r}}, \frac{2}{M_2 + \varepsilon} - \frac{f^u (M_1 + \varepsilon)^{1/2} r}{4(m_2 - \varepsilon)(m_1 - \varepsilon)^{1/2}} \right\} \\ & \quad - \alpha \frac{c^u (M_1 + \varepsilon)^{1/2}}{4m^l (m_2 - \varepsilon)^r (m_1 - \varepsilon)^{1/2}} - \alpha \frac{c^u (M_2 + \varepsilon)^r (1-r)}{4(m_1 - \varepsilon)(m_2 - \varepsilon)^r} > \delta. \end{aligned} \quad (3.2)$$

Since (H_1) and (H_2) hold, for any positive solutions $(x_1(k), y_1(k))$ and $(x_2(k), y_2(k))$ of system (1.4), it follows from Theorem 2.4 that

$$\begin{aligned} m_1 & \leq \liminf_{k \rightarrow +\infty} x_i(k) \leq \limsup_{k \rightarrow +\infty} x_i(k) \leq M_1, \\ m_2 & \leq \liminf_{k \rightarrow +\infty} y_i(k) \leq \limsup_{k \rightarrow +\infty} y_i(k) \leq M_2, \quad i = 1, 2. \end{aligned} \quad (3.3)$$

For above ε and (3.3), there exists a $k_4 > 0$ such that for all $k > k_4$,

$$m_1 - \varepsilon \leq x_i(k) \leq M_1 + \varepsilon, \quad m_2 - \varepsilon \leq x_i(k) \leq M_2 + \varepsilon, \quad i = 1, 2. \quad (3.4)$$

Let

$$V_1(k) = |\ln x_1(k) - \ln x_2(k)|. \quad (3.5)$$

Then from the first equation of system (1.3), we have

$$\begin{aligned} V_1(k+1) &= |\ln x_1(k+1) - \ln x_2(k+1)| \\ &\leq |\ln x_1(k) - \ln x_2(k) - b(k)(x_1(k) - x_2(k))| \\ &\quad + c(k) \left| \frac{y_1(k)}{m(k)y_1^r(k) + x_1(k)} - \frac{y_2(k)}{m(k)y_2^r(k) + x_2(k)} \right|. \end{aligned} \quad (3.6)$$

Using the Mean Value Theorem, we get

$$\begin{aligned} x_1(k) - x_2(k) &= \exp(\ln x_1(k)) - \exp(\ln x_2(k)) = \xi_1(k)(\ln x_1(k) - \ln x_2(k)), \\ y_1^{1-r}(k) - y_2^{1-r}(k) &= (1-r)\xi_2^{-r}(k)(y_1(k) - y_2(k)), \end{aligned} \quad (3.7)$$

where $\xi_1(k)$ lies between $x_1(k)$ and $x_2(k)$, $\xi_2(k)$ lies between $y_1(k)$ and $y_2(k)$.

It follows from (3.6), (3.7) that

$$\begin{aligned} V_1(k+1) &\leq |\ln x_1(k) - \ln x_2(k)| - \left(\frac{1}{\xi_1(k)} - \left| \frac{1}{\xi_1(k)} - b(k) \right| \right) |x_1(k) - x_2(k)| \\ &\quad + \left| \frac{c(k)y_1(k)}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))} \right| |x_1(k) - x_2(k)| \\ &\quad + \left| \frac{c(k)x_1(k)}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))} \right| |y_1(k) - y_2(k)| \\ &\quad + \left| \frac{c(k)m(k)y_1^r(k)y_2^r(k)}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))} \frac{1-r}{\xi_2^r(k)} \right| |y_1(k) - y_2(k)|. \end{aligned} \quad (3.8)$$

And so, for $k > k_4$

$$\begin{aligned}
 \Delta V_1 \leq & -\min\left\{b^l, \frac{2}{M_1 + \varepsilon} - b^u\right\}|x_1(k) - x_2(k)| \\
 & + \frac{c^u(M_2 + \varepsilon)^{1-(r/2)}}{4m^l(m_2 - \varepsilon)^{r/2}(m_1 - \varepsilon)}|x_1(k) - x_2(k)| \\
 & + \frac{c^u(M_1 + \varepsilon)^{1/2}}{4m^l(m_2 - \varepsilon)^r(m_1 - \varepsilon)^{1/2}}|y_1(k) - y_2(k)| \\
 & + \frac{c^u(M_2 + \varepsilon)^r(1 - r)}{4(m_1 - \varepsilon)(m_2 - \varepsilon)^r}|y_1(k) - y_2(k)|.
 \end{aligned} \tag{3.9}$$

Let

$$V_2(k) = |\ln y_1(k) - \ln y_2(k)|. \tag{3.10}$$

Then from the second equation of system (1.4), we have

$$\begin{aligned}
 V_2(k+1) &= |\ln y_1(k+1) - \ln y_2(k+1)| \\
 &= \left| \ln y_1(k) - \ln y_2(k) + f(k) \left(\frac{x_1(k)}{m(k)y_1^r(k) + x_1(k)} - \frac{x_2(k)}{m(k)y_2^r(k) + x_2(k)} \right) \right| \\
 &\leq \left| \ln y_1(k) - \ln y_2(k) - \frac{f(k)m(k)x_1(k)(y_1^r(k) - y_2^r(k))}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))} \right| \\
 &\quad + \left| \frac{f(k)m(k)y_1^r(k)(x_1(k) - x_2(k))}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))} \right|.
 \end{aligned} \tag{3.11}$$

Using the Mean Value Theorem, we get

$$\begin{aligned}
 y_1(k) - y_2(k) &= \exp(\ln y_1(k)) - \exp(\ln y_2(k)) = \xi_3(k)(\ln y_1(k) - \ln y_2(k)), \\
 y_1^r(k) - y_2^r(k) &= r\xi_4^{r-1}(k)(y_1(k) - y_2(k)),
 \end{aligned} \tag{3.12}$$

where $\xi_3(k)$, $\xi_4(k)$ lie between $y_1(k)$ and $y_2(k)$, respectively. Then, it follows from (3.11), (3.12) that for $k > k_4$,

$$\begin{aligned}
\Delta V_2 &\leq - \left(\frac{1}{\xi_3(k)} - \left| \frac{1}{\xi_3(k)} - \frac{f(k)m(k)x_1(k)r}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))\xi_4^{1-r}(k)} \right| \right) \\
&\quad \times |y_1(k) - y_2(k)| \\
&\quad + \frac{f(k)m(k)y_1^r(k)}{(m(k)y_1^r(k) + x_1(k))(m(k)y_2^r(k) + x_2(k))} |x_1(k) - x_2(k)| \\
&\leq - \min \left\{ \frac{f^l m^l (m_1 - \varepsilon)r}{[m^u(M_2 + \varepsilon)^r + (M_1 + \varepsilon)]^2 (M_2 + \varepsilon)^{1-r}}, \frac{2}{M_2 + \varepsilon} - \frac{f^u(M_1 + \varepsilon)^{1/2}r}{4(m_2 - \varepsilon)(m_1 - \varepsilon)^{1/2}} \right\} \\
&\quad \times |y_1(k) - y_2(k)| \\
&\quad + \frac{f^u(M_1 + \varepsilon)^{r/2}}{4(m_1 - \varepsilon)(m_2 - \varepsilon)^{r/2}} |x_1(k) - x_2(k)|.
\end{aligned} \tag{3.13}$$

Now we define a Lyapunov function as follows:

$$V(k) = \alpha V_1(k) + \beta V_2(k). \tag{3.14}$$

Calculating the difference of V along the solution of system (1.4), for $k > k_4$, it follows from (3.9) and (3.13) that

$$\begin{aligned}
\Delta V &\leq \left[-\alpha \min \left\{ b^l, \frac{2}{M_1 + \varepsilon} - b^u \right\} - \alpha \frac{c^u(M_2 + \varepsilon)^{1-(r/2)}}{4m^l(m_2 - \varepsilon)^{r/2}(m_1 - \varepsilon)} - \beta \frac{f^u(M_1 + \varepsilon)^{r/2}}{4(m_1 - \varepsilon)(m_2 - \varepsilon)^{r/2}} \right] \\
&\quad \times |x_1(k) - x_2(k)| \\
&\quad - \left[\beta \min \left\{ \frac{f^l m^l (m_1 - \varepsilon)r}{[m^u(M_2 + \varepsilon)^r + (M_1 + \varepsilon)]^2 (M_2 + \varepsilon)^{1-r}}, \frac{2}{M_2 + \varepsilon} - \frac{f^u(M_1 + \varepsilon)^{1/2}r}{4(m_2 - \varepsilon)(m_1 - \varepsilon)^{1/2}} \right\} \right. \\
&\quad \left. - \alpha \frac{c^u(M_1 + \varepsilon)^{1/2}}{4m^l(m_2 - \varepsilon)^r(m_1 - \varepsilon)^{1/2}} - \alpha \frac{c^u(M_2 + \varepsilon)^r(1-r)}{4(m_1 - \varepsilon)(m_2 - \varepsilon)^r} \right] \times |y_1(k) - y_2(k)| \\
&\leq -\delta(|x_1(k) - x_2(k)| + |y_1(k) - y_2(k)|).
\end{aligned} \tag{3.15}$$

Summing both sides of the above inequalities from k_4 to k , we have

$$\sum_{p=k_4}^k (V(p+1) - v(p)) \leq -\delta \sum_{p=k_4}^k (|x_1(p) - x_2(p)| + |y_1(p) - y_2(p)|), \tag{3.16}$$

which implies

$$V(k+1) + \delta \sum_{p=k_4}^k (|x_1(p) - x_2(p)| + |y_1(p) - y_2(p)|) \leq V(k_4). \quad (3.17)$$

It follows that

$$\sum_{p=k_4}^k (|x_1(p) - x_2(p)| + |y_1(p) - y_2(p)|) \leq \frac{V(k_4)}{\delta}. \quad (3.18)$$

Using the fundamental theorem of positive series, there exists small enough positive constant $\varepsilon > 0$ such that

$$\sum_{p=k_4}^{+\infty} (|x_1(p) - x_2(p)| + |y_1(p) - y_2(p)|) \leq \frac{V(k_4)}{\delta}, \quad (3.19)$$

which implies that

$$\lim_{k \rightarrow +\infty} (|x_1(k) - x_2(k)| + |y_1(k) - y_2(k)|) = 0, \quad (3.20)$$

that is

$$\lim_{k \rightarrow +\infty} |x_1(k) - x_2(k)| = 0, \quad \lim_{k \rightarrow +\infty} |y_1(k) - y_2(k)| = 0. \quad (3.21)$$

This completes the proof of Theorem 3.1. $\square$

4. Extinction of the Predator Species

This section is devoted to study the extinction of the predator species y .

Theorem 4.1. *Assume that*

$$-d^l + f^u < 0. \quad (H_5)$$

Then, the species y will be driven to extinction, and the species x is permanent, that is, for any positive solution $(x(k), y(k))$ of system (1.4),

$$\begin{aligned} \lim_{k \rightarrow +\infty} y(k) &= 0, \\ m_* &\leq \liminf_{k \rightarrow +\infty} x(k) \leq \limsup_{k \rightarrow +\infty} x(k) \leq M_1, \end{aligned} \quad (4.1)$$

where

$$\begin{aligned} m_* &= \frac{a^l}{b^u} \exp\{a^l - b^u M_1\}, \\ M_1 &= \frac{1}{b^l} \exp(a^u - 1). \end{aligned} \quad (4.2)$$

Proof. For condition (H_5) , there exists small enough positive $\gamma > 0$, such that

$$-d^l + f^u < -\gamma < 0 \quad (4.3)$$

for all $k \in N$, from (4.3) and the second equation of the system (1.4), one can easily obtain that

$$\begin{aligned} y(k+1) &= y(k) \exp\left\{-d(k) + \frac{f(k)x(k)}{m(k)y(k) + x(k)}\right\} \\ &< y(k) \exp\{-d^l + f^u\} \\ &< y(k) \exp\{-\gamma\}. \end{aligned} \quad (4.4)$$

Therefore,

$$y(k+1) < y(0) \exp\{-k\gamma\}, \quad (4.5)$$

which yields

$$\lim_{k \rightarrow +\infty} y(k) = 0. \quad (4.6)$$

From the proof of Theorem 3.1, we have

$$\limsup_{k \rightarrow +\infty} x(k) \leq M_1. \quad (4.7)$$

For enough small positive constant $\varepsilon > 0$,

$$a^l - \frac{c^u \varepsilon^{1-r}}{m^l} > 0. \quad (4.8)$$

For above ε , from (2.9) and (4.6), there exists a $k_5 > 0$ such that for all $k > k_5$,

$$x(k) < M_1 + \varepsilon, \quad y(k) < \varepsilon. \quad (4.9)$$

From the first equation of (1.4), we have

$$x(k+1) \geq x(k) \exp \left\{ a^l - \frac{c^u \varepsilon^{1-r}}{m^l} - b^u x(k) \right\}. \quad (4.10)$$

By Lemma 2.3, we have

$$\liminf_{k \rightarrow +\infty} x(k) \geq \frac{a^l - (c^u \varepsilon^{1-r} / m^l)}{b^u} \exp \left\{ a^l - \frac{c^u \varepsilon^{1-r}}{m^l} - b^u (M_1 + \varepsilon) \right\}. \quad (4.11)$$

Setting $\varepsilon \rightarrow 0$ in (4.11) leads to

$$\liminf_{k \rightarrow +\infty} x(k) \geq \frac{a^l}{b^u} \exp \left\{ a^l - b^u M_1 \right\} \stackrel{\text{def}}{=} m_*. \quad (4.12)$$

The proof of Theorem 4.1 is completed. $\square$

5. Example

The following example shows the feasibility of the main results.

Example 5.1. Consider the following system:

$$\begin{aligned} x(k+1) &= x(k) \exp \left\{ 1.41 + 0.12 \cos(k) - 1.78x(k) - \frac{0.33y(k)}{2.16y^{1/2}(k) + x(k)} \right\}, \\ y(k+1) &= y(k) \exp \left\{ -0.62 + \frac{1.79x(k)}{2.16y^{1/2}(k) + x(k)} \right\}. \end{aligned} \quad (5.1)$$

One could easily see that there exist positive constants $\alpha = 0.01$, $\beta = 0.05$, $\delta = 0.001$ such that

$$\begin{aligned} a^l - \frac{c^u M_2^{1-r}}{m^l} &\approx 2.3281 > 0, \\ f^l &> d^u \approx 1.1700 > 0, \\ \alpha \min \left\{ b^l, \frac{2}{M_1} - b^u \right\} - \alpha \frac{c^u M_2^{1-(r/2)}}{4m^l m_2} - \beta \frac{f^u M_1^{1/2}}{4m_1 m_2^{r/2}} &\approx 0.0011 > \delta, \\ \beta \min \left\{ \frac{f^l m^l m_1 r}{(m^u M_2^r + M_1)^2 M_2^{1-r}}, \frac{2}{M_2} - \frac{f^u M_1^{1/2} r}{4m_2 m_1^{1/2}} \right\} - \alpha \frac{c^u M_1^{1/2}}{4m^l m_2^r m_1^{1/2}} - \alpha \frac{c^u M_2^r (1-r)}{4m_1 m_2^r} &\approx 0.0107 > \delta. \end{aligned} \quad (5.2)$$

Figure 1: Dynamics behavior of system (1.4) with initial conditions $(x(0), y(0)) = (0.9, 0.6), (0.7, 0.4), (0.5, 0.5)$, respectively.

Clearly, conditions (H_1) – (H_4) are satisfied. It follows from Theorems 2.4 and 3.1, that the system is permanent and globally attractive. Numerical simulation from Figure 1 shows that solutions do converge and system is permanent.

6. Conclusion

In this paper, we have obtained sufficient conditions for the permanence and global attractivity of the system (1.4), where $r \in (0, 1)$. If $r = 1$ in the system (1.4), the system (1.4) is a discrete ratio-dependent predator-prey model with Holling-II functional response, in this case, HUO and LI gave sufficient conditions for the permanence of the system in [24], however, they did not provide the condition for the extinction of the predator species y . In this paper, Theorem 2.4 gives the same conditions as that of Huo and Li's condition for the permanence of the system. Furthermore, Theorem 4.1 gives sufficient conditions which ensure the extinction the predator of the system (1.4) when $r = 1$. If $a^l - c^u/m^l > 0$ holds, then the prey species x is permanence. If $r = 0$ in the system of (1.4), the system is a discrete predator-prey model with Holling-II function response, Theorem 4.1 also holds for the case $r = 0$.

References

- [1] A. A. Berryman, "The origins and evolution of predator-prey theory," *Ecology*, vol. 73, no. 5, pp. 1530–1535, 1992.
- [2] H. I. Freedman, *Deterministic Mathematical Models in Population Ecology*, vol. 57 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 1980.
- [3] R. Arditi and H. Saiah, "Empirical evidence of the role of heterogeneity in ratio-dependent consumption," *Ecology*, vol. 73, no. 5, pp. 1544–1551, 1992.
- [4] R. Arditi, L. R. Ginzburg, and H. R. Akcakaya, "Variation in plankton densities among lakes: a case for ratio-dependent predation models," *The American Naturalist*, vol. 138, no. 5, pp. 1287–1296, 1991.
- [5] B. Dai, H. Su, and D. Hu, "Periodic solution of a delayed ratio-dependent predator-prey model with monotonic functional response and impulse," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 70, no. 1, pp. 126–134, 2009.
- [6] F. Chen and J. Shi, "On a delayed nonautonomous ratio-dependent predator-prey model with Holling type functional response and diffusion," *Applied Mathematics and Computation*, vol. 192, no. 2, pp. 358–369, 2007.

- [7] A. P. Gutierrez, "Physiological basis of ratio-dependent predator-prey theory: the metabolic pool model as a paradigm," *Ecology*, vol. 73, no. 5, pp. 1552–1563, 1992.
- [8] R. Xu, F. A. Davidson, and M. A. J. Chaplain, "Persistence and stability for a two-species ratio-dependent predator-prey system with distributed time delay," *Journal of Mathematical Analysis and Applications*, vol. 269, no. 1, pp. 256–277, 2002.
- [9] R. Arditi and L. R. Ginzburg, "Coupling in predator-prey dynamics: ratio-dependence," *Journal of Theoretical Biology*, vol. 139, no. 3, pp. 311–326, 1989.
- [10] M. P. Hassell and G. C. Varley, "New inductive population model for insect parasites and its bearing on biological control," *Nature*, vol. 223, no. 5211, pp. 1133–1137, 1969.
- [11] S.-B. Hsu, T.-W. Hwang, and Y. Kuang, "Global dynamics of a predator-prey model with Hassell-Varley type functional response," *Discrete and Continuous Dynamical Systems*, vol. 10, no. 4, pp. 857–871, 2008.
- [12] J. Yang, "Dynamics behaviors of a discrete ratio-dependent predator-prey system with holling type III functional response and feedback controls," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 186539, 19 pages, 2008.
- [13] F. Chen, "Permanence in a discrete Lotka-Volterra competition model with deviating arguments," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 5, pp. 2150–2155, 2008.
- [14] F. Chen, "Permanence and global attractivity of a discrete multispecies Lotka-Volterra competition predator-prey systems," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 3–12, 2006.
- [15] X. Li and W. Yang, "Permanence of a discrete predator-prey systems with Beddington-deAngelis functional response and feedback controls," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 149267, 8 pages, 2008.
- [16] X. Chen and C. Fengde, "Stable periodic solution of a discrete periodic Lotka-Volterra competition system with a feedback control," *Applied Mathematics and Computation*, vol. 181, no. 2, pp. 1446–1454, 2006.
- [17] Z. Zhou and X. Zou, "Stable periodic solutions in a discrete periodic logistic equation," *Applied Mathematics Letters*, vol. 16, no. 2, pp. 165–171, 2003.
- [18] L. Chen, J. Xu, and Z. Li, "Permanence and global attractivity of a delayed discrete predator-prey system with general Holling-type functional response and feedback controls," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 629620, 17 pages, 2008.
- [19] L. L. Wang and Y. H. Fan, "Permanence for a discrete Nicholson's blowflies model with feedback control and delay," *International Journal of Biomathematics*, vol. 1, no. 4, pp. 433–442, 2008.
- [20] F. Chen, "Permanence for the discrete mutualism model with time delays," *Mathematical and Computer Modelling*, vol. 47, no. 3-4, pp. 431–435, 2008.
- [21] F. Chen, L. Wu, and Z. Li, "Permanence and global attractivity of the discrete Gilpin-Ayala type population model," *Computers & Mathematics with Applications*, vol. 53, no. 8, pp. 1214–1227, 2007.
- [22] F. Chen, "Permanence of a discrete N -species cooperation system with time delays and feedback controls," *Applied Mathematics and Computation*, vol. 186, no. 1, pp. 23–29, 2007.
- [23] F. Chen, "Permanence and global stability of nonautonomous Lotka-Volterra system with predator-prey and deviating arguments," *Applied Mathematics and Computation*, vol. 173, no. 2, pp. 1082–1100, 2006.
- [24] H.-F. Huo and W.-T. Li, "Permanence and global stability of positive solutions of a nonautonomous discrete ratio-dependent predator-prey model," *Discrete Dynamics in Nature and Society*, vol. 2005, no. 2, pp. 135–144, 2005.