Research Article

On the Solutions of the System of Difference Equations $x_{n+1} = \max\{A/x_n, y_n/x_n\}$, $y_{n+1} = \max\{A/y_n, x_n/y_n\}$

Dağıstan Simsek,¹ Bilal Demir,² and Cengiz Cinar²

¹ Department of Industrial Engineering, Engineering-Architecture Faculty, Selcuk University, Kambiis, 42075 Konya, Turkey

² Mathematics Department, Education Faculty, Selcuk University, Meram Yeni Yol, 42090 Konya, Turkey

Correspondence should be addressed to Dağıstan Simsek, dsimsek@selcuk.edu.tr

Received 17 January 2009; Accepted 13 April 2009

Recommended by Guang Zhang

We study the behavior of the solutions of the following system of difference equations $x_{n+1} = \max\{A/x_n, y_n/x_n\}$, $y_{n+1} = \max\{A/y_n, x_n/y_n\}$ where the constant A and the initial conditions are positive real numbers.

Copyright © 2009 Dağıstan Simsek et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Recently, there has been a great interest in studying the periodic nature of nonlinear difference equations. Although difference equations are relatively simple in form, it is, unfortunately, extremely difficult to understand thoroughly the periodic behavior of their solutions. The periodic nature of nonlinear difference equations of the max type has been investigated by many authors. See, for example [1–24].

In this paper we study the behavior of the solutions of the following system of difference equations:

$$x_{n+1} = \max\left\{\frac{A}{x_n}, \frac{y_n}{x_n}\right\}, \quad y_{n+1} = \max\left\{\frac{A}{y_n}, \frac{x_n}{y_n}\right\}, \quad (1.1)$$

where the constant A and the initial conditions are positive real numbers.

2. Main Result

Definition 2.1. Fibonacci sequence is $f_1 = 1, f_2 = 1$ and for $n \geq 3, f_n = f_{n-1} + f_{n-2}$.

Definition 2.2. The symbol $[[\]]$ symbolizes the greatest integer function.

Definition 2.3. The sequence of $a(n) \equiv n \pmod{2}$.

Definition 2.4. The sequence of

$$k(n) = \begin{cases} n, & n = 0, 2, 4, \dots, \\ n + 1, & n = 1, 3, 5, \dots \end{cases} \quad (2.1)$$

Theorem 2.5. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $A < x_0 < y_0$ and $y_0/x_0 > A$.

If $n \geq 1$, then

$$x_n = \left(\frac{A^{f_{k(n)-1}-a(n)} x_0^{f_{k(n)}}}{y_0^{f_{k(n)}}} \right)^{(-1)^n}, \quad (2.2)$$

$y_1 = x_0/y_0$, and if $n \geq 2$

$$y_n = \left(\frac{y_0^{f_{k(n-1)}+1}}{A^{f_{k(n-1)}+a(n)-1} x_0^{f_{k(n-1)}+1}} \right)^{(-1)^n}. \quad (2.3)$$

Proof. Let $y_0 > A$, then

$$x_1 = \max \left\{ \frac{A}{x_0}, \frac{y_0}{x_0} \right\} = \frac{y_0}{x_0} > A,$$

$$y_1 = \max \left\{ \frac{A}{y_0}, \frac{x_0}{y_0} \right\} = \frac{x_0}{y_0} < A,$$

$$x_2 = \max \left\{ \frac{A}{x_1}, \frac{y_1}{x_1} \right\} = \frac{Ax_0}{y_0} < A,$$

$$y_2 = \max \left\{ \frac{A}{y_1}, \frac{x_1}{y_1} \right\} = \frac{x_1}{y_1} = \frac{y_0^2}{x_0^2} > A,$$

$$x_3 = \max \left\{ \frac{A}{x_2}, \frac{y_2}{x_2} \right\} = \frac{y_2}{x_2} = \frac{y_0^3}{Ax_0^3} > A,$$

$$y_3 = \max \left\{ \frac{A}{y_2}, \frac{x_2}{y_2} \right\} = \frac{A}{y_2} = \frac{Ax_0^2}{y_0^2} < A,$$

$$\begin{aligned}
x_4 &= \max \left\{ \frac{A}{x_3}, \frac{y_3}{x_3} \right\} = \frac{A^2 x_0^3}{y_0^3} < A, \\
y_4 &= \max \left\{ \frac{A}{y_3}, \frac{x_3}{y_3} \right\} = \frac{x_3}{y_3} = \frac{y_0^5}{A^2 x_0^5} > A, \\
x_5 &= \max \left\{ \frac{A}{x_4}, \frac{y_4}{x_4} \right\} = \frac{y_4}{x_4} = \frac{y_0^8}{A^3 x_0^8} > A, \\
y_5 &= \max \left\{ \frac{A}{y_4}, \frac{x_4}{y_4} \right\} = \frac{A}{y_4} = \frac{A^3 x_0^5}{y_0^5} < A, \\
&\vdots
\end{aligned} \tag{2.4}$$

$n \geq 1$ then

$$x_n = \left(\frac{A^{f_{k(n)}-1-a(n)} x_0^{f_{k(n)}}}{y_0^{f_{k(n)}}} \right)^{(-1)^n}, \tag{2.5}$$

$y_1 = x_0/y_0$ then $n \geq 2$,

$$y_n = \left(\frac{y_0^{f_{k(n-1)}+1}}{A^{f_{k(n-1)}+a(n)-1} x_0^{f_{k(n-1)}+1}} \right)^{(-1)^n}. \tag{2.6}$$

□

Theorem 2.6. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $A < y_0 < x_0$. $x_1 = x_0/y_0$ and if $n \geq 2$,

$$x_n = \left(\frac{A^{f_{k(n-1)}+a(n)-1} x_0^{f_{k(n-1)}+1}}{y_0^{f_{k(n-1)}+1}} \right)^{(-1)^n}. \tag{2.7}$$

If $n \geq 1$, then

$$y_n = \left(\frac{y_0^{f_{k(n)}}}{A^{f_{k(n)}-1-a(n)} x_0^{f_{k(n)}}} \right)^{(-1)^n}. \tag{2.8}$$

Proof. Similarly we can obtain the proof as the proof of Theorem 2.5. □

Theorem 2.7. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $A < x_0 < y_0$ and $(y_0/x_0) > A$.

$$\begin{aligned}
 (a) \quad & \lim_{n \rightarrow \infty} x_{2n} = 0, \\
 & \lim_{n \rightarrow \infty} y_{2n} = \infty. \\
 (b) \quad & \lim_{n \rightarrow \infty} x_{2n+1} = \infty, \\
 & \lim_{n \rightarrow \infty} y_{2n+1} = 0.
 \end{aligned} \tag{2.9}$$

Proof. (a) We obtain that

$$\begin{aligned}
 \lim_{n \rightarrow \infty} x_{2n} &= \lim_{n \rightarrow \infty} \left(\frac{A^{f_{k(2n)}-1-a(2n)} x_0^{f_{k(2n)}}}{y_0^{f_{k(2n)}}} \right)^{(-1)^{2n}} \\
 &= \lim_{n \rightarrow \infty} \left(\frac{A^{f_{k(2n)}-1-a(2n)} x_0^{f_{k(2n)}}}{y_0^{f_{k(2n)}}} \right)^{(-1)^{2k}} \\
 &= \lim_{n \rightarrow \infty} \left(\frac{A^{f_{2n}-1-a(2n)} x_0^{f_{2n}}}{y_0^{f_{2n}}} \right) \\
 &= 0, \\
 \lim_{n \rightarrow \infty} y_{2n} &= \lim_{n \rightarrow \infty} \left(\frac{y_0^{f_{k(2n-1)}+1}}{A^{f_{k(2n-1)}+a(2n)-1} x_0^{f_{k(2n-1)}+1}} \right)^{(-1)^{2n}} \\
 &= \lim_{n \rightarrow \infty} \left(\frac{y_0^{f_{k(2n-1)}+1}}{A^{f_{k(2n-1)}+a(2n)-1} x_0^{f_{k(2n-1)}+1}} \right)^{(-1)^{2k}} \\
 &= \lim_{k \rightarrow \infty} \left(\frac{y_0^{f_{2n}+1}}{A^{f_{2n}-1} x_0^{f_{2n}+1}} \right) \\
 &= \infty.
 \end{aligned} \tag{2.10}$$

(b) Similarly we can obtain the proof of (b) as the proof of (a). □

Theorem 2.8. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $A < y_0 < x_0$ and $(x_0/y_0) > A$.

$$\begin{aligned}
 (a) \quad & \lim_{n \rightarrow \infty} x_{2n} = \infty, \\
 & \lim_{n \rightarrow \infty} y_{2n} = 0. \\
 (b) \quad & \lim_{n \rightarrow \infty} x_{2n+1} = 0, \\
 & \lim_{n \rightarrow \infty} y_{2n+1} = \infty.
 \end{aligned} \tag{2.11}$$

Proof. Similarly we can obtain the proof as the proof of Theorem 2.7. $\square$

Theorem 2.9. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $1 < x_0 < y_0 < A$.

If $n \geq 1$, then

$$x_n = \left(\frac{x_0}{A^{a(n)}} \right)^{(-1)^n}. \tag{2.12}$$

If $n \geq 1$, then

$$y_n = \left(\frac{y_0}{A^{a(n)}} \right)^{(-1)^n}. \tag{2.13}$$

Proof. Let

$$\begin{aligned}
 x_1 &= \max \left\{ \frac{A}{x_0}, \frac{y_0}{x_0} \right\} = \frac{A}{x_0} < A, \\
 y_1 &= \max \left\{ \frac{A}{y_0}, \frac{x_0}{y_0} \right\} = \frac{A}{y_0} < A, \\
 x_2 &= \max \left\{ \frac{A}{x_1}, \frac{y_1}{x_1} \right\} = \frac{Ax_0}{A} = x_0 < A, \\
 y_2 &= \max \left\{ \frac{A}{y_1}, \frac{x_1}{y_1} \right\} = \frac{A}{y_1} = y_0 < A, \\
 x_3 &= \max \left\{ \frac{A}{x_2}, \frac{y_2}{x_2} \right\} = \frac{A}{x_2} = \frac{A}{x_0} < A, \\
 y_3 &= \max \left\{ \frac{A}{y_2}, \frac{x_2}{y_2} \right\} = \frac{A}{y_2} = \frac{A}{y_0} < A, \\
 x_4 &= \max \left\{ \frac{A}{x_3}, \frac{y_3}{x_3} \right\} = \frac{A}{x_3} = x_0 < A,
 \end{aligned}$$

$$\begin{aligned}
y_4 &= \max\left\{\frac{A}{y_3}, \frac{x_3}{y_3}\right\} = \frac{A}{y_3} = y_0 < A, \\
x_5 &= \max\left\{\frac{A}{x_4}, \frac{y_4}{x_4}\right\} = \frac{A}{x_4} = \frac{A}{x_0} < A, \\
y_5 &= \max\left\{\frac{A}{y_4}, \frac{x_4}{y_4}\right\} = \frac{A}{y_4} = \frac{A}{y_0} < A, \\
&\vdots
\end{aligned} \tag{2.14}$$

$n \geq 1$, then

$$\begin{aligned}
x_n &= \left(\frac{x_0}{A^{a(n)}}\right)^{(-1)^n}, \\
y_n &= \left(\frac{y_0}{A^{a(n)}}\right)^{(-1)^n}.
\end{aligned} \tag{2.15}$$

□

Theorem 2.10. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $1 < x_0 < y_0 < A$.

$$\begin{aligned}
\text{(a)} \quad &\lim_{n \rightarrow \infty} x_{2n} = x_0, \\
&\lim_{n \rightarrow \infty} y_{2n} = y_0. \\
\text{(b)} \quad &\lim_{n \rightarrow \infty} x_{2n+1} = \frac{A}{x_0}, \\
&\lim_{n \rightarrow \infty} y_{2n+1} = \frac{A}{y_0}.
\end{aligned} \tag{2.16}$$

Proof. (a) We obtain that

$$\begin{aligned}
\lim_{n \rightarrow \infty} x_{2n} &= \lim_{n \rightarrow \infty} \left(\frac{x_0}{A^{a(2n)}}\right)^{(-1)^{2n}} = \lim_{n \rightarrow \infty} \left(\frac{x_0}{A^0}\right)^{(-1)^{2n}} = \lim_{n \rightarrow \infty} \left(\frac{x_0}{A^0}\right) = x_0, \\
\lim_{n \rightarrow \infty} y_{2n} &= \lim_{n \rightarrow \infty} \left(\frac{y_0}{A^{a(2n)}}\right)^{(-1)^{2n}} = \lim_{n \rightarrow \infty} \left(\frac{y_0}{A^0}\right)^{(-1)^{2n}} = \lim_{n \rightarrow \infty} \left(\frac{y_0}{A^0}\right) = y_0.
\end{aligned} \tag{2.17}$$

(b) Similarly we can obtain the proof of (b) as the proof of (a). □

Lemma 2.11. Let (x_0, y_0) be the initial condition of (1.1) for $0 < x_0 < 1 < y_0 < A$; there is at least an $i_0 \in \mathbb{N}$ such that every $n \in \mathbb{N}$ for $n > i_0$, $y_0/x_0^n > A$.

Proof. We consider that $x_0 < 1$ hence $\lim_{n \rightarrow \infty} (y_0/x_0^n) = \infty$ and that proves the existing of i_0 defined in hypothesis. □

Theorem 2.12. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $0 < x_0 < 1 < y_0 < A$, and i_0 is the number, defined by Lemma 2.11.

$$1 \leq n \leq i_0$$

$$x_n = A^{a(n)} (x_0)^{(-1)^n}, \quad (2.18)$$

$$1 \leq n \leq i_0$$

$$y_n = A^{a(n)} \left(\frac{y_0}{x_0^{\lfloor n/2 \rfloor}} \right)^{(-1)^n}, \quad (2.19)$$

and when $n > i_0$, the solutions will be different for every different constant A .

Proof. Let $y_0 < A$, then

$$\begin{aligned} x_1 &= \max \left\{ \frac{A}{x_0}, \frac{y_0}{x_0} \right\} = \frac{A}{x_0} > 1, \\ y_1 &= \max \left\{ \frac{A}{y_0}, \frac{x_0}{y_0} \right\} = \frac{A}{y_0} > 1, \\ x_2 &= \max \left\{ \frac{A}{x_1}, \frac{y_1}{x_1} \right\} = \frac{A}{x_1} = x_0 < 1, \\ y_2 &= \max \left\{ \frac{A}{y_1}, \frac{x_1}{y_1} \right\} = \frac{x_1}{y_1} = \frac{y_0}{x_0} > 1, \\ x_3 &= \max \left\{ \frac{A}{x_2}, \frac{y_2}{x_2} \right\} = \frac{A}{x_2} = \frac{A}{x_0} > 1, \\ y_3 &= \max \left\{ \frac{A}{y_2}, \frac{x_2}{y_2} \right\} = \frac{A}{y_2} = \frac{Ax_0}{y_0}, \\ x_4 &= \max \left\{ \frac{A}{x_3}, \frac{y_3}{x_3} \right\} = \frac{A}{x_3} = x_0 < 1, \\ y_4 &= \max \left\{ \frac{A}{y_3}, \frac{x_3}{y_3} \right\} = \frac{x_3}{y_3} = \frac{y_0}{x_0^2}, \\ x_5 &= \max \left\{ \frac{A}{x_4}, \frac{y_4}{x_4} \right\} = \frac{A}{x_4} = \frac{A}{x_0} > 1, \\ y_5 &= \max \left\{ \frac{A}{y_4}, \frac{x_4}{y_4} \right\} = \frac{A}{y_4} = \frac{Ax_0^2}{y_0}, \end{aligned} \quad (2.20)$$

$$1 \leq n \leq i_0,$$

$$\begin{aligned} x_n &= A^{a(n)} (x_0)^{(-1)^n}, \\ y_n &= A^{a(n)} \left(\frac{y_0}{x_0^{\lfloor [n/2] \rfloor}} \right)^{(-1)^n}. \end{aligned} \quad (2.21)$$

□

Lemma 2.13. Let (x_0, y_0) be the initial condition of (1.1) for $0 < y_0 < 1 < x_0 < A$; there is at least an $i_0 \in \mathbb{N}$ such that every $n \in \mathbb{N}$ for $n > i_0$, $x_0/y_0^n > A$.

Proof. Similarly we can obtain the proof as the proof of Lemma 2.11. □

Theorem 2.14. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $0 < y_0 < 1 < x_0 < A$, and i_0 is the number, defined by Lemma 2.13.

$$1 \leq n \leq i_0$$

$$x_n = A^{a(n)} \left(\frac{x_0}{y_0^{\lfloor [n/2] \rfloor}} \right)^{(-1)^n}, \quad (2.22)$$

$$1 \leq n \leq i_0$$

$$y_n = A^{a(n)} (y_0)^{(-1)^n}, \quad (2.23)$$

and when $n > i_0$, the solutions will be different for every different constant A .

Proof. Similarly we can obtain the proof of be as the proof of Theorem 2.12. □

Lemma 2.15. Let (x_0, y_0) be the initial condition of (1.1) for $0 < y_0 < x_0 < 1$; there is at least an $i_0 \in \mathbb{N}$ such that every $n \in \mathbb{N}$ for $n > i_0$, $(x_0/y_0)^n > A$.

Proof. We consider that $y_0 < x_0$ hence $\lim_{n \rightarrow \infty} (x_0/y_0)^n = \infty$ and that proofs the existing of i_0 defined in hypothesis. □

Theorem 2.16. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $0 < y_0 < x_0 < 1$, $A > 1$, and i_0 is the number, defined by Lemma 2.15.

$$x_1 = A/x_0 \text{ and if } 1 < n \leq i_0,$$

$$x_n = A^{a(n)} \left(\frac{x_0}{y_0} \right)^{\lfloor [n/2] \rfloor (-1)^n}, \quad (2.24)$$

$$y_1 = A/y_0, \text{ and if } 1 < n \leq i_0,$$

$$y_n = A^{a(n)} \left(\frac{y_0}{x_0} \right)^{(-1)^n}, \quad (2.25)$$

and when $n > i_0$, the solutions will be different for every different constant A .

Proof. Let $y_0 < A$, then

$$\begin{aligned}
 x_1 &= \max\left\{\frac{A}{x_0}, \frac{y_0}{x_0}\right\} = \frac{A}{x_0} > 1, \\
 y_1 &= \max\left\{\frac{A}{y_0}, \frac{x_0}{y_0}\right\} = \frac{A}{y_0} > 1, \\
 x_2 &= \max\left\{\frac{A}{x_1}, \frac{y_1}{x_1}\right\} = \frac{y_1}{x_1} = \frac{x_0}{y_0} > 1, \\
 y_2 &= \max\left\{\frac{A}{y_1}, \frac{x_1}{y_1}\right\} = \frac{x_1}{y_1} = \frac{y_0}{x_0} < 1, \\
 x_3 &= \max\left\{\frac{A}{x_2}, \frac{y_2}{x_2}\right\} = \frac{A}{x_2} = \frac{Ay_0}{x_0}, \\
 y_3 &= \max\left\{\frac{A}{y_2}, \frac{x_2}{y_2}\right\} = \frac{A}{y_2} = \frac{Ax_0}{y_0} > 1, \\
 x_4 &= \max\left\{\frac{A}{x_3}, \frac{y_3}{x_3}\right\} = \frac{y_3}{x_3} = \frac{x_0^2}{y_0^2} > 1, \\
 y_4 &= \max\left\{\frac{A}{y_3}, \frac{x_3}{y_3}\right\} = \frac{A}{y_3} = \frac{y_0}{x_0} < 1,
 \end{aligned} \tag{2.26}$$

$n > 1$ then

$$x_n = A^{a(n)} \left(\frac{x_0}{y_0} \right)^{[n/2](-1)^n}, \tag{2.27}$$

$n \geq 1$ then

$$y_n = A^{a(n)} \left(\frac{y_0}{x_0} \right)^{(-1)^n}. \tag{2.28}$$

□

Lemma 2.17. Let (x_0, y_0) be the initial condition of (1.1) for $0 < y_0 < x_0 < 1$; there is at least an $i_0 \in \mathbb{N}$ such that every $n \in \mathbb{N}$ for $n > i_0$, $(y_0/x_0)^n > A$.

Proof. Similarly we can obtain the proof as the proof of Lemma 2.15. □

Theorem 2.18. Let (x_n, y_n) be the solution of the system of difference equations (1.1) for $0 < x_0 < y_0 < 1$, $A > 1$, and i_0 is the number, defined by Lemma 2.17.

$x_1 = A/x_0$, and if $1 < n \leq i_0$,

$$x_n = A^{a(n)} \left(\frac{x_0}{y_0} \right)^{(-1)^n}, \tag{2.29}$$

$y_1 = A/y_0$, and if $1 < n \leq i_0$,

$$y_n = A^{a(n)} \left(\frac{y_0}{x_0} \right)^{[n/2](-1)^n}, \quad (2.30)$$

and when $n > i_0$, the solutions will be different for every different constant A .

Proof. Similarly we can obtain the proof as the proof of Theorem 2.16, which completes the proofs of theorems. $\square$

References

- [1] A. M. Amleh, *Boundedness periodicity and stability of some difference equations*, Ph.D. thesis, University of Rhode Island, Kingston, Rhode Island, USA, 1998.
- [2] C. Çinar, S. Stević, and İ. Yalçinkaya, "On positive solutions of a reciprocal difference equation with minimum," *Journal of Applied Mathematics & Computing*, vol. 17, no. 1-2, pp. 307–314, 2005.
- [3] S. N. Elaydi, *An Introduction to Difference Equations*, Undergraduate Texts in Mathematics, Springer, New York, NY, USA, 1996.
- [4] J. Feuer, "Periodic solutions of the Lyness max equation," *Journal of Mathematical Analysis and Applications*, vol. 288, no. 1, pp. 147–160, 2003.
- [5] A. Gelişken, C. Çinar, and R. Karataş, "A note on the periodicity of the Lyness max equation," *Advances in Difference Equations*, vol. 2008, Article ID 651747, 5 pages, 2008.
- [6] A. Gelişken, C. Çinar, and İ. Yalçinkaya, "On the periodicity of a difference equation with maximum," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 820629, 11 pages, 2008.
- [7] E. Janowski, V. L. Kocic, G. Ladas, and G. Tzanetopoulos, "Global behavior of solutions of $x_{n+1} = [\max\{x_n^k, A\}]/x_{n-1}$," *Journal of Difference Equations and Applications*, vol. 3, no. 3-4, pp. 297–310, 1998.
- [8] M. R. S. Kulenević and G. Ladas, *Dynamics of Second Order Rational Difference Equations with Open Problems and Conjecture*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 2002.
- [9] D. P. Mishev, W. T. Patula, and H. D. Voulou, "A reciprocal difference equation with maximum," *Computers & Mathematics with Applications*, vol. 43, no. 8-9, pp. 1021–1026, 2002.
- [10] L. A. Moybé and A. S. Kapadia, *Difference Equations with Public Health Applications*, CRC Press, New York, NY, USA, 2000.
- [11] G. Papaschinopoulos and V. Hatzifilippidis, "On a max difference equation," *Journal of Mathematical Analysis and Applications*, vol. 258, no. 1, pp. 258–268, 2001.
- [12] G. Papaschinopoulos, J. Schinas, and V. Hatzifilippidis, "Global behavior of the solutions of a max-equation and of a system of two max-equations," *Journal of Computational Analysis and Applications*, vol. 5, no. 2, pp. 237–254, 2003.
- [13] W. T. Patula and H. D. Voulou, "On a max type recurrence relation with periodic coefficients," *Journal of Difference Equations and Applications*, vol. 10, no. 3, pp. 329–338, 2004.
- [14] G. Stefanidou and G. Papaschinopoulos, "The periodic nature of the positive solutions of a nonlinear fuzzy max-difference equation," *Information Sciences*, vol. 176, no. 24, pp. 3694–3710, 2006.
- [15] S. Stević, "On the recursive sequence $x_{n+1} = \max\{c, x_n^p/x_{n-1}^p\}$," *Applied Mathematics Letters*, vol. 21, no. 8, pp. 791–796, 2008.
- [16] D. Şimsek, C. Çinar, and İ. Yalçinkaya, "On the solutions of the difference equation $x_{n+1} = \max\{1/x_{n-1}, x_{n-1}\}$," *International Journal of Contemporary Mathematical Sciences*, vol. 1, no. 9–12, pp. 481–487, 2006.
- [17] D. Şimsek, B. Demir, and A. S. Kurbanlı, " $x_{n+1} = \max\{1/x_n, y_n/x_n\}, y_{n+1} = \max\{1/y_n, x_n/y_n\}$," *Denklemler Sisteminin Çözümleri Üzerine*. In press.
- [18] C. T. Teixeira, *Existence stability boundedness and periodicity of some difference equations*, Ph.D. thesis, University of Rhode Island, Kingston, Rhode Island, USA, 2000.
- [19] S. Valicenti, *Periodicity and global attractivity of some difference equations*, Ph.D. thesis, University of Rhode Island, Kingston, Rhode Island, USA, 1999.
- [20] H. D. Voulou, "On the periodic character of some difference equations," *Journal of Difference Equations and Applications*, vol. 8, no. 9, pp. 799–810, 2002.

- [21] H. D. Voulfov, "Periodic solutions to a difference equation with maximum," *Proceedings of the American Mathematical Society*, vol. 131, no. 7, pp. 2155–2160, 2003.
- [22] İ. Yalçınkaya, B. D. Iričanin, and C. Çinar, "On a max-type difference equation," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 47264, 10 pages, 2007.
- [23] İ. Yalçınkaya, C. Çinar, and M. Atalay, "On the solutions of systems of difference equations," *Advances in Difference Equations*, vol. 2008, Article ID 143943, 9 pages, 2008.
- [24] X. Yan, X. Liao, and C. Li, "On a difference equation with maximum," *Applied Mathematics and Computation*, vol. 181, pp. 1–5, 2006.