Research Article

***K-nacci* Sequences in Finite Triangle Groups**

Erdal Karaduman and Ömür Deveci

Department of Mathematics, Faculty of Science, Atatürk University, 25240 Erzurum, Turkey

Correspondence should be addressed to Erdal Karaduman, eduman@atauni.edu.tr

Received 10 March 2009; Revised 16 July 2009; Accepted 14 September 2009

Recommended by Leonid Shaikhet

A *k-nacci* sequence in a finite group is a sequence of group elements $x_0, x_1, x_2, \dots, x_n, \dots$ for which, given an initial (seed) set $x_0, x_1, x_2, \dots, x_{j-1}$, each element is defined by $x_n = x_0 x_1 \dots x_{n-1}$, for $j \leq n < k$, and $x_n = x_{n-k} x_{n-k+1} \dots x_{n-1}$, for $n \geq k$. We also require that the initial elements of the sequence, $x_0, x_1, x_2, \dots, x_{j-1}$, generate the group, thus forcing the *k-nacci* sequence to reflect the structure of the group. The *K-nacci* sequence of a group generated by $x_0, x_1, x_2, \dots, x_{j-1}$ is denoted by $F_k(G; x_0, x_1, \dots, x_{j-1})$ and its period is denoted by $P_k(G; x_0, x_1, \dots, x_{j-1})$. In this paper, we obtain the period of *K-nacci* sequences in finite polyhedral groups and the extended triangle groups.

Copyright © 2009 E. Karaduman and Ö. Deveci. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The Fibonacci sequences and their related higher-order (tribonacci, quaternacci, *k-nacci*) are generally viewed as sequences of integers. In [1] the Fibonacci length of a 2-generator group is defined, thus extending the idea of forming a sequence of group elements based on a Fibonacci-like recurrence relation first introduced by Wall in [2]. There he considered the Fibonacci length of the cyclic group C_n . The concept of Fibonacci length for more than two generators has also been considered, see, for example [3, 4]. Also, the theory has been expanded to the nilpotent groups, see, for example [5–7]. Other works on Fibonacci length are discussed in, for example, [8–12]. Knox proved that the periods of *k-nacci* (*k*-step Fibonacci) sequences in dihedral groups are equal to $2k + 2$ [13]. Campbell and Campbell, examined the behaviour of the Fibonacci length of the finite polyhedral, binary polyhedral groups, and related groups in [14].

This paper discusses the period of *k-nacci* Fibonacci sequences in the polyhedral groups $(2, 2, 2)$, $(n, 2, 2)$, $(2, n, 2)$, $(2, 2, n)$ for any n and in the extended triangle groups $E(2, 2, 2)$, $E(n, 2, 2)$, $E(2, n, 2)$, $E(2, 2, n)$ for any $n > 2$. We consider polyhedral groups both as 2-generator and as 3-generator groups. A 2-step Fibonacci sequence in the integers modulo m can be written as $F_2(\mathbb{Z}_m; 0, 1)$. A 2-step Fibonacci sequence of group elements is called

a *Fibonacci sequence of a finite group*. A finite group G is *k-nacci sequenceable* if there exists a *k-nacci* sequence of G such that every element of the group appears in the sequence. A sequence of group elements is *periodic* if, after a certain point, it consists only of repetitions of a fixed subsequence. The number of elements in the repeating subsequence is called the *period of the sequence*. For example, the sequence $x_0, x_1, x_2, x_3, x_4, x_1, x_2, x_3, x_4, x_1, x_2, x_3, x_4, \dots$ is periodic after the initial element x_0 and has period 4. A sequence of group elements is *simply periodic* with period k if the first k elements in the sequence form a repeating subsequence. For example, the sequence $x_0, x_1, x_2, x_3, x_4, x_0, x_1, x_2, x_3, x_4, \dots$ is simply periodic with period 5. It is important to note that the Fibonacci length depends on the chosen generating n -tuple for a group.

Definition 1.1. For a finitely generated group $G = \langle A \rangle$ where $A = \{a_1, a_2, \dots, a_n\}$ the sequence $x_i = a_{i+1}$, $0 \leq i \leq n-1$, $x_{i+n} = \prod_{j=1}^n x_{i+j-1}$, $i \geq 0$, is called the *Fibonacci orbit* of G with respect to the generating set A , denoted $F_A(G)$.

Notice that the orbit of a k -generated group is a *k-nacci* sequence. The orbits of $(n, 2, 2)$, $(2, n, 2)$, $(2, 2, n)$ for any $n > 2$ and $E(2, q, 2)$ for any $q > 2$ are studied in [14].

2. The Groups $(2, 2, 2)$, $(n, 2, 2)$, $(2, n, 2)$, and $(2, 2, n)$

Definition 2.1. The *polyhedral group* (l, m, n) for $l, m, n > 1$ is defined by the presentation

$$\langle x, y, z : x^l = y^m = z^n = xyz = e \rangle \quad (2.1)$$

or

$$\langle x, y : x^l = y^m = (xy)^n = e \rangle. \quad (2.2)$$

The *polyhedral group* (l, m, n) is finite if and only if the number

$$\mu = lmn \left(\frac{1}{l} + \frac{1}{m} + \frac{1}{n} - 1 \right) = mn + nl + lm - lmn \quad (2.3)$$

is positive, that is, in the case $(2, 2, n)$, $(2, 3, 3)$, $(2, 3, 3)$, $(2, 3, 4)$, $(2, 3, 5)$. Its order is $2lmn/\mu$. Using Tietze transformations, we may show that $(l, m, n) \cong (m, n, l) \cong (n, l, m)$. For more information on these groups see [15] and [16, pages 67–68]. The groups considered in Theorems 2.3 and 2.4 are the same group, namely, D_n , the dihedral group of $2n$ elements, except the generators x, y , and z are different from one theorem to the other.

Theorem 2.2. Let G_2 be the group defined by the presentation $G_2 = \langle x, y, z : x^2 = y^2 = z^2 = xyz = e \rangle$. Then $P_k(G_2, x, y, z) = k + 1$.

Proof. Firstly, let us consider the 2-generator case. Notice that G_2 is $Z_2 \oplus Z_2$ and $P_k(Z_2; 0, 1) = k + 1$. Under these identifications, since the period of a Fibonacci sequence in a direct product of groups is the least common multiple of the periods in each the factors we get $P_k(G_2; x, y) = k + 1$. On the other hand, since $z = xy$ the formulas in the “three generator case” with recurrences of period $k + 1$ are the same as the formulas the two generator case as long as $k \geq 4$. $\square$

Theorem 2.3. Let $G_n, n > 2$, be the group defined by the presentation $\langle x, y, z : x^n = y^2 = z^2 = xyz = e \rangle$. Then $P_k(G_n; x, y, z) = 2k + 2$.

Proof. Let us consider the 3-generator case. We first note that the orders of x, y , and z are $n, 2, 2$, respectively. If $k = 2$, we have the sequence

$$x, y, z, yz, zyz, z, x, y, \dots, \quad (2.4)$$

which has period 6. If $k = 3$, we have the sequence

$$x, y, z, xyz = e, yz, zyz, z, e, x, y, z, \dots, \quad (2.5)$$

which has period 8. If $k \geq 4$, the first k elements of sequence are

$$x_0 = x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = xyz, \quad x_4 = (xyz)^2, \dots, \quad x_{k-1} = (xyz)^{2^{k-3}}. \quad (2.6)$$

Thus, using the above information the sequence reduces to

$$x_0 = x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = e, \quad x_4 = e, \dots, e, \quad (2.7)$$

where $x_j = e$ for $3 \leq j \leq k-1$. Thus,

$$\begin{aligned} x_k &= \prod_{i=0}^{k-1} x_i = (xy)^{2^{k-2}} = e, & x_{k+1} &= \prod_{i=1}^k x_i = yz = x^{n-1}, & x_{k+2} &= \prod_{i=2}^{k+1} x_i = zyz = xz, \\ x_{k+3} &= \prod_{i=3}^{k+2} x_i = z, & x_{k+4} &= \prod_{i=4}^{k+3} x_i = e, \dots, e. \end{aligned} \quad (2.8)$$

It follows that $x_{k+j} = e$ for $4 \leq j \leq k$. We also have,

$$\begin{aligned} x_{k+k+1} &= \prod_{i=k+1}^{k+k} x_i = e, & x_{k+k+2} &= \prod_{i=k+2}^{k+k+1} x_i = x, \\ x_{k+k+3} &= \prod_{i=k+3}^{k+k+2} x_i = y, & x_{k+k+4} &= \prod_{i=k+4}^{k+k+3} x_i = z. \end{aligned} \quad (2.9)$$

Since the elements succeeding $x_{2k+2}, x_{2k+3}, x_{2k+4}$, depend on x, y , and z for their values, the cycle begins again with the $2k + 2$ nd element; that is, $x_0 = x_{2k+2}, x_1 = x_{2k+3}, x_2 = x_{2k+4}, \dots$. Thus, $P_k(G_n; x, y, z) = 2k + 2$.

Similarly, it is easy to show that for 2-generator, $P_k(G_n; x, y, z) = 2k + 2$ in $(n, 2, 2)$, and it can be shown that $P_k(G_n; x, y, z) = 2k + 2$ for $(2, n, 2)$.

Because of $(n, 2, 2) \cong (2, n, 2) \cong (2, 2, n) \cong D_n$ for any $n > 2$ and using Tietze transformations we can obtain the same presentation for this groups, it is easy to show that for 2-generator $P_k(G_n; x, y) = 2k + 2$ in the groups $(n, 2, 2)$, $(2, n, 2)$, and $(2, 2, n)$. $\square$

Theorem 2.4. Let G_n , $n > 2$, be the group defined by the presentation $\langle x, y, z : x^2 = y^2 = z^n = xyz = e \rangle$

- (i) $P_2(G_n; y, x, z) = 6$:
(ii)

$$P_4(G_n; x, y, z) = \begin{cases} n\left(\frac{5}{2}\right), & n \equiv 0 \pmod{4}, \\ 5n, & n \equiv 2 \pmod{4}, \\ 10n, & \text{otherwise}, \end{cases} \quad (2.10)$$

- (iii) $k \geq 5$.

(1) If there is no $t \in [3, k-2]$ such that t is a odd factor of n , then

$$P_k(G_n; x, y, z) = \begin{cases} n\left(\frac{k+1}{2}\right), & n \equiv 0 \pmod{4}, \\ n(k+1), & n \equiv 2 \pmod{4}, \\ 2n(k+1), & \text{otherwise}. \end{cases} \quad (2.11)$$

(2) Let α be the biggest odd factor of n in $[3, k-2]$. Then two cases occur:

- (i') if $\alpha 3^j \notin [3, k-2]$ for $j \in \mathbb{N}$, then

$$P_k(G_n; x, y, z) = \begin{cases} \alpha\left(n\left(\frac{k+1}{2}\right)\right), & n \equiv 0 \pmod{4}, \\ \alpha(n(k+1)), & n \equiv 2 \pmod{4}, \\ \alpha(2n(k+1)), & \text{otherwise}; \end{cases} \quad (2.12)$$

- (ii') if β is the biggest odd number which is in $[3, k-2]$ and $\beta = \alpha 3^j$ for $j \in \mathbb{N}$, then

$$P_k(G_n; x, y, z) = \begin{cases} \beta\left(n\left(\frac{k+1}{2}\right)\right), & n \equiv 0 \pmod{4}, \\ \beta(n(k+1)), & n \equiv 2 \pmod{4}, \\ \beta(2n(k+1)), & \text{otherwise}. \end{cases} \quad (2.13)$$

Proof. We consider G_n as D_n , the dihedral group of $2n$ elements. Now D_n being the group of symmetries of the regular polygon with n elements admits a presentation as the group generated by the two matrices:

$$a := \begin{pmatrix} \cos\left(\frac{2\pi}{n}\right) & -\sin\left(\frac{2\pi}{n}\right) \\ \sin\left(\frac{2\pi}{n}\right) & \cos\left(\frac{2\pi}{n}\right) \end{pmatrix}, \quad b := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.14)$$

Under these identifications, we can take $z = \begin{pmatrix} \cos(2\pi/n) & -\sin(2\pi/n) \\ \sin(2\pi/n) & \cos(2\pi/n) \end{pmatrix}$, $y = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, and $x = \begin{pmatrix} \cos(2\pi/n) & -\sin(2\pi/n) \\ -\sin(2\pi/n) & -\cos(2\pi/n) \end{pmatrix}$.

(i) If $k = 2$, we have the sequence

$$\begin{aligned} x_0 = y, \quad x_1 = x, \quad x_2 = z, \quad x_3 = \begin{pmatrix} \cos\left(\frac{4\pi}{n}\right) & -\sin\left(\frac{4\pi}{n}\right) \\ -\sin\left(\frac{4\pi}{n}\right) & -\cos\left(\frac{4\pi}{n}\right) \end{pmatrix} = xz, \quad x_4 = x, \\ x_5 = \begin{pmatrix} \cos\left(\frac{2\pi}{n}\right) & \sin\left(\frac{2\pi}{n}\right) \\ -\sin\left(\frac{2\pi}{n}\right) & \cos\left(\frac{2\pi}{n}\right) \end{pmatrix} = xy, \quad x_6 = y, \quad x_7 = x, \quad x_8 = z, \dots \end{aligned} \quad (2.15)$$

Thus we get $P_2(G_n; y, x, z) = 6$.

(ii) If $k = 4$, we have the sequence

$$\begin{aligned} x, y, z, xyz = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = e, \quad (xyz)^2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = e, \quad \begin{pmatrix} \cos\left(\frac{2\pi}{n}\right) & -\sin\left(\frac{2\pi}{n}\right) \\ -\sin\left(\frac{2\pi}{n}\right) & -\cos\left(\frac{2\pi}{n}\right) \end{pmatrix} = x, \\ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = y, \quad \begin{pmatrix} \cos\left(\frac{2\pi}{n}\right) & \sin\left(\frac{2\pi}{n}\right) \\ -\sin\left(\frac{2\pi}{n}\right) & \cos\left(\frac{2\pi}{n}\right) \end{pmatrix} = xy, \quad \begin{pmatrix} \cos\left(\frac{4\pi}{n}\right) & \sin\left(\frac{4\pi}{n}\right) \\ -\sin\left(\frac{4\pi}{n}\right) & \cos\left(\frac{4\pi}{n}\right) \end{pmatrix} = z^{-2}, \\ \begin{pmatrix} \cos\left(\frac{6\pi}{n}\right) & \sin\left(\frac{6\pi}{n}\right) \\ \sin\left(\frac{6\pi}{n}\right) & -\cos\left(\frac{6\pi}{n}\right) \end{pmatrix} = z^4 x, \quad \begin{pmatrix} \cos\left(\frac{14\pi}{n}\right) & \sin\left(\frac{14\pi}{n}\right) \\ \sin\left(\frac{14\pi}{n}\right) & -\cos\left(\frac{14\pi}{n}\right) \end{pmatrix} = z^8 x, \\ \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = y, \quad \begin{pmatrix} \cos\left(\frac{2\pi}{n}\right) & -\sin\left(\frac{2\pi}{n}\right) \\ \sin\left(\frac{2\pi}{n}\right) & \cos\left(\frac{2\pi}{n}\right) \end{pmatrix} = z, \quad \begin{pmatrix} \cos\left(\frac{8\pi}{n}\right) & -\sin\left(\frac{8\pi}{n}\right) \\ \sin\left(\frac{8\pi}{n}\right) & \cos\left(\frac{8\pi}{n}\right) \end{pmatrix} = z^4, \\ \begin{pmatrix} \cos\left(\frac{24\pi}{n}\right) & -\sin\left(\frac{24\pi}{n}\right) \\ \sin\left(\frac{24\pi}{n}\right) & \cos\left(\frac{24\pi}{n}\right) \end{pmatrix} = z^{12}, \quad \begin{pmatrix} \cos\left(\frac{34\pi}{n}\right) & -\sin\left(\frac{34\pi}{n}\right) \\ -\sin\left(\frac{34\pi}{n}\right) & -\cos\left(\frac{34\pi}{n}\right) \end{pmatrix} = xz^{16}, \dots \end{aligned} \quad (2.16)$$

Now we consider what happens to the 4-*nacci* sequence when we have a section of the form $\dots, z^\tau x, zx, z, \dots$:

$$\begin{aligned} z^\tau x, zx &= y, z, z^\epsilon, z^{\tau+\epsilon}, xz^{2\epsilon+\tau}, y, xzx = xy, xz^{\epsilon+2}x = z^{-(\epsilon+2)}, \\ xz^{3\epsilon+\tau+4}x &= z^{-(3\epsilon+\tau+4)}, z^{4\epsilon+\tau+8}x, zx = y, z, \dots \end{aligned} \quad (2.17)$$

The 4-*nacci* sequence can be said to form layers of length 10. Using the above, the 4-*nacci* sequence becomes

$$\begin{aligned} x_0 &= x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = e, \dots, \\ x_{10} &= z^8x, \quad x_{11} = zx = y, \quad x_{12} = z, \quad x_{13} = z^4, \dots, \\ x_{20} &= z^{32}x, \quad x_{21} = zx = y, \quad x_{22} = z, \quad x_{23} = z^8, \dots, \\ x_{10i} &= z^{8i^2}x, \quad x_{10i+1} = zx = y, \quad x_{10i+2} = z, \quad x_{10i+3} = z^{4i}, \dots, \end{aligned} \quad (2.18)$$

where $z^{8i^2} = \begin{pmatrix} \cos 8i^2(2\pi/n) & -\sin 8i^2(2\pi/n) \\ \sin 8i^2(2\pi/n) & \cos 8i^2(2\pi/n) \end{pmatrix}$ and $z^{4i} = \begin{pmatrix} \cos 4i(2\pi/n) & -\sin 4i(2\pi/n) \\ \sin 4i(2\pi/n) & \cos 4i(2\pi/n) \end{pmatrix}$.

So, we need the smallest $i \in N$ such that $8i^2 = nv_1$ and $4i = nv_2$ for $v_1, v_2 \in N$.

If $n \equiv 0 \pmod{4}$, $z^{8i^2} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = n/4$.

Thus, $10i = (5/2)n$ and $P_4 = (G_n; x, y, z) = n((k+1)/2) = (5/2)n$.

If $n \equiv 2 \pmod{4}$, $z^{8i^2} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = n/2$.

Thus, $10i = 5n$ and $P_4 = (G_n; x, y, z) = n(k+1) = 5n$.

If $n \equiv 1 \pmod{4}$ or $n \equiv 3 \pmod{4}$, $z^{8i^2} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = n$.

Thus, $10i = 10n$ and $P_4 = (G_n; x, y, z) = 2n(k+1) = 10n$.

(iii) If $k \geq 5$, the first $k+1$ elements of the sequence are

$$x_0 = x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = z^n, \quad x_4 = z^{2n}, \dots, \quad x_k = z^{2^{k-3}n}. \quad (2.19)$$

Thus, using the above information, the sequence reduces

$$x_0 = x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = e, \quad x_4 = e, \dots, e, \quad (2.20)$$

where $x_j = e$ for $3 \leq j \leq k$.

Now we consider what happens to the k -*nacci* sequence when we have a section of the form $\dots, z^\tau x, zx, z, \dots$:

$$\begin{aligned} x_{2k+2} &= \prod_{i=k+2}^{2k+1} x_i = z^\tau x, \quad x_{2k+2+1} = \prod_{i=k+3}^{2k+2} x_i = zx = y, \quad x_{2k+2+2} = \prod_{i=k+4}^{2k+3} x_i = z, \\ x_{2k+2+3} &= \prod_{i=k+5}^{2k+4} x_i = z^\epsilon, \quad x_{2k+2+4} = \prod_{i=k+6}^{2k+5} x_i = z^c, \quad x_{2k+2+5} = \prod_{i=k+7}^{2k+6} x_i = z^{u_1}, \dots, \\ x_{2k+2+k} &= \prod_{i=2k+2}^{3k+1} x_i = z^{u_{k-4}}, \dots \end{aligned} \quad (2.21)$$

The k -nacci sequence can be said to form layers of length $(2k+2)$. Using the above, the k -nacci sequence becomes

$$\begin{aligned} x_0 &= x, & x_1 &= y, & x_2 &= z, & x_3 &= e, \dots, & x_k &= z^{2^{k-3}n} = e, \dots, \\ x_{i(2k+2)} &= z^T x, & x_{i(2k+2)+1} &= zx, & x_{i(2k+2)+2} &= z, & x_{i(2k+2)+3} &= z^{4i}, \\ x_{i(2k+2)+4} &= z^{8i^2+4i}, & x_{i(2k+2)+5} &= z^{u_1}, \dots, & x_{i(2k+2)+k} &= z^{u_{k-4}}, \dots \end{aligned} \quad (2.22)$$

So, we need the smallest $i \in N$ such that $4i = nv_1$ and $8i^2 + 4i = nv_2$ for $v_1, v_2 \in N$.

(1) If there is no $t \in [3, k-2]$ such that t is an odd factor of n , there are 3 subcases.

Case 1. If $n \equiv 0 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = n/4$. So, we get $P_k = (G_n; x, y, z) = n((k+1)/2)$ since $i(2k+2) = n((k+1)/2)$ (where by $n \mid \tau$ we mean that n divides τ).

Case 2. If $n \equiv 2 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = n/2$. So, we get $P_k = (G_n; x, y, z) = n(k+1)$ since $i(2k+2) = n(k+1)$.

Case 3. If $n \equiv 1 \pmod{4}$ or $n \equiv 3 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = n$. So, we get $P_k = (G_n; x, y, z) = 2n(k+1)$ since $i(2k+2) = 2n(k+1)$.

(2) Let α odd be the biggest factor of n in $[3, k-2]$. Then two cases occur:

(i') If $\alpha 3^j \notin [3, k-2]$ for $j \in N$, then there are 3 subcases.

Case 1. If $n \equiv 0 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = \alpha(n/4)$. So, we get $P_k = (G_n; x, y, z) = \alpha(n((k+1)/2))$ since $i(2k+2) = \alpha(n((k+1)/2))$.

Case 2. If $n \equiv 2 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = \alpha(n/2)$. So, we get $P_k = (G_n; x, y, z) = \alpha(n(k+1))$ since $i(2k+2) = \alpha(n(k+1))$.

Case 3. If $n \equiv 1 \pmod{4}$ or $n \equiv 3 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = \alpha n$. So, we get $P_k = (G_n; x, y, z) = \alpha(2n(k+1))$ since $i(2k+2) = \alpha(2n(k+1))$.

(ii') If β is the biggest odd number which is in $[3, k-2]$ and $\beta = \alpha 3^j$ for $j \in N$, then there are 3 subcases.

Case 1. If $n \equiv 0 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = \beta(n/4)$. So, we get $P_k = (G_n; x, y, z) = \beta(n((k+1)/2))$ since $i(2k+2) = \beta(n((k+1)/2))$.

Case 2. If $n \equiv 2 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = \beta(n/2)$. So, we get $P_k = (G_n; x, y, z) = \beta(n(k+1))$ since $i(2k+2) = \beta(n(k+1))$.

Case 3. If $n \equiv 1 \pmod{4}$ or $n \equiv 3 \pmod{4}$ and $n \mid \tau$, $n \mid u_1, \dots, n \mid u_{k-4}$, $z^{4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, and $z^{8i^2+4i} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ for $i = \beta n$. So, we get $P_k = (G_n; x, y, z) = \beta(2n(k+1))$ since $i(2k+2) = \beta(2n(k+1))$.

This completes the proof. $\square$

In the case of 2-generator the group has the presentation $\langle x, y : x^2 = y^2 = (xy)^n = e \rangle$ and the period is the same as in the 3-generator case and proof is similar.

3. The Groups $E(2, 2, 2)$, $E(n, 2, 2)$, $E(2, n, 2)$, and $E(2, 2, n)$

Definition 3.1. The extended triangle group $E(p, q, r)$, for $p, q, r > 1$, is defined by the presentation

$$\langle x, y, z : x^2 = y^2 = z^2 = (xy)^p = (yz)^q = (zx)^r = e \rangle. \quad (3.1)$$

The *extended triangle groups* are a very important class of groups closely linked to automorphism groups of regular maps, see [17]. The *triangle groups* (*polyhedral groups*), (p, q, r) are index two subgroups of *extended triangle groups*. To see this, let $X = xy$, $Y = yz$ and $Z = zx$ in $E(p, q, r)$ and then use the obvious epimorphism. We get the following three cases for $E(p, q, r)$:

- (1) the Euclidean case if $1/p + 1/q + 1/r = 1$,
- (2) the elliptic case if $1/p + 1/q + 1/r > 1$,
- (3) the hyperbolic case if $1/p + 1/q + 1/r < 1$.

The group $E(p, q, r)$ is finite if and only if $1/p + 1/q + 1/r > 1$.

For more information on these groups, see [14, 18].

Theorem 3.2. Let E_2 be the group defined by the presentation $\langle x, y, z : x^2 = y^2 = z^2 = (xy)^2 = (yz)^2 = (zx)^2 = e \rangle$. Then $P_k(E_2; x, y, z) = k + 1$ for $k > 2$.

Proof. Since E_2 can be identified with $Z_2 \oplus Z_2 \oplus Z_2$ and x, y, z with $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$, respectively, from a similar argument applied to Theorem 2.2, we get $P_k(E_2; x, y, z) = k + 1$. $\square$

Theorem 3.3. Let $E_n, n > 2$, be the group defined by the presentation $\langle x, y, z : x^2 = y^2 = z^2 = (xy)^2 = (yz)^2 = (zx)^n = e \rangle$

(i)

$$P_{4,5}(E_n; x, y, z) = \begin{cases} n\left(\frac{k+1}{2}\right), & n \equiv 0 \pmod{4}, \\ n(k+1), & n \equiv 2 \pmod{4}, \\ 2n(k+1), & \text{otherwise;} \end{cases} \quad (3.2)$$

(ii) let $k \geq 6$.

(1) If there is no $t \in [3, k-3]$ such that t is an odd factor of n , then

$$P_k(E_n; x, y, z) = \begin{cases} n\left(\frac{k+1}{2}\right), & n \equiv 0 \pmod{4}, \\ n(k+1), & n \equiv 2 \pmod{4}, \\ 2n(k+1), & \text{otherwise.} \end{cases} \quad (3.3)$$

(2) Let α be the biggest odd factor of n in $[3, k-3]$. Then two cases occur:

(i') if $\alpha 3^j \notin [3, k-3]$ for $j \in \mathbb{N}$, then

$$P_k(E_n; x, y, z) = \begin{cases} \alpha\left(n\left(\frac{k+1}{2}\right)\right), & n \equiv 0 \pmod{4}, \\ \alpha(n(k+1)), & n \equiv 2 \pmod{4}, \\ \alpha(2n(k+1)), & \text{otherwise;} \end{cases} \quad (3.4)$$

(ii') if β is the biggest odd number which is in $[3, k-3]$ and $\beta = \alpha 3^j$ for $j \in \mathbb{N}$, then

$$P_k(E_r; x, y, z) = \begin{cases} \beta\left(n\left(\frac{k+1}{2}\right)\right), & n \equiv 0 \pmod{4}, \\ \beta(n(k+1)), & n \equiv 2 \pmod{4}, \\ \beta(2n(k+1)), & \text{otherwise.} \end{cases} \quad (3.5)$$

Proof. Since y has order 2 and commutes with x and z it follows that $E_n = Z_2 \oplus D_n$. As a group of matrices, the can be identified with a group of 3×3 matrices of form

$$\begin{pmatrix} \pm 1 & 0 \\ 0 & a \end{pmatrix}, \quad (3.6)$$

where a is a 2×2 matrix in dihedral group generated by a and b shown at (2.14). Here,

$$x = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos\left(\frac{2\pi}{n}\right) & -\sin\left(\frac{2\pi}{n}\right) \\ 0 & -\sin\left(\frac{2\pi}{n}\right) & -\cos\left(\frac{2\pi}{n}\right) \end{pmatrix}, \quad y = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad z = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \quad (3.7)$$

Now, since the period of a Fibonacci sequence in a direct product of groups is the least common multiple of the periods in each the factors and from a similar argument applied to Theorem 2.4 the proof is done. $\square$

Acknowledgment

The authors thank the referees for their valuable suggestions which improved the presentation of the paper.

References

- [1] C. M. Campbell, H. Doostie, and E. F. Robertson, "Fibonacci length of generating pairs in groups," in *Applications of Fibonacci Numbers*, G. E. Bergum, Ed., vol. 3, pp. 27–35, Kluwer Academic Publisher, Dordrecht, The Netherlands, 1990.
- [2] D. D. Wall, "Fibonacci series module m ," *The American Mathematical Monthly*, vol. 67, pp. 525–532, 1969.
- [3] C. M. Campbell and P. P. Campbell, "On the Fibonacci length of powers of dihedral groups," in *Applications of Fibonacci Numbers*, F. T. Howard, Ed., vol. 9, pp. 69–85, Kluwer Academic Publisher, Dordrecht, The Netherlands, 2004.
- [4] H. Doostie and C. M. Campbell, "Fibonacci length of automorphism groups involving Tribonacci numbers," *Vietnam Journal of Mathematics*, vol. 28, no. 1, pp. 57–65, 2000.
- [5] H. Aydın and R. Dikici, "General Fibonacci sequences in finite groups," *The Fibonacci Quarterly*, vol. 36, no. 3, pp. 216–221, 1998.
- [6] H. Aydın and G. C. Smith, "Finite p -quotients of some cyclically presented groups," *Journal of the London Mathematical Society*, vol. 49, no. 1, pp. 83–92, 1994.
- [7] E. Karaduman and H. Aydın, "On the relationship between the recurrences in nilpotent groups and the binomial formula," *Indian Journal of Pure and Applied Mathematics*, vol. 35, no. 9, pp. 1093–1103, 2004.
- [8] C. M. Campbell and P. P. Campbell, "The Fibonacci length of certain centro-polyhedral groups," *Journal of Applied Mathematics & Computing*, vol. 19, no. 1-2, pp. 231–240, 2005.
- [9] H. Doostie and M. Maghasedi, "Fibonacci length of direct products of groups," *Vietnam Journal of Mathematics*, vol. 33, no. 2, pp. 189–197, 2005.
- [10] H. Doostie and M. Hashemi, "Fibonacci lengths involving the Wall number $k(n)$," *Journal of Applied Mathematics & Computing*, vol. 20, no. 1-2, pp. 171–180, 2006.
- [11] E. Özkan, "On truncated Fibonacci sequences," *Indian Journal of Pure and Applied Mathematics*, vol. 38, no. 4, pp. 241–251, 2007.
- [12] H. J. Wilcox, "Fibonacci sequences of period n in groups," *The Fibonacci Quarterly*, vol. 24, no. 4, pp. 356–361, 1986.
- [13] Steven W. Knox, "Fibonacci sequences in finite groups," *The Fibonacci Quarterly*, vol. 30, no. 2, pp. 116–120, 1992.
- [14] C. M. Campbell and P. P. Campbell, "The Fibonacci lengths of binary polyhedral groups and related groups," *Congressus Numerantium*, vol. 194, pp. 95–102, 2009.
- [15] J. H. Conway, H. S. M. Coxeter, and G. C. Shephard, "The centre of a finitely generated group," *Tensor. New Series*, vol. 25, pp. 405–418, 1972.
- [16] H. S. M. Coxeter and W. O. J. Moser, *Generators and Relations for Discrete Groups*, Springer, Berlin, Germany, 3rd edition, 1972.
- [17] M. D. E. Conder, "Group actions on graphs, maps and surfaces with maximum symmetry," in *Groups St. Andrews 2001 in Oxford*, C. M. Campbell, E. F. Robertson, and G. C. Smith, Eds., vol. 1, pp. 63–91, Cambridge University Press, Cambridge, UK, 2003.
- [18] M. A. Albar and W. M. Al-Hamdan, "The triangle groups," *Rendiconti del Seminario Matematico della Università di Padova*, vol. 89, pp. 103–111, 1993.