Research Article

Permanence of a Discrete Nonlinear Prey-Competition System with Delays

Hongying Lu

*School of Mathematics and Quantitative Economics, Dongbei University of Finance and Economics,
 Dalian, Liaoning 116025, China*

Correspondence should be addressed to Hongying Lu, hongyinglu543@163.com

Received 3 July 2009; Accepted 16 September 2009

Recommended by Xue-Zhong He

A discrete nonlinear prey-competition system with m -preys and $(n-m)$ -predators and delays is considered. Two sets of sufficient conditions on the permanence of the system are obtained. One set is delay independent, while the other set is delay dependent.

Copyright © 2009 Hongying Lu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In this paper, we investigate the following discrete nonlinear prey-competition system with delays:

$$\begin{aligned}
 x_i(k+1) &= x_i(k) \exp \left[r_i(k) - \sum_{j=1}^n a_{ij}(k) x_j^{\alpha_{ij}}(k) - \sum_{j=1}^n b_{ij}(k) x_j^{\beta_{ij}}(k - \tau_{ij}(k)) \right], \\
 &\quad i = 1, 2, \dots, m, \\
 x_i(k+1) &= x_i(k) \exp \left[-r_i(k) + \sum_{j=1}^m a_{ij}(k) x_j^{\alpha_{ij}}(k) + \sum_{j=1}^m b_{ij}(k) x_j^{\beta_{ij}}(k - \tau_{ij}(k)) \right. \\
 &\quad \left. - \sum_{j=m+1}^n a_{ij}(k) x_j^{\alpha_{ij}}(k) - \sum_{j=m+1}^n b_{ij}(k) x_j^{\beta_{ij}}(k - \tau_{ij}(k)) \right], \\
 &\quad i = m+1, 2, \dots, n,
 \end{aligned} \tag{1.1}$$

where $x_i(k)$ ($i = 1, 2, \dots, m$) is the density of prey species i at k th generation, $x_i(k)$ ($i = m + 1, \dots, n$) is the density of predator species i at k th generation. In this system, the competition among predator species and among prey species is simultaneously considered. For more background and biological adjustments of system (1.1), we can see [1–5] and the references cited therein.

Throughout this paper, we always assume that for all $i, j = 1, 2, \dots, n$,

(H₁) $r_i(k), a_{ij}(k), b_{ij}(k)$ are all bounded nonnegative sequences and $a_{ii}^l \geq 0, b_{ii}^l \geq 0, a_{ii}^l + b_{ii}^l > 0$. Here, for any bounded sequence $f^u = \sup_{k \in N} f(k), f^l = \inf_{k \in N} f(k)$;

(H₂) $\tau_{ij}(k)$ are bounded nonnegative integer sequences, and α_{ij}, β_{ij} are all positive constants.

By a solution of system (1.1), we mean a sequence $\{x_1(k), \dots, x_n(k)\}$ which defined for $N = \{0, 1, \dots\}$ and which satisfies system (1.1) for $N = \{0, 1, \dots\}$. Motivated by application of system (1.1) in population dynamics, we assume that solutions of system (1.1) satisfy the following initial conditions:

$$x_i(\theta) = \phi_i(\theta), \quad \theta \in N[-\tau, 0] = [-\tau, -\tau + 1, \dots, 0], \quad \phi_i(0) > 0, \quad (1.2)$$

where $\tau = \max\{\tau_{ij}(k), i, j = 1, 2, \dots, n\}$. The exponential forms of system (1.1) assure that the solution of system (1.1) with initial conditions (1.2) remains positive.

Recently, Chen et al. in [1] proposed the following nonlinear prey-competition system with delays:

$$\begin{aligned} \dot{x}_i(t) &= x_i(t) \left[r_i(t) - \sum_{j=1}^n a_{ij}(t) x_j^{\alpha_{ij}}(t) - \sum_{j=1}^n b_{ij}(t) x_j^{\beta_{ij}}(t - \tau_{ij}(t)) \right], \quad i = 1, 2, \dots, m, \\ \dot{x}_i(t) &= x_i(t) \left[-r_i(t) + \sum_{j=1}^m a_{ij}(t) x_j^{\alpha_{ij}}(t) + \sum_{j=1}^m b_{ij}(t) x_j^{\beta_{ij}}(t - \tau_{ij}(t)) \right. \\ &\quad \left. - \sum_{j=m+1}^n a_{ij}(t) x_j^{\alpha_{ij}}(t) - \sum_{j=m+1}^n b_{ij}(t) x_j^{\beta_{ij}}(t - \tau_{ij}(t)) \right], \quad i = m + 1, 2, \dots, n. \end{aligned} \quad (1.3)$$

By using Gaines and Mawhins continuation theorem of coincidence degree theory and by constructing an appropriate Lyapunov functional, they obtained a set of sufficient conditions which guarantee the existence and global attractivity of positive periodic solutions of the system (1.3). In addition, sufficient conditions are obtained for the permanence of the system (1.3) in [2].

On the other hand, though most population dynamics are based on continuous models governed by differential equations, the discrete time models are more appropriate than the continuous ones when the size of the population is rarely small or the population has nonoverlapping generations [3–15]. Therefore, it is reasonable to study discrete time prey-competition models governed by difference equations.

As we know, a more important theme that interested mathematicians as well as biologists is whether all species in a multispecies community would survive in the long run, that is, whether the ecosystems are permanent. In fact, no such work has been done for system (1.1).

The main purpose of this paper is, by developing the analytical technique of [4, 8, 16], to obtain two sets of sufficient conditions which guarantee the permanence of system (1.1).

2. Main Results

Firstly, we introduce a definition and some lemmas which will be useful in the proof of the main results of this section.

Definition 2.1. System (1.1) is said to be permanent, if there are positive constants m and M , such that each positive solution $(x_1(k), \dots, x_n(k))$ of system (1.1) satisfies

$$m \leq \liminf_{k \rightarrow +\infty} x_i(k) \leq \limsup_{k \rightarrow +\infty} x_i(k) \leq M, \quad i = 1, 2, \dots, n. \quad (2.1)$$

Lemma 2.2 (see [8]). Assume that $\{x(k)\}$ satisfies $x(k) > 0$ and

$$x(k+1) \leq x(k) \exp\{r(k)(1 - ax(k))\} \quad (2.2)$$

for $k \in [k_1, +\infty)$, where a is a positive constant. Then

$$\limsup_{k \rightarrow +\infty} x(k) \leq \frac{1}{ar^u} \exp(r^u - 1). \quad (2.3)$$

Lemma 2.3 (see [8]). Assume that $\{x(k)\}$ satisfies

$$x(k+1) \geq x(k) \exp\{r(k)(1 - ax(k))\}, \quad k \geq K_0, \quad (2.4)$$

$\limsup_{k \rightarrow +\infty} x(k) \leq x^*$ and $x(K_0) > 0$, where a is a constant such that $ax^* > 1$ and $K_0 \in \mathbb{N}$. Then

$$\liminf_{k \rightarrow +\infty} x(k) \geq \frac{1}{a} \exp(r^u(1 - ax^*)). \quad (2.5)$$

For system (1.1), we will consider two cases, $a_{ii}^l > 0, b_{ii}^l \geq 0$ and $a_{ii}^l \geq 0, b_{ii}^l > 0$ respectively, and then we obtain Lemmas 2.4–2.6.

Lemma 2.4. Assume that $a_{ii}^l > 0$. Then for every positive solution $(x_1(k), \dots, x_n(k))$ of system (1.1) with initial condition (1.2), one has

$$\limsup_{k \rightarrow +\infty} x_i(k) \leq M_i, \quad i = 1, 2, \dots, n, \quad (2.6)$$

where

$$\begin{aligned} M_i &= \left(\frac{1}{\alpha_{ii} a_{ii}^l} \right)^{1/\alpha_{ii}} \exp \left[r_i^u - \frac{1}{\alpha_{ii}} \right], \quad i = 1, 2, \dots, m, \\ M_i &= \left(\frac{1}{\alpha_{ii} a_{ii}^l} \right)^{1/\alpha_{ii}} \exp \left[-r_i^l + \sum_{j=1}^n a_{ij}^u M_j^{\alpha_{ij}} + \sum_{j=1}^n b_{ij}^u M_j^{\beta_{ij}} - \frac{1}{\alpha_{ii}} \right], \quad i = m+1, \dots, n. \end{aligned} \quad (2.7)$$

Proof. Let $x(k) = (x_1(k), \dots, x_n(k))$ be any positive solution of system (1.1) with initial condition (1.2), for $i = 1, 2, \dots, m$, it follows from system (1.1) that

$$x_i(k+1) \leq x_i(k) \exp [r_i(k) - a_{ii}(k) x_i^{\alpha_{ii}}(k)], \quad (2.8)$$

thus

$$x_i^{\alpha_{ii}}(k+1) \leq x_i^{\alpha_{ii}}(k) \exp [\alpha_{ii}(r_i(k) - a_{ii}(k) x_i^{\alpha_{ii}}(k))]. \quad (2.9)$$

Let $u_i(k) = x_i^{\alpha_{ii}}(k)$, we can have

$$\begin{aligned} u_i(k+1) &\leq u_i(k) \exp [\alpha_{ii}(r_i(k) - a_{ii}(k) u_i(k))] \\ &\leq u_i(k) \exp \left[\alpha_{ii} r_i(k) \left(1 - \frac{a_{ii}^l}{r_i^u} u_i(k) \right) \right]. \end{aligned} \quad (2.10)$$

By applying Lemma 2.2 to (2.10), we obtain

$$\limsup_{k \rightarrow +\infty} u_i(k) \leq \left(\frac{1}{\alpha_{ii} a_{ii}^l} \right) \exp [\alpha_{ii} r_i^u - 1] \doteq L_i; \quad (2.11)$$

so, we immediately get

$$\limsup_{k \rightarrow +\infty} x_i(k) \leq M_i, \quad i = 1, 2, \dots, m. \quad (2.12)$$

For any $\varepsilon > 0$ small enough, it follows from (2.12) that there exists enough large K_1 such that for all $i = 1, 2, \dots, m$ and $k \geq K_1$

$$x_i(k) \leq M_i + \varepsilon. \quad (2.13)$$

For $i = m + 1, \dots, n$ and $k \geq K_1 + \tau$, (2.13) combining with the i -th equation of system (1.1) leads to

$$x_i(k+1) \leq x_i(k) \exp \left[-r_i(k) + \sum_{j=1}^m a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k)x_i^{\alpha_{ii}}(k) \right], \quad (2.14)$$

thus

$$x_i^{\alpha_{ii}}(k+1) \leq x_i^{\alpha_{ii}}(k) \exp \left[\alpha_{ii} \left(-r_i(k) + \sum_{j=1}^m a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k)x_i^{\alpha_{ii}}(k) \right) \right]. \quad (2.15)$$

Similarly, let $u_i(k) = x_i^{\alpha_{ii}}(k)$, we get

$$\begin{aligned} u_i(k+1) &\leq u_i(k) \exp \left[\alpha_{ii} \left(-r_i(k) + \sum_{j=1}^m a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k)u_i(k) \right) \right] \\ &\leq u_i(k) \exp \left[\alpha_{ii} \left(-r_i(k) + \sum_{j=1}^m a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} \right) \right. \\ &\quad \left. \times \left(1 - \frac{a_{ii}^l}{-r_i^l + \sum_{j=1}^m a_{ij}^u(M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^u(M_j + \varepsilon)^{\beta_{ij}}} u_i(k) \right) \right]. \end{aligned} \quad (2.16)$$

By using (2.16), for $i = m + 1, \dots, n$, according to Lemma 2.2, it follows that

$$\limsup_{k \rightarrow +\infty} u_i(k) \leq \left(\frac{1}{\alpha_{ii} a_{ii}^l} \right) \exp \left[\alpha_{ii} \left(-r_i^l + \sum_{j=1}^n a_{ij}^u(M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^n b_{ij}^u(M_j + \varepsilon)^{\beta_{ij}} \right) - 1 \right]; \quad (2.17)$$

setting $\varepsilon \rightarrow 0$ in above inequality, we have

$$\limsup_{k \rightarrow +\infty} u_i(k) \leq \left(\frac{1}{\alpha_{ii} a_{ii}^l} \right) \exp \left[\alpha_{ii} \left(-r_i^l + \sum_{j=1}^n a_{ij}^u M_j^{\alpha_{ij}} + \sum_{j=1}^n b_{ij}^u M_j^{\beta_{ij}} \right) - 1 \right] \doteq L_i, \quad (2.18)$$

then

$$\limsup_{k \rightarrow +\infty} x_i(k) \leq M_i, \quad i = m+1, \dots, n. \quad (2.19)$$

This completes the proof. $\square$

For convenience, we introduce the following notation.

For $i = 1, 2, \dots, m$

$$\begin{aligned} A_i &= \frac{a_{ii}^u}{r_i^l - \sum_{j=1, j \neq i}^n a_{ij}^u M_j^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}^u M_j^{\beta_{ij}}}, \\ R_i^u &= r_i^u - \sum_{j=1, j \neq i}^n a_{ij}^l M_j^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}^l M_j^{\beta_{ij}}. \end{aligned} \quad (2.20)$$

For $i = m+1, \dots, n$

$$\begin{aligned} A_i &= \frac{a_{ii}^u}{-r_i^u + \sum_{j=1}^m a_{ij}^l m_j^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^l m_j^{\beta_{ij}} - \sum_{j=m+1, j \neq i}^n a_{ij}^u M_j^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}^u M_j^{\beta_{ij}}}, \\ R_i^u &= -r_i^l + \sum_{j=1}^m a_{ij}^u m_j^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^u m_j^{\beta_{ij}} - \sum_{j=m+1, j \neq i}^n a_{ij}^l M_j^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}^l M_j^{\beta_{ij}}. \end{aligned} \quad (2.21)$$

Lemma 2.5. Assume that $a_{ii}^l > 0$ and

$$\min_{1 \leq i \leq n} L_i A_i > 1, \quad (2.22)$$

hold. Then for any positive solution $(x_1(k), \dots, x_n(k))$ of system (1.1) with initial condition (1.2), one has

$$\liminf_{k \rightarrow +\infty} x_i(k) \geq m_i, \quad (2.23)$$

where

$$m_i = \left(\frac{1}{A_i} \right)^{1/\alpha_{ii}} \exp[R_i^u(1 - A_i L_i)], \quad i = 1, 2, \dots, n. \quad (2.24)$$

Proof. Let $x(k) = (x_1(k), \dots, x_n(k))$ be any positive solution of system (1.1) with initial condition (1.2). From Lemma 2.4, we know that there exists $K_2 > K_1 + \tau$, such that for $i = 1, 2, \dots, n$ and $k \geq K_2$

$$x_i(k) \leq M_i + \varepsilon. \quad (2.25)$$

For $i = 1, \dots, m$ and $k \geq K_2 + \tau$, (2.25) combining with the i -th equation of system (1.1) lead to

$$x_i(k+1) \geq x_i(k) \exp \left[r_i(k) - \sum_{j=1, j \neq i}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k)x_i^{\alpha_{ii}}(k) \right], \quad (2.26)$$

thus

$$x_i^{\alpha_{ii}}(k+1) \geq x_i^{\alpha_{ii}}(k) \exp \left[\alpha_{ii} \left(r_i(k) - \sum_{j=1, j \neq i}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k)x_i^{\alpha_{ii}}(k) \right) \right]; \quad (2.27)$$

let $u_i(k) = x_i^{\alpha_{ii}}(k)$, we can have

$$\begin{aligned} u_i(k+1) &\geq u_i(k) \exp \left[\alpha_{ii} \left(r_i(k) - \sum_{j=1, j \neq i}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k)u_i(k) \right) \right] \\ &\geq u_i(k) \exp[\alpha_{ii} R_{i\varepsilon}(k)(1 - A_{i\varepsilon} u_i(k))], \end{aligned} \quad (2.28)$$

where

$$\begin{aligned} R_{i\varepsilon}(k) &= r_i(k) - \sum_{j=1, j \neq i}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}}, \\ A_{i\varepsilon} &= \frac{a_{ii}^u}{r_i^l - \sum_{j=1, j \neq i}^n a_{ij}^u(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}^u(M_j + \varepsilon)^{\beta_{ij}}}. \end{aligned} \quad (2.29)$$

According to Lemma 2.3, we obtain

$$\liminf_{k \rightarrow +\infty} u_i(k) \geq \frac{1}{A_{i\varepsilon}} \exp[\alpha_{ii} R_{i\varepsilon}^u (1 - A_{i\varepsilon} L_i)], \quad (2.30)$$

where

$$R_{i\varepsilon}^u = r_i^u - \sum_{j=1, j \neq i}^n a_{ij}^l (M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}^l (M_j + \varepsilon)^{\beta_{ij}}. \quad (2.31)$$

Setting $\varepsilon \rightarrow 0$ in (2.30) leads to

$$\liminf_{k \rightarrow +\infty} u_i(k) \geq \frac{1}{A_i} \exp[\alpha_{ii} R_i^u (1 - A_i L_i)], \quad (2.32)$$

therefore

$$\liminf_{k \rightarrow +\infty} x_i(k) \geq m_i, \quad i = 1, 2, \dots, m. \quad (2.33)$$

For any $\varepsilon > 0$ small enough, it follows from (2.33) that there exists enough large $K_3 > K_2 + \tau$ such that for all $i = 1, \dots, m$ and $k \geq K_3$

$$x_i(k) \geq m_i - \varepsilon, \quad (2.34)$$

and so, for $i = m+1, \dots, n$ and $k \geq K_3 + \tau$, it follows from system (1.1) that

$$\begin{aligned} x_i(k+1) &\geq x_i(k) \exp \left[-r_i(k) + \sum_{j=1}^m a_{ij}(k) (m_j - \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k) (m_j - \varepsilon)^{\beta_{ij}} \right. \\ &\quad \left. - \sum_{j=m+1, j \neq i}^n a_{ij}(k) (M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}(k) (M_j + \varepsilon)^{\beta_{ij}} - a_{ii}(k) x_i^{\alpha_{ii}}(k) \right] \\ &\geq x_i(k) \exp [R_{i\varepsilon}(k) (1 - A_{i\varepsilon} x_i^{\alpha_{ii}}(k))], \end{aligned} \quad (2.35)$$

where

$$\begin{aligned}
 R_{i\varepsilon}(k) &= -r_i(k) + \sum_{j=1}^m a_{ij}(k)(m_j - \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k)(m_j - \varepsilon)^{\beta_{ij}} \\
 &\quad - \sum_{j=m+1, j \neq i}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}}, \\
 A_{i\varepsilon} &= a_{ii}^u / \left\{ -r_i^u + \sum_{j=1}^m a_{ij}^l(m_j - \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^n b_{ij}^l(m_j - \varepsilon)^{\beta_{ij}} \right. \\
 &\quad \left. - \sum_{j=m+1, j \neq i}^n a_{ij}^u(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}^u(M_j + \varepsilon)^{\beta_{ij}} \right\},
 \end{aligned} \tag{2.36}$$

by using (2.35), similarly to the analysis of (2.33), for $i = m+1, \dots, n$

$$\liminf_{k \rightarrow +\infty} u_i(k) \geq \frac{1}{A_i} \exp[\alpha_{ii} R_i^u (1 - A_i L_i)], \tag{2.37}$$

and therefore, we easily get

$$\liminf_{k \rightarrow +\infty} x_i(k) \geq m_i, \quad i = m+1, \dots, n. \tag{2.38}$$

This ends the proof of Lemma 2.5. □

Denote for $i = 1, 2, \dots, m$

$$\begin{aligned}
 \bar{L}_i &= \frac{1}{\beta_{ii} b_{ii}^l} \exp[\beta_{ii} r_i^u (\tau + 1) - 1]; \\
 \Gamma_i &= r_i^l - \sum_{j=1}^n a_{ij}^u M_j^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}^u M_j^{\beta_{ij}}; \\
 \bar{A}_i &= \frac{b_{ii}^u \exp[-\beta_{ii} \Gamma_i \tau]}{r_i^l - \sum_{j=1}^n a_{ij}^u M_j^{\alpha_{ij}} - \sum_{j=1, j \neq i}^n b_{ij}^u M_j^{\beta_{ij}}}; \\
 \bar{R}_i^u &= r_i^u - \sum_{j=1}^n a_{ij}^l M_j^{\alpha_{ij}} - \sum_{j=1, j \neq i}^n b_{ij}^l M_j^{\beta_{ij}}.
 \end{aligned} \tag{2.39}$$

For $i = m + 1, \dots, n$

$$\begin{aligned}
 \Upsilon_i &= -r_i^l + \sum_{j=1}^m a_{ij}^u M_j^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^u M_j^{\beta_{ij}}; \\
 \bar{L}_i &= \frac{1}{\beta_{ii} b_{ii}^l} \exp[\beta_{ii} \Upsilon_i(\tau + 1) - 1]; \\
 \Gamma_i &= -r_i^u + \sum_{j=1}^m a_{ij}^l m_j^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^l m_j^{\beta_{ij}} - \sum_{j=m+1}^n a_{ij}^u M_j^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}^u (M_j + \varepsilon)^{\beta_{ij}}; \\
 \bar{A}_i &= \frac{b_{ii}^u \exp[-\beta_{ii} \Gamma_i \tau]}{-r_i^u + \sum_{j=1}^m a_{ij}^l m_j^{\alpha_{ij}} + \sum_{j=1}^n b_{ij}^l m_j^{\beta_{ij}} - \sum_{j=m+1}^n a_{ij}^u M_j^{\alpha_{ij}} - \sum_{j=m+1, j \neq i}^n b_{ij}^u m_j^{\beta_{ij}}}; \\
 \bar{R}_i^u &= -r_i^l + \sum_{j=1}^m a_{ij}^u m_j^{\alpha_{ij}} + \sum_{j=1}^n b_{ij}^u m_j^{\beta_{ij}} - \sum_{j=m+1}^n a_{ij}^l M_j^{\alpha_{ij}} - \sum_{j=m+1, j \neq i}^n b_{ij}^l m_j^{\beta_{ij}}.
 \end{aligned} \tag{2.40}$$

Lemma 2.6. Assume that $b_{ii}^l > 0$ and

$$(\bar{H}_3) \min_{1 \leq i \leq n} \bar{L}_i \bar{A}_i > 1, \tag{2.41}$$

hold. Then for any positive solution $(x_1(k), \dots, x_n(k))$ of system (1.1) with initial condition (1.2), one has

$$\bar{m}_i \leq \liminf_{k \rightarrow +\infty} x_i(k) \leq \limsup_{k \rightarrow +\infty} x_i(k) \leq \bar{M}_i, \quad i = 1, 2, \dots, n. \tag{2.42}$$

where

$$\bar{M}_i = \bar{L}_i^{1/\beta_{ii}}, \quad \bar{m}_i = \left(\frac{1}{\bar{A}_i} \right)^{1/\beta_{ii}} \exp[\bar{R}_i^u (1 - \bar{A}_i \bar{L}_i)]. \tag{2.43}$$

Proof. Let $x(k) = (x_1(k), \dots, x_n(k))$ be any positive solution of system (1.1) with initial condition (1.2), for $i = 1, 2, \dots, m$, it follows from system (1.1) that

$$x_i(k+1) \leq x_i(k) \exp[r_i(k)] \leq x_i(k) \exp[r_i^u], \tag{2.44}$$

$$x_i(k+1) \leq x_i(k) \exp[r_i(k) - b_{ii}(k) x_i^{\beta_{ii}}(k - \tau_{ii}(k))], \tag{2.45}$$

It follows from (2.44) that

$$\prod_{j=k-\tau_{ii}(k)}^{k-1} \frac{x_i(j+1)}{x_i(j)} \leq \prod_{j=k-\tau_{ii}(k)}^{k-1} \exp[r_i^u] \leq \exp[r_i^u \tau], \tag{2.46}$$

and hence

$$x_i(k - \tau_{ii}(k)) \geq x_i(k) \exp[-r_i^u \tau], \quad (2.47)$$

which, together with (2.45), produces

$$\begin{aligned} x_i(k+1) &\leq x_i(k) \exp \left[r_i(k) - b_{ii}(k) \exp[-\beta_{ii} r_i^u \tau] x_i^{\beta_{ii}}(k) \right] \\ &\leq x_i(k) \exp \left[r_i(k) \left(1 - \frac{b_{ii}^l \exp[-\beta_{ii} r_i^u \tau]}{r_i^u} x_i^{\beta_{ii}}(k) \right) \right], \end{aligned} \quad (2.48)$$

similar to the analysis of (2.11) and (2.12), for $i = 1, 2, \dots, m$

$$\limsup_{k \rightarrow +\infty} u_i(k) \leq \bar{L}_i, \quad (2.49)$$

and thus, we immediately get

$$\limsup_{k \rightarrow +\infty} x_i(k) \leq \bar{M}_i, \quad i = 1, 2, \dots, m. \quad (2.50)$$

For any $\varepsilon > 0$ small enough, it follows from (2.50) that there exists enough large $\bar{K}_1$ such that for all $i = 1, 2, \dots, m$ and $k \geq \bar{K}_1$

$$x_i(k) \leq \bar{M}_i + \varepsilon. \quad (2.51)$$

For $i = m+1, \dots, n$ and $k \geq \bar{K}_1 + \tau$, (2.51) combining with the i -th equation of system (1.1) lead to

$$\begin{aligned} x_i(k+1) &\leq x_i(k) \exp \left[-r_i^l + \sum_{j=1}^m a_{ij}^u (M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^u (M_j + \varepsilon)^{\beta_{ij}} \right] \\ &= x_i(k) \exp[\Upsilon_{ie}], \end{aligned} \quad (2.52)$$

$$\begin{aligned} x_i(k+1) &\leq x_i(k) \exp \left[-r_i(k) + \sum_{j=1}^m a_{ij}(k) (M_j + \varepsilon)^{\alpha_{ij}} \right. \\ &\quad \left. + \sum_{j=1}^m b_{ij}(k) (M_j + \varepsilon)^{\beta_{ij}} - b_{ii}(k) x_i^{\beta_{ii}}(k - \tau_{ii}(k)) \right], \end{aligned} \quad (2.53)$$

from (2.53), similar to the argument of (2.44) and (2.47), for $k \geq \bar{K}_1 + \tau$, we have

$$x_i(k - \tau_{ii}(k)) \geq x_i(k) \exp[-\Upsilon_{ie} \tau], \quad (2.54)$$

substituting (2.54) into (2.53), we get

$$x_i(k+1) \leq x_i(k) \exp \left[-r_i(k) + \sum_{j=1}^m a_{ij}(k) (M_j + \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k) (M_j + \varepsilon)^{\beta_{ij}} - b_{ii}(k) \exp[-\beta_{ii} \Upsilon_{i\varepsilon} \tau] x_i^{\beta_{ii}}(k) \right], \quad (2.55)$$

similar to the analysis of (2.18) and (2.19), for $i = m+1, \dots, n$

$$\limsup_{k \rightarrow +\infty} u_i(k) \leq \bar{L}_i, \quad (2.56)$$

then,

$$\limsup_{k \rightarrow +\infty} x_i(k) \leq \bar{M}_i, i = m+1, \dots, n. \quad (2.57)$$

For any $\varepsilon > 0$ small enough, it follows from (2.51) and (2.57) that there exists enough large $\bar{K}_2 > \bar{K}_1 + \tau$ such that for all $i = 1, 2, \dots, n$ and $k \geq \bar{K}_2$

$$x_i(k) \leq \bar{M}_i + \varepsilon. \quad (2.58)$$

Hence, for $i = 1, 2, \dots, m$, and $k \geq \bar{K}_2 + \tau$, it follows from system (1.1) that

$$x_i(k+1) \geq x_i(k) \exp \left[r_i^l - \sum_{j=1}^n a_{ij}^u (M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1}^n b_{ij}^u (M_j + \varepsilon)^{\beta_{ij}} \right] \quad (2.59)$$

$$= x_i(k) \exp[\Gamma_{i\varepsilon}],$$

$$x_i(k+1) \geq x_i(k) \exp \left[r_i(k) - \sum_{j=1}^n a_{ij}(k) (M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1, j \neq i}^n b_{ij}(k) (M_j + \varepsilon)^{\beta_{ij}} - b_{ii}(k) x_i^{\beta_{ii}}(k - \tau_{ii}(k)) \right], \quad (2.60)$$

from (2.59), similar to the argument of (2.44) and (2.47), for $k \geq \bar{K}_2 + \tau$, we have

$$x_i(k - \tau_{ii}(k)) \leq x_i(k) \exp[-\Gamma_{i\varepsilon} \tau], \quad (2.61)$$

and this combined with (2.60) gives

$$x_i(k+1) \geq x_i(k) \exp \left[r_i(k) - \sum_{j=1}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=1, j \neq i}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - b_{ii}(k) \exp[-\beta_{ii}\Gamma_{i\varepsilon}\tau] x_i^{\beta_{ii}}(k) \right]. \quad (2.62)$$

Similar to the argument of (2.32) and (2.33), for $k \geq \bar{K}_2 + \tau$, we obtain

$$\liminf_{k \rightarrow +\infty} u_i(k) \geq \frac{1}{\bar{A}_i} \exp \left[\beta_{ii} \bar{R}_i^u (1 - \bar{A}_i \bar{L}_i) \right], \quad (2.63)$$

then

$$\liminf_{k \rightarrow +\infty} x_i(k) \geq \bar{m}_i, \quad i = 1, 2, \dots, m. \quad (2.64)$$

For any $\varepsilon > 0$ small enough, it follows from (2.63) that there exists enough large $\bar{K}_3 > \bar{K}_2 + \tau$ such that for all $i = 1, \dots, m$ and $k \geq \bar{K}_3$

$$x_i(k) \geq \bar{m}_i - \varepsilon, \quad (2.65)$$

and so, for $i = m+1, \dots, n$ and $k \geq \bar{K}_3 + \tau$, it follows from system (1.1) that

$$\begin{aligned} x_i(k+1) &\geq x_i(k) \exp \left[-r_i^u + \sum_{j=1}^m a_{ij}^l(m_j - \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}^l(m_j - \varepsilon)^{\beta_{ij}} \right. \\ &\quad \left. - \sum_{j=m+1}^n a_{ij}^u(M_j + \varepsilon)^{\alpha_{ij}} - \sum_{j=m+1}^n b_{ij}^u(M_j + \varepsilon)^{\beta_{ij}} \right] = x_i(k) \exp[\Gamma_{i\varepsilon}], \\ x_i(k+1) &\geq x_i(k) \exp \left[-r_i(k) + \sum_{j=1}^m a_{ij}(k)(m_j - \varepsilon)^{\alpha_{ij}} + \sum_{j=1}^m b_{ij}(k)(m_j - \varepsilon)^{\beta_{ij}} - \sum_{j=m+1}^n a_{ij}(k)(M_j + \varepsilon)^{\alpha_{ij}} \right. \\ &\quad \left. - \sum_{j=m+1, j \neq i}^n b_{ij}(k)(M_j + \varepsilon)^{\beta_{ij}} - b_{ii}(k) x_i^{\beta_{ii}}(k - \tau_{ii}(k)) \right]. \end{aligned} \quad (2.66)$$

Similar to the argument of (2.61) and (2.62), for $k \geq \bar{K}_3 + \tau$, we have

$$\liminf_{k \rightarrow +\infty} u_i(k) \geq \frac{1}{\bar{A}_i} \exp \left[\beta_{ii} \bar{R}_i^u (1 - \bar{A}_i \bar{L}_i) \right], \quad (2.67)$$

then

$$\liminf_{k \rightarrow +\infty} x_i(k) \geq \bar{m}_i, \quad i = m+1, \dots, n. \quad (2.68)$$

This ends the proof of Lemma 2.6. $\square$

Denote (H_3)

$$a_{ii}^l > 0, \quad \min_{1 \leq i \leq n} L_i A_i > 1, \quad (2.69)$$

or

$$b_{ii}^l > 0, \quad \min_{1 \leq i \leq n} \bar{L}_i \bar{A}_i > 1, \quad (2.70)$$

Our main result in this paper is the following theorem about the permanence of system (1.1).

Theorem 2.7. *Assume that (H_1) , (H_2) , and (H_3) hold, then system (1.1) is permanent.*

Proof. Let $x(k) = (x_1(k), \dots, x_n(k))$ be any positive solution of system (1.1) with initial condition (1.2). Suppose $M = \max_{i=1, \dots, n} \{M_i, \bar{M}_i\}$, $m = \min_{i=1, \dots, n} \{m_i, \bar{m}_i\}$. By Lemmas 2.4–2.6, if system (1.1) satisfies (H_1) , (H_2) , and (H_3) , then we have

$$m \leq \liminf_{k \rightarrow +\infty} x_i(k) \leq \limsup_{k \rightarrow +\infty} x_i(k) \leq M, \quad i = 1, 2, \dots, n. \quad (2.71)$$

The proof is completed. $\square$

In this paper, we study a discrete nonlinear predator-prey system with m -preys and $(n-m)$ -predators and delays, which can be seen as the modification of the traditional Lotka-Volterra prey-competition model. From our main results, Theorem 2.7 gives two sets of sufficient conditions on the permanence of the system (1.1). One set is delay independent, while the other set is delay dependent.

Acknowledgments

The author is grateful to the Associate Editor, Professor Xue-Zhong He, and two referees for a number of helpful suggestions that have greatly improved her original submission. This research is supported by the Nation Natural Science Foundation of China (no. 10171010), Key Project on Science and Technology of Education Ministry of China (no. 01061), and Innovation Group Program of Liaoning Educational Committee No. 2007T050.

References

- [1] F. Chen, X. Xie, and J. Shi, "Existence, uniqueness and stability of positive periodic solution for a nonlinear prey-competition model with delays," *Journal of Computational and Applied Mathematics*, vol. 194, no. 2, pp. 368–387, 2006.
- [2] A. Huang, "Permanence of a nonlinear prey-competition model with delays," *Applied Mathematics and Computation*, vol. 197, no. 1, pp. 372–381, 2008.
- [3] X. Liao, S. Zhou, and Y. N. Raffoul, "On the discrete-time multi-species competition-predation system with several delays," *Applied Mathematics Letters*, vol. 21, no. 1, pp. 15–22, 2008.
- [4] X. Liao, S. Zhou, and Y. Chen, "On permanence and global stability in a general Gilpin-Ayala competition predator-prey discrete system," *Applied Mathematics and Computation*, vol. 190, no. 1, pp. 500–509, 2007.
- [5] F. Chen, L. Wu, and Z. Li, "Permanence and global attractivity of the discrete Gilpin-Ayala type population model," *Computers & Mathematics with Applications*, vol. 53, no. 8, pp. 1214–1227, 2007.
- [6] R. P. Agarwal and P. J. Y. Wong, *Advanced Topics in Difference Equations*, vol. 404 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [7] R. P. Agarwal, *Difference Equations and Inequalities: Theory, Methods, and Applications*, vol. 228 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 2nd edition, 2000.
- [8] X. Yang, "Uniform persistence and periodic solutions for a discrete predator-prey system with delays," *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 161–177, 2006.
- [9] F. Chen, "Permanence in a discrete Lotka-Volterra competition model with deviating arguments," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 5, pp. 2150–2155, 2008.
- [10] W. Yang and X. Li, "Permanence for a delayed discrete ratio-dependent predator-prey model with monotonic functional responses," *Nonlinear Analysis: Real World Applications*, vol. 10, no. 2, pp. 1068–1072, 2009.
- [11] Y. Muroya, "Partial survival and extinction of species in discrete nonautonomous Lotka-Volterra systems," *Tokyo Journal of Mathematics*, vol. 28, no. 1, pp. 189–200, 2005.
- [12] Y. Saito, W. Ma, and T. Hara, "A necessary and sufficient condition for permanence of a Lotka-Volterra discrete system with delays," *Journal of Mathematical Analysis and Applications*, vol. 256, no. 1, pp. 162–174, 2001.
- [13] L. Chen and L. Chen, "Permanence of a discrete periodic volterra model with mutual interference," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 205481, 9 pages, 2009.
- [14] M. Fazly and M. Hesaaraki, "Periodic solutions for a discrete time predator-prey system with monotone functional responses," *Comptes Rendus Mathématique*, vol. 345, no. 4, pp. 199–202, 2007.
- [15] W. Wang and Z. Lu, "Global stability of discrete models of Lotka-Volterra type," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 35, no. 8, pp. 1019–1030, 1999.
- [16] X. Meng and L. Chen, "Periodic solution and almost periodic solution for a nonautonomous Lotka-Volterra dispersal system with infinite delay," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 1, pp. 125–145, 2008.