Research Article

Permanence and Global Attractivity of a Discrete Two-Prey One-Predator Model with Infinite Delay

Zhixiang Yu and Zhong Li

College of Mathematics and Computer Science, Fuzhou University, Fuzhou, Fujian 350108, China

Correspondence should be addressed to Zhong Li, lizhong04108@163.com

Received 4 February 2009; Revised 2 July 2009; Accepted 15 August 2009

Recommended by Leonid Berezensky

A discrete two-prey one-predator model with infinite delay is proposed. A set of sufficient conditions which guarantee the permanence of the system is obtained. By constructing a suitable Lyapunov functional, we also obtain sufficient conditions ensuring the global attractivity of the system. An example together with its numerical simulation shows the feasibility of the main results.

Copyright © 2009 Z. Yu and Z. Li. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The aim of this paper is to investigate the persistence and stability property of the following discrete two-prey one-predator model with infinite delays:

$$\begin{aligned}x_1(n+1) &= x_1(n) \exp \left[a_1(n) - b_1(n)x_1(n) - c_1(n) \sum_{s=-\infty}^n H_1(n-s)x_3(s) \right], \\x_2(n+1) &= x_2(n) \exp \left[a_2(n) - b_2(n)x_2(n) - c_2(n) \sum_{s=-\infty}^n H_2(n-s)x_3(s) \right], \\x_3(n+1) &= x_3(n) \exp \left[-b_3(n)x_3(n) + d_1(n) \sum_{s=-\infty}^n H_3(n-s)x_1(s) \right. \\&\quad \left. + d_2(n) \sum_{s=-\infty}^n H_4(n-s)x_2(s) \right],\end{aligned}\tag{1.1}$$

where $x_i(n)$, $i = 1, 2$, are the densities of the prey species i at the n th generation; $x_3(n)$ is the density of the predator at the n th generation; $\{a_i(n)\}$, $\{c_i(n)\}$, $\{d_i(n)\}$ ($i = 1, 2$), $\{b_i(n)\}$

$(i = 1, 2, 3); \{H_i(n)\} (i = 1, \dots, 4)$ are all bounded nonnegative sequences such that

$$0 < a_i^l \leq a_i^u, \quad 0 < c_i^l \leq c_i^u, \quad 0 < d_i^l \leq d_i^u, \quad 0 < b_i^l \leq b_i^u, \quad \sum_{n=0}^{\infty} H_i(n) = 1. \quad (1.2)$$

Here, for any bounded sequence $\{g(n)\}$, set $g^u = \sup_{n \in \mathbb{N}} \{g(n)\}$ and $g^l = \inf_{n \in \mathbb{N}} \{g(n)\}$.

From the point of view of biology, in the sequel, we assume that

$$x_i(s) = \Phi_i(s) \geq 0, \quad \Phi_i(0) > 0, \quad i = 1, 2, 3, \quad (1.3)$$

where $s = \dots, -n, -n+1, \dots, -1, 0$. Then system (1.1) with the initial condition (1.3) has a unique positive solution $(x_1(n), x_2(n), x_3(n))^T$.

As one of the dominant themes in mathematical biology, the predator-prey relationship has been studied in a number of ways (see [1–4] and the references therein). In 1970, Parrish and Saila [5] firstly proposed the one-prey two-predator model as follows:

$$\begin{aligned} \dot{x}_1(t) &= x_1(t)(b_1 - x_1(t) - \alpha x_2(t) - \eta z(t)), \\ \dot{x}_2(t) &= x_2(t)(b_2 - \beta x_1(t) - x_2(t) - \nu z(t)), \\ \dot{z}(t) &= z(t)(-b_3 + d\eta x_1(t) + \nu x_2(t)). \end{aligned} \quad (1.4)$$

Gramer and May [6] studied the stability of the positive equilibrium of system (1.4); Takcuchi and Adachi [7] investigated the existence of the positive equilibrium and Hopf Bifurcation of the above system.

Recently, Elettrey [8] proposed the following two-prey one-predator model:

$$\begin{aligned} \dot{x}(t) &= ax(t)(1 - x(t)) - (x(t)z(t)), \\ \dot{y}(t) &= by(t)(1 - y(t)) - (y(t)z(t)), \\ \dot{z}(t) &= -cz(t)^2 + dx(t)z(t) + ey(t)z(t), \end{aligned} \quad (1.5)$$

where all the parameters in system (1.5) are positive constants. By applying differential inequality theory and iterative scheme, he showed that the unique positive equilibrium of system (1.5) is globally attractive. It is well known that a suitable ecosystem should incorporate some phase of the state of system, which is represented by time delays. Li et al. [9] studied the two-prey one-predator model with delays:

$$\begin{aligned} \dot{x}_1(t) &= x_1(t)(b_1 - x_1(t) - \alpha x_2(t) - r_1 y(t)), \\ \dot{x}_2(t) &= x_2(t)(b_2 - \beta x_1(t) - x_2(t) - r_2 y(t)), \\ \dot{y}(t) &= -b_3 y(t) + dr_1 x_1(t - \tau) y(t - \tau) + dr_2 x_2(t - \tau) y(t - \tau), \end{aligned} \quad (1.6)$$

where $x_1(t)$, $x_2(t)$, $y(t)$ are the densities of the prey and predator at the time t , respectively, $b_1, b_2, b_3, \alpha, \beta, r_1, r_2, d$ are all positive constants. They investigated the Hopf bifurcation of

system (1.6). Corresponding to system (1.5), Huang [10] proposed and studied the following system with infinite delays:

$$\begin{aligned} \dot{x}_1(t) &= ax_1(t) \left(1 - x_1(t) - \frac{1}{a} \int_{-\infty}^t K_1(t-s)x_3(s)ds \right), \\ \dot{x}_2(t) &= bx_2(t) \left(1 - x_2(t) - \frac{1}{b} \int_{-\infty}^t K_2(t-s)x_3(s)ds \right), \\ \dot{x}_3(t) &= x_3(t) \left(-cx_3(t) + d \int_{-\infty}^t K_3(t-s)x_1(s)ds + e \int_{-\infty}^t K_4(t-s)x_2(s)ds \right), \end{aligned} \quad (1.7)$$

where all the coefficients a, b, c, d, e are positive constants, and $K_i : [0, +\infty) \rightarrow (0, +\infty)$ are continue functions such that $\int_0^\infty K_i(s)ds = 1$, $i = 1, 2, 3, 4$. By applying iterative scheme, he showed that the unique positive equilibrium of the system is globally attractive.

On the other hand, it is well known that the discrete time model governed by difference equations are more appropriate than the continuous ones when the populations have nonoverlapping generations. Corresponding to traditional continuous Logistic model governed by differential equations, Mohamad and Gopalsamy [11] proposed the following single species discrete model:

$$x(n+1) = x(n) \exp \left[r(n) \left(1 - \frac{x(n)}{K(n)} \right) \right]. \quad (1.8)$$

They tried to obtain a set of sufficient conditions which ensure that (1.8) admits a unique positive and globally asymptotically stable almost periodic solution. However, Zhou and Zou [12] gave an counterexample which shows that the main results of [11] are not correct. By developing some new analysis technique, Zhou and Zou [12] obtained sufficient conditions which ensure the existence of a positive and globally asymptotically stable ω -periodic solution of system (1.8). Chen and Zhou [13] further generalized system (1.8) to the following two-species Lotka-Volterra competition system:

$$\begin{aligned} x(n+1) &= x(n) \exp \left[r_1(n) \left(1 - \frac{x(n)}{K_1(n)} - \mu_2(n)y(n) \right) \right], \\ y(n+1) &= y(n) \exp \left[r_2(n) \left(1 - \mu_1(n)x(n) - \frac{y(n)}{K_2(n)} \right) \right]. \end{aligned} \quad (1.9)$$

They obtained the sufficient conditions which guarantee the persistence of the system (1.3). Also, for the periodic case, they obtained the sufficient conditions which guarantee the existence of a globally stable periodic solution of the system. Wang and Lu [14] proposed the following Lotka-Volterra model:

$$x_i(k+1) = x_i(k) \exp \left[r_i(k) - \sum_{j=1}^n a_{ij}(k)x_j(k) \right], \quad i = 1, 2, \dots, n, \quad (1.10)$$

where $x_i(k)$ is the density of population i at k th generation; $r_i(k)$ is the growth rate of population i at k th generation; $a_{ij}(k)$ measures the intensity of intraspecific competition or interspecific action of species. By constructing a suitable Lyapunov function and using the finite covering theorem of Mathematic Analysis, they obtained a set of sufficient conditions which ensures the system to be globally asymptotically stable. Similar to the continuous ones, some scholars also argued that the discrete model should incorporate some past state of the species and thus should consider the discrete model with time delay. Recently, Chen [15] investigated the persistent property of the following discrete two species Lotka-Volterra competition model with deviating arguments:

$$\begin{aligned} x_1(n+1) &= x_1(n) \exp \left[r_1(n) \left(1 - \frac{x_1(n)}{K_1(n)} - \mu_2(n) \sum_{s=-\infty}^n H_1(n-s)x_2(s) \right) \right], \\ x_2(n+1) &= x_2(n) \exp \left[r_2(n) \left(1 - \mu_1(n) \sum_{s=-\infty}^n H_2(n-s)x_1(s) - \frac{x_2(n)}{K_2(n)} \right) \right], \end{aligned} \quad (1.11)$$

where $x_i(n)$, $i = 1, 2$, are the densities of competition species i at n th generation. By establishing a new difference inequality, Chen [15] showed that under the same conditions as that of Chen and Zhou [13], (1.11) is also permanent, which means that with some suitable restriction on the coefficients of the system, delay has no influence on the persistent property of the system. Chen [16] also investigated the persistent property of a discrete n -species nonautonomous Lotka-Volterra competitive systems with infinite delays and feedback controls. As we can see, both [15] and [16] considered the persistent property of the system, but they did not investigate the stability property of the system. Recently, Chen et al. [17] investigated the dynamic behaviors of the following general discrete nonautonomous system of plankton allelopathy with finite time delay:

$$\begin{aligned} N_1(k+1) &= N_1(k) \exp \left[r_1(k) - \sum_{l=0}^m a_{1l}(k)N_1(k-l) - \sum_{l=0}^m b_{1l}(k)N_2(k-l) \right. \\ &\quad \left. - \sum_{l=0}^m c_{1l}(k)N_1(k)N_2(k-l) \right], \\ N_2(k+1) &= N_2(k) \exp \left[r_2(k) - \sum_{l=0}^m a_{2l}(k)N_2(k-l) - \sum_{l=0}^m b_{2l}(k)N_1(k-l) \right. \\ &\quad \left. - \sum_{l=0}^m c_{2l}(k)N_2(k)N_1(k-l) \right], \end{aligned} \quad (1.12)$$

where $N_i(k)$ represents the densities of population i at the k th generation; $r_i(k)$ is the intrinsic growth rate of population i at the k th generation; $a_{il}(k)$ measures the intraspecific influence of the $(k-l)$ th generation of population i on the density of own population; $b_{il}(k)$ stands for the inter-specific influence of the $(k-l)$ th generation of population i on the density of own population and $c_{il}(k)$ stands for the effect of toxic inhibition of population i by population j at the $(k-l)$ th generation, $i, j = 1, 2$ and $i \neq j$. $\{r_i(k)\}$, $\{a_{il}(k)\}$, $\{b_{il}(k)\}$ and $\{c_{il}(k)\}$ are all bounded nonnegative sequences defined for $k \in N$. They obtained sufficient

conditions which guarantee the permanence, global attractivity and partial extinction of the above system.

Concerned with the discrete predator-prey system, by giving the detail analysis of the right hand side of the system, Yang [18] obtained a set of sufficient conditions which ensures the uniform persistence of the system investigated. Recently, Chen and Chen [19] proposed the following discrete periodic Volterra model with mutual interference and Holling II type functional response

$$\begin{aligned} x(n+1) &= x(n) \exp \left[r_1(n) - b_1(n)x(n) - \frac{c_1(n)}{k+x(n)} y^m(n) \right], \\ y(n+1) &= y(n) \exp \left[-r_2(n) - b_2(n)y(n) - \frac{c_2(n)x(n)}{k+x(n)} y^{m-1}(n) \right]. \end{aligned} \quad (1.13)$$

They also obtained sufficient conditions which ensure the permanence of the system. For more works on discrete population model, one could refer to [11–42] and the references cited therein.

However, to the best of the authors knowledge, to this day, no scholars propose and study the discrete predator-prey model with infinite delays. This motivates us to propose and study (1.1). The aim of this paper is to investigate the persistent and stability property of system (1.1).

The rest of the paper is arranged as follows: some useful lemmas are stated in the following section. Sufficient conditions which ensure the permanence and global attractivity of system (1.1) are stated and proved in Section 3. In Section 4, an example together with its numeric simulation shows the feasibility of the main results. We end this paper by a brief discussion.

2. Preliminaries

Now let us state several lemmas which will be useful in proving main results.

Lemma 2.1 (see [29]). *Assume that $\{x(n)\}$ satisfies $x(n) > 0$ and*

$$x(n+1) \leq x(n) \exp[r(n) - a(n)x(n)], \quad (2.1)$$

for $n \in \mathbb{N}$, where $\{r(n)\}$ and $\{a(n)\}$ are all positive sequences bounded above and below by positive constants. Then

$$\limsup_{n \rightarrow +\infty} x(n) \leq \frac{1}{a^l} \exp[r^u - 1]. \quad (2.2)$$

Lemma 2.2 (see [29]). *Assume that $\{x(n)\}$ satisfies*

$$x(n+1) \geq x(n) \exp[r(n) - a(n)x(n)], \quad k \geq N_0, \quad (2.3)$$

$\limsup_{n \rightarrow +\infty} x(n) \leq x^*$ and $x(N_0) > 0$, where $\{r(n)\}$ and $\{a(n)\}$ are all positive sequences bounded above and below by positive constants and $N_0 \in \mathbb{N}$. Then

$$\liminf_{n \rightarrow +\infty} x(n) \geq \frac{r^l \exp[r^l - a^u x^*]}{a^u}. \quad (2.4)$$

Lemma 2.3 (see [43]). Let $x : \mathbb{Z} \rightarrow \mathbb{R}$ be a nonnegative bounded sequences, and let $H : \mathbb{N} \rightarrow \mathbb{R}$ be a nonnegative sequences such that $\sum_{n=0}^{\infty} H(n) = 1$. Then

$$\liminf_{n \rightarrow +\infty} x(n) \leq \liminf_{n \rightarrow +\infty} \sum_{s=-\infty}^n H(n-s)x(s) \leq \limsup_{n \rightarrow +\infty} \sum_{s=-\infty}^n H(n-s)x(s) \leq \limsup_{n \rightarrow +\infty} x(n). \quad (2.5)$$

3. Main Results

Now, we investigate the persistence property and stability property of system (1.1).

Theorem 3.1. Assume that

$$a_1^l - c_1^u M_3 > 0, \quad (H_1)$$

$$a_2^l - c_2^u M_3 > 0 \quad (H_2)$$

hold, then system (1.1) is permanent, that is,

$$m_i \leq \liminf_{n \rightarrow +\infty} x_i(n) \leq \limsup_{n \rightarrow +\infty} x_i(n) \leq M_i, \quad i = 1, 2, 3, \quad (3.1)$$

where

$$\begin{aligned} M_i &= \frac{1}{b_i^l} \exp[a_i^u - 1], \quad i = 1, 2, \quad M_3 = \frac{1}{b_3^l} \exp[d_1^u M_1 + d_2^u M_2 - 1], \\ m_i &= \frac{a_i^l - c_i^u M_3}{b_i^u} \exp[a_i^l - c_i^u M_3 - b_i^u M_i], \quad i = 1, 2, \\ m_3 &= \frac{d_1^l m_1 + d_2^l m_2}{b_3} \exp[d_1^l m_1 + d_2^l m_2 - b_3^u M_3]. \end{aligned} \quad (3.2)$$

Proof. It follows from the first two equations of system (1.1) that

$$x_i(n+1) \leq x_i(n) \exp[a_i(n) - b_i(n)x_i(n)], \quad i = 1, 2. \quad (3.3)$$

So, as a consequence of Lemma 2.1, for any solution $(x_1(n), x_2(n), x_3(n))_{n=0}^{\infty}$ of system (1.1) with initial condition (1.3), one has

$$\limsup_{n \rightarrow +\infty} x_i(n) \leq \frac{1}{b_i^l} \exp[a_i^u - 1] \stackrel{\text{def}}{=} M_i, \quad i = 1, 2. \quad (3.4)$$

According to Lemma 2.3, from (3.4), we have

$$\limsup_{n \rightarrow +\infty} \sum_{s=-\infty}^n H_3(n-s)x_1(s) \leq M_1, \quad \limsup_{n \rightarrow +\infty} \sum_{s=-\infty}^n H_4(n-s)x_2(s) \leq M_2. \quad (3.5)$$

For any small positive constant $\varepsilon > 0$, it follows from (3.5) that there exists a positive integer n_1 such that for all $n \geq n_1$,

$$\sum_{s=-\infty}^n H_3(n-s)x_1(s) \leq M_1 + \varepsilon, \quad \sum_{s=-\infty}^n H_4(n-s)x_2(s) \leq M_2 + \varepsilon. \quad (3.6)$$

Thus, for all $n \geq n_1$, from the last equation of system (1.1), it follows that

$$x_3(n+1) \leq x_3(n) \exp \left[-b_3^l x_3(n) + d_1^u(M_1 + \varepsilon) + d_2^u(M_2 + \varepsilon) \right]. \quad (3.7)$$

By applying Lemma 2.1 to (3.7), we have

$$\limsup_{n \rightarrow +\infty} x_3(n) \leq \frac{1}{b_3^l} \exp \left[d_1^u(M_1 + \varepsilon) + d_2^u(M_2 + \varepsilon) - 1 \right]. \quad (3.8)$$

Setting $\varepsilon \rightarrow 0$, it follows that

$$\limsup_{n \rightarrow +\infty} x_3(n) \leq \frac{1}{b_3^l} \exp \left[d_1^u M_1 + d_2^u M_2 - 1 \right] \stackrel{\text{def}}{=} M_3. \quad (3.9)$$

Next, we show that under the assumption of Theorem 3.1,

$$\liminf_{n \rightarrow +\infty} x_i(n) \geq m_i, \quad i = 1, 2, 3. \quad (3.10)$$

According to Lemma 2.3, from (3.9) we have

$$\limsup_{n \rightarrow \infty} \sum_{s=-\infty}^n H_i(n-s)x_3(s) \leq M_3, \quad i = 1, 2. \quad (3.11)$$

Conditions (H_1) and (H_2) imply that for enough small positive constant ε_1 , we have

$$a_i^l - C_i^u(M_3 + \varepsilon_1) > 0, \quad i = 1, 2. \quad (3.12)$$

For ε_1 , it follows from (3.11) that there exists a positive integer $n_2 \geq n_1$ such that for all $n > n_2$,

$$\sum_{s=-\infty}^n H_i(n-s)x_3(s) \leq M_3 + \varepsilon_1, \quad i = 1, 2. \quad (3.13)$$

For $n > n_2$, from (3.13) and the first two equations of system (1.1), we have

$$x_i(n+1) \geq x_i(n) \exp \left[a_i^l - b_i^u x_i(n) - c_i^u (M_3 + \varepsilon_1) \right], \quad i = 1, 2. \quad (3.14)$$

Thus, according to Lemma 2.2, one has

$$\liminf_{n \rightarrow +\infty} x_i(n) \geq \frac{a_i^l - c_i^u (M_3 + \varepsilon_1)}{b_i^u} \exp \left[a_i^l - c_i^u (M_3 + \varepsilon_1) - b_i^u M_i \right], \quad i = 1, 2. \quad (3.15)$$

Setting $\varepsilon_1 \rightarrow 0$, it follows that

$$\liminf_{n \rightarrow +\infty} x_i(n) \geq \frac{a_i^l - c_i^u M_3}{b_i^u} \exp \left[a_i^l - c_i^u M_3 - b_i^u M_i \right] \stackrel{\text{def}}{=} m_i, \quad i = 1, 2. \quad (3.16)$$

According to Lemma 2.3, from (3.16) we have

$$\liminf_{n \rightarrow +\infty} \sum_{s=-\infty}^n H_3(n-s) x_1(s) \geq m_1, \quad \liminf_{n \rightarrow +\infty} \sum_{s=-\infty}^n H_4(n-s) x_2(s) \geq m_2. \quad (3.17)$$

For any $\varepsilon_2 > 0$ small enough, without loss of generality, we may assume that $\varepsilon_2 < (1/2) \min\{m_1, m_2\}$. From (3.17), it follows that there exists a n_3 , such that for all $n \geq n_3$,

$$\sum_{s=-\infty}^n H_3(n-s) x_1(s) \geq m_1 - \varepsilon_2, \quad \sum_{s=-\infty}^n H_4(n-s) x_2(s) \geq m_2 - \varepsilon_2. \quad (3.18)$$

For $n \geq n_3$, from (3.18) and the last equation of (1.1), we have

$$x_3(n+1) \geq x_3(n) \exp \left[-b_3^u x_3(n) + d_1^l (m_1 - \varepsilon_2) + d_2^l (m_2 - \varepsilon_2) \right]. \quad (3.19)$$

By applying Lemma 2.2 to (3.19), it follows that

$$\begin{aligned} \liminf_{n \rightarrow +\infty} x_3(n) &\geq \frac{d_1^l (m_1 - \varepsilon_2) + d_2^l (m_2 - \varepsilon_2)}{b_3^u} \\ &\quad \times \exp \left[d_1^l (m_1 - \varepsilon_2) + d_2^l (m_2 - \varepsilon_2) - b_3^u M_3 \right]. \end{aligned} \quad (3.20)$$

Setting $\varepsilon_2 \rightarrow 0$, it follows that

$$\liminf_{n \rightarrow +\infty} x_3(n) \geq \frac{d_1^l m_1 + d_2^l m_2}{b_3^u} \exp \left[d_1^l m_1 + d_2^l m_2 - b_3^u M_3 \right] \stackrel{\text{def}}{=} m_3. \quad (3.21)$$

This ends the proof of Theorem 3.1. □

Theorem 3.2. Assume that (H_1) and (H_2) hold. Assume further that there exist positive constants α, β, γ and δ such that

$$\alpha \min \left\{ b_1^l, \frac{2}{M_1} - b_1^u \right\} - \gamma d_1^u > \delta, \quad (H_3)$$

$$\beta \min \left\{ b_2^l, \frac{2}{M_2} - b_2^u \right\} - \gamma d_2^u > \delta, \quad (H_4)$$

$$\gamma \min \left\{ b_3^l, \frac{2}{M_3} - b_3^u \right\} - \alpha c_1^u - \beta c_2^u > \delta, \quad (H_5)$$

hold. Then for any two positive solutions $(x_1(n), x_2(n), x_3(n))^T$ and $(y_1(n), y_2(n), y_3(n))^T$ of system (1.1), one has

$$\lim_{n \rightarrow +\infty} (x_i(n) - y_i(n)) = 0, \quad i = 1, 2, 3. \quad (3.22)$$

Proof. From conditions (H_3) – (H_5) , there exists an enough small positive constant ε_3 such that

$$\alpha \min \left\{ b_1^l, \frac{2}{M_1 + \varepsilon_3} - b_1^u \right\} - \gamma d_1^u > \delta, \quad (3.23)$$

$$\beta \min \left\{ b_2^l, \frac{2}{M_2 + \varepsilon_3} - b_2^u \right\} - \gamma d_2^u > \delta, \quad (3.24)$$

$$\gamma \min \left\{ b_3^l, \frac{2}{M_3 + \varepsilon_3} - b_3^u \right\} - \alpha c_1^u - \beta c_2^u > \delta. \quad (3.25)$$

Since (H_1) and (H_2) hold, for any solutions $(x_1(n), x_2(n), x_3(n))^T$ and $(y_1(n), y_2(n), y_3(n))^T$ of system (1.1) with the initial conditions (1.3), it follows from Theorem 3.1 that

$$\limsup_{n \rightarrow +\infty} x_i(n) \leq M_i, \quad \limsup_{n \rightarrow +\infty} y_i(n) \leq M_i, \quad i = 1, 2, 3. \quad (3.26)$$

For the above ε_3 and (3.26), there exists an $n^* > 0$ such that for all $n > n^*$,

$$x_i(n) \leq M_i + \varepsilon_3, \quad y_i(n) \leq M_i + \varepsilon_3. \quad (3.27)$$

Firstly, let

$$V_{11}(n) = |\ln x_1(n) - \ln y_1(n)|. \quad (3.28)$$

Then from the first equation of the system (1.1), we have

$$\begin{aligned}
 V_{11}(n+1) &= |\ln x_1(n+1) - \ln y_1(n+1)| \\
 &\leq |\ln x_1(n) - \ln y_1(n) - b_1(n)(x_1(n) - y_1(n))| \\
 &\quad + c_1(n) \left| \sum_{s=-\infty}^n H_1(n-s) |x_3(s) - y_3(s)| \right|.
 \end{aligned} \tag{3.29}$$

Using the Mean Value Theorem, we get

$$x_1(n) - y_1(n) = \exp(\ln x_1(n)) - \exp(\ln y_1(n)) = \xi_1(n)(\ln x_1(n) - \ln y_1(n)), \tag{3.30}$$

where $\xi_1(n)$ lies between $x_1(n)$ and $y_1(n)$, then it follows that

$$\begin{aligned}
 V_{11}(n+1) &\leq |\ln x_1(n) - \ln y_1(n)| - \left(\frac{1}{\xi_1(n)} - \left| \frac{1}{\xi_1(n)} - b_1(n) \right| \right) |x_1(n) - y_1(n)| \\
 &\quad + c_1(n) \left| \sum_{s=-\infty}^n H_1(n-s) |x_3(s) - y_3(s)| \right|,
 \end{aligned} \tag{3.31}$$

and so

$$\begin{aligned}
 \Delta V_{11} &\leq - \left(\frac{1}{\xi_1(n)} - \left| \frac{1}{\xi_1(n)} - b_1(n) \right| \right) |x_1(n) - y_1(n)| \\
 &\quad + c_1(n) \left| \sum_{s=-\infty}^n H_1(n-s) |x_3(s) - y_3(s)| \right|.
 \end{aligned} \tag{3.32}$$

Secondly, let

$$V_{12}(n) = \sum_{s=0}^{\infty} H_1(s) \sum_{\theta=n-s}^{n-1} c_1(\theta+s) |x_3(\theta) - y_3(\theta)|. \tag{3.33}$$

then, similar to the aforementioned analysis, we have

$$\begin{aligned}
 \Delta V_{12} &= V_{12}(n+1) - V_{12}(n) \\
 &\leq c_1^n |x_3(n) - y_3(n)| - c_1(n) \sum_{s=-\infty}^n H_1(n-s) |x_3(s) - y_3(s)|.
 \end{aligned} \tag{3.34}$$

Now, set

$$V_1(n) = V_{11}(n) + V_{12}(n), \quad (3.35)$$

then from (3.32) and (3.34), we have

$$\begin{aligned} \Delta V_1 &= \Delta V_{11} + \Delta V_{12} \\ &\leq -\left(\frac{1}{\xi_1(n)} - \left|\frac{1}{\xi_1(n)} - b_1(n)\right|\right) |x_1(n) - y_1(n)| \\ &\quad + c_1^u |x_3(n) - y_3(n)|. \end{aligned} \quad (3.36)$$

Let

$$V_2(n) = V_{21}(n) + V_{22}(n), \quad (3.37)$$

where

$$\begin{aligned} V_{21}(n) &= |\ln x_2(n) - \ln y_2(n)|, \\ V_{22}(n) &= \sum_{s=0}^{\infty} H_2(s) \sum_{\theta=n-s}^{n-1} c_2(\theta + s) |x_3(\theta) - y_3(\theta)|. \end{aligned} \quad (3.38)$$

Similar to the aforementioned analysis, we have

$$\begin{aligned} \Delta V_2 &= \Delta V_{21} + \Delta V_{22} \\ &\leq -\left(\frac{1}{\xi_2(n)} - \left|\frac{1}{\xi_2(n)} - b_2(n)\right|\right) |x_2(n) - y_2(n)| + c_2^u |x_3(n) - y_3(n)|, \end{aligned} \quad (3.39)$$

where $\xi_2(n)$ lies between $x_2(n)$ and $y_2(n)$.

Let

$$V_3(n) = V_{31}(n) + V_{32}(n), \quad (3.40)$$

where

$$\begin{aligned} V_{31}(n) &= |\ln x_3(n) - \ln y_3(n)|, \\ V_{32}(n) &= \sum_{s=0}^{\infty} H_3(s) \sum_{\theta=n-s}^{n-1} d_1(\theta + s) |x_1(\theta) - y_1(\theta)| \\ &\quad + \sum_{s=0}^{\infty} H_4(s) \sum_{\theta=n-s}^{n-1} d_2(\theta + s) |x_2(\theta) - y_2(\theta)|. \end{aligned} \quad (3.41)$$

Similar to the aforementioned analysis, we have

$$\begin{aligned}\Delta V_3 &= \Delta V_{31} + \Delta V_{32} \\ &\leq -\left(\frac{1}{\xi_3(n)} - \left|\frac{1}{\xi_3(n)} - b_3(n)\right|\right)|x_3(n) - y_3(n)| \\ &\quad + d_1^u|x_1(n) - y_1(n)| + d_2^u|x_2(n) - y_2(n)|,\end{aligned}\tag{3.42}$$

where $\xi_3(n)$ lies between $x_3(n)$ and $y_3(n)$.

Now, we define a Lyapunou functional as follows:

$$V(n) = \alpha V_1(n) + \beta V_2(n) + \gamma V_3(n).\tag{3.43}$$

Calculating the difference of V along the solution of system (1.1), for $n > n^*$, it follows from (3.23), (3.24), (3.25), (3.27), (3.36), (3.39) and (3.42) that

$$\begin{aligned}\Delta V &\leq -\left[\alpha \min\left\{b_1^l, \frac{2}{M_1 + \varepsilon_3} - b_1^u\right\} - \gamma d_1^u\right]|x_1(n) - y_1(n)| \\ &\quad - \left[\beta \min\left\{b_2^l, \frac{2}{M_2 + \varepsilon_3} - b_2^u\right\} - \gamma d_2^u\right]|x_2(n) - y_2(n)| \\ &\quad - \left[\gamma \min\left\{b_3^l, \frac{2}{M_3 + \varepsilon_3} - b_3^u\right\} - \alpha c_1^u - \beta c_2^u\right]|x_3(n) - y_3(n)| \\ &\leq -\delta \sum_{i=1}^n |x_i(n) - y_i(n)|.\end{aligned}\tag{3.44}$$

Summating both sides of the above inequalities from n^* to n , we have

$$\sum_{p=n^*}^n (V(p+1) - V(p)) \leq -\delta \sum_{p=n^*}^n \sum_{i=1}^3 |x_i(p) - y_i(p)|,\tag{3.45}$$

which implies

$$V(n+1) + \delta \sum_{p=n^*}^n \sum_{i=1}^3 |x_i(p) - y_i(p)| \leq V(n^*).\tag{3.46}$$

It follows that

$$\sum_{p=n^*}^n \sum_{i=1}^3 |x_i(p) - y_i(p)| \leq \frac{V(n^*)}{\delta}, \quad (3.47)$$

then

$$\sum_{p=n^*}^n \sum_{i=1}^3 |x_i(p) - y_i(p)| \leq \frac{V(n^*)}{\delta} < +\infty, \quad (3.48)$$

which means that $\lim_{n \rightarrow \infty} \sum_{i=1}^3 |x_i(n) - y_i(n)| = 0$, that is

$$\lim_{n \rightarrow +\infty} (x_i(n) - y_i(n)) = 0, \quad i = 1, 2, 3. \quad (3.49)$$

This completes the proof of Theorem 3.2. □

4. Example

The following example shows the feasibility of the main results.

Example 4.1. Consider the following system:

$$\begin{aligned} x_1(n+1) &= x_1(n) \exp \left[\left(1.1 + 0.1 \sin(\sqrt{2}n) \right) - \left(1.2 + 0.2 \sin(\sqrt{2}n) \right) x_1(n) \right. \\ &\quad \left. - (0.3 + 0.2 \cos(n)) \sum_{s=-\infty}^n \frac{e-1}{e} e^{-(n-s)} x_3(s) \right], \\ x_2(n+1) &= x_2(n) \exp \left[\left(1 + 0.1 \sin(\sqrt{3}n) \right) - (1.1 + 0.1 \sin(n)) x_2(n) \right. \\ &\quad \left. - \left(0.2 + 0.02 \cos(\sqrt{3}n) \right) \sum_{s=-\infty}^n \frac{e^2-1}{e^2} e^{-2(n-s)} x_3(s) \right], \\ x_3(n+1) &= x_3(n) \exp \left[- \left(1.15 + 0.15 \sin(\sqrt{3}n) \right) x_3(n) \right. \\ &\quad \left. + \left(0.12 + 0.03 \cos(\sqrt{2}n) \right) \sum_{s=-\infty}^n \frac{e^3-1}{e^3} e^{-3(n-s)} x_1(s) \right. \\ &\quad \left. + (0.13 + 0.03 \cos(n)) \sum_{s=-\infty}^n \frac{e^4-1}{e^4} e^{-4(n-s)} x_2(s) \right]. \end{aligned} \quad (4.1)$$

Figure 1: Dynamics behaviors of system (4.1) with initial conditions $(x_1(s), x_2(s), x_3(s))^T = (0.5, 1, 0.3)^T, (1.1, 1.1, 0.1)^T, (0.9, 0.6, 0.2)^T, s = \dots, -n, -n+1, \dots, -1, 0$.

One could easily see that there exist $\alpha = 0.06$, $\beta = 0.05$, $\gamma = 0.05$ and $\delta = 0.001$ such that

$$\begin{aligned}
 a_1^l - c_1^u M_3 &= 1 - 0.5M - 3 \approx 0.7364 > 0, \\
 a_2^l - c_2^u M_3 &= 1 - 0.22M_3 \approx 0.8840 > 0, \\
 \alpha \min \left\{ b_1^l, \frac{2}{M_1} - b_1^u \right\} - \gamma d_1^u &\approx 0.0082 > \delta, \\
 \beta \min \left\{ b_2^l, \frac{2}{M_2} - b_2^u \right\} - \gamma d_2^u &\approx 0.0241 > \delta, \\
 \gamma \min \left\{ b_3^l, \frac{2}{M_3} - b_3^u \right\} - \alpha c_1^u - \beta c_2^u &\approx 0.009 > \delta.
 \end{aligned} \tag{4.2}$$

Clearly, conditions (H_1) – (H_5) are satisfied. From Theorems 3.1 and 3.2, (1.1) is permanent and globally attractive. Numeric simulation (Figure 1) strongly supports our results.

5. Discussion

In this paper, we propose a discrete two-prey one-predator model with infinite delay. Theorem 3.1 shows that to ensure the permanence of the system, one should ensure $a_i^l, i = 1, 2$ and b_3^l enough large, that is, the net birth rate of prey species and the density restriction of predator species should be large enough. We also obtain a set of sufficient conditions which ensures the global attractivity of the system.

Acknowledgments

The authors are grateful to anonymous referees for their excellent suggestions, which greatly improve the presentation of the paper. Also, this work was supported by the Program for New Century Excellent Talents in Fujian Province University (0330-003383).

References

- [1] X. Liu and L. Chen, "Complex dynamics of Holling type II Lotka-Volterra predator-prey system with impulsive perturbations on the predator," *Chaos, Solitons & Fractals*, vol. 16, no. 2, pp. 311–320, 2003.
- [2] B. Liu, Y. Zhang, and L. Chen, "Dynamic complexities of a Holling I predator-prey model concerning periodic biological and chemical control," *Chaos, Solitons & Fractals*, vol. 22, no. 1, pp. 123–134, 2004.
- [3] S. Zhang, D. Tan, and L. Chen, "Chaos in periodically forced Holling type II predator-prey system with impulsive perturbations," *Chaos, Solitons & Fractals*, vol. 28, no. 2, pp. 367–376, 2006.
- [4] F. Wang and G. Zeng, "Chaos in a Lotka-Volterra predator-prey system with periodically impulsive ratio-harvesting the prey and time delays," *Chaos, Solitons & Fractals*, vol. 32, no. 4, pp. 1499–1512, 2007.
- [5] J. D. Parrish and S. B. Saila, "Interspecific competition, predation and species diversity," *Journal of Theoretical Biology*, vol. 27, no. 2, pp. 207–220, 1970.
- [6] F. Cramer and J. May, *Chaos and Order-The Complex Structure of Living Systems*, VCH, New York, NY, USA, 1993.
- [7] Y. Takeuchi and N. Adachi, "Existence and bifurcation of stable equilibrium in two-prey, one-predator communities," *Bulletin of Mathematical Biology*, vol. 45, no. 6, pp. 877–900, 1983.
- [8] M. F. Elettrey, "Two-prey one-predator model," *Chaos, Solitons & Fractals*, vol. 39, no. 5, pp. 2018–2027, 2009.
- [9] S. Y. Li, Z. L. Xiong, and X. Wang, "Hopf Bifurcation of a two-prey one-predator system with time delays," *Journal of Nanchang University (Natural Science)*, vol. 30, no. 1, pp. 19–23, 2008.
- [10] S. Y. Huang, "Global attractivity of a delayed two-prey one-predator model," *Journal of Fuzhou University (Natural Sciences Edition)*. In press.
- [11] S. Mohamad and K. Gopalsamy, "Extreme stability and almost periodicity in a discrete logistic equation," *The Tohoku Mathematical Journal*, vol. 52, no. 1, pp. 107–125, 2000.
- [12] Z. Zhou and X. Zou, "Stable periodic solutions in a discrete periodic logistic equation," *Applied Mathematics Letters*, vol. 16, no. 2, pp. 165–171, 2003.
- [13] Y. Chen and Z. Zhou, "Stable periodic solution of a discrete periodic Lotka-Volterra competition system," *Journal of Mathematical Analysis and Applications*, vol. 277, no. 1, pp. 358–366, 2003.
- [14] W. Wang and Z. Lu, "Global stability of discrete models of Lotka-Volterra type," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 35, no. 8, pp. 1019–1030, 1999.
- [15] F. Chen, "Permanence in a discrete Lotka-Volterra competition model with deviating arguments," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 5, pp. 2150–2155, 2008.
- [16] X. Chen, "Permanence in a discrete n -species non-autonomous Lotka-Volterra competitive systems with infinite delays and feedback control," *Advances in Complex Systems*, vol. 10, no. 4, pp. 463–477, 2007.
- [17] Y. Chen, F. Chen, and Z. Li, "Dynamic behaviors of a general discrete nonautonomous system of plankton allelopathy with delays," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 310425, 22 pages, 2008.
- [18] X. Yang, "Uniform persistence and periodic solutions for a discrete predator-prey system with delays," *Journal of Mathematical Analysis and Applications*, vol. 316, no. 1, pp. 161–177, 2006.
- [19] L. Chen and L. Chen, "Permanence of a discrete periodic Volterra model with mutual interference," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 205481, 9 pages, 2009.
- [20] F. Chen, Z. Li, and Y. Huang, "Note on the permanence of a competitive system with infinite delay and feedback controls," *Nonlinear Analysis: Real World Applications*, vol. 8, no. 2, pp. 680–687, 2007.
- [21] H.-F. Huo and W.-T. Li, "Permanence and global stability for nonautonomous discrete model of plankton allelopathy," *Applied Mathematics Letters*, vol. 17, no. 9, pp. 1007–1013, 2004.
- [22] F. Chen, "Permanence and global attractivity of a discrete multispecies Lotka-Volterra competition predator-prey systems," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 3–12, 2006.

- [23] F. Chen, L. Wu, and Z. Li, "Permanence and global attractivity of the discrete Gilpin-Ayala type population model," *Computers & Mathematics with Applications*, vol. 53, no. 8, pp. 1214–1227, 2007.
- [24] F. Chen, "Permanence for the discrete mutualism model with time delays," *Mathematical and Computer Modelling*, vol. 47, no. 3-4, pp. 431–435, 2008.
- [25] F. Chen, "Permanence of a discrete n -species food-chain system with time delays," *Applied Mathematics and Computation*, vol. 185, no. 1, pp. 719–726, 2007.
- [26] Z. Li and F. Chen, "Extinction in two dimensional nonautonomous Lotka-Volterra systems with the effect of toxic substances," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 684–690, 2006.
- [27] Y. Saito, W. Ma, and T. Hara, "A necessary and sufficient condition for permanence of a Lotka-Volterra discrete system with delays," *Journal of Mathematical Analysis and Applications*, vol. 256, no. 1, pp. 162–174, 2001.
- [28] F. Chen, "Some new results on the permanence and extinction of nonautonomous Gilpin-Ayala type competition model with delays," *Nonlinear Analysis: Real World Applications*, vol. 7, no. 5, pp. 1205–1222, 2006.
- [29] F. Chen, "Permanence and global stability of nonautonomous Lotka-Volterra system with predator-prey and deviating arguments," *Applied Mathematics and Computation*, vol. 173, no. 2, pp. 1082–1100, 2006.
- [30] Y. Song and H. Tian, "Periodic and almost periodic solutions of nonlinear discrete Volterra equations with unbounded delay," *Journal of Computational and Applied Mathematics*, vol. 205, no. 2, pp. 859–870, 2007.
- [31] Y. Song, "Almost periodic solutions of discrete Volterra equations," *Journal of Mathematical Analysis and Applications*, vol. 314, no. 1, pp. 174–194, 2006.
- [32] B. Dai and J. Zou, "Periodic solutions of a discrete-time diffusive system governed by backward difference equations," *Advances in Difference Equations*, vol. 2005, no. 3, pp. 263–274, 2005.
- [33] X. Chen and F. Chen, "Stable periodic solution of a discrete periodic Lotka-Volterra competition system with a feedback control," *Applied Mathematics and Computation*, vol. 181, no. 2, pp. 1446–1454, 2006.
- [34] Y.-H. Fan and W.-T. Li, "Permanence for a delayed discrete ratio-dependent predator-prey system with Holling type functional response," *Journal of Mathematical Analysis and Applications*, vol. 299, no. 2, pp. 357–374, 2004.
- [35] X. Li and W. Yang, "Permanence of a discrete predator-prey systems with Beddington-deAngelis functional response and feedback controls," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 149267, 8 pages, 2008.
- [36] Y. Muroya, "Partial survival and extinction of species in discrete nonautonomous Lotka-Volterra systems," *Tokyo Journal of Mathematics*, vol. 28, no. 1, pp. 189–200, 2005.
- [37] Y. Li and L. Lu, "Positive periodic solutions of discrete n -species food-chain systems," *Applied Mathematics and Computation*, vol. 167, no. 1, pp. 324–344, 2005.
- [38] W. Wendi, G. Mulone, F. Salemi, and V. Salone, "Global stability of discrete population models with time delays and fluctuating environment," *Journal of Mathematical Analysis and Applications*, vol. 264, no. 1, pp. 147–167, 2001.
- [39] R. Y. Zhang, Z. C. Wang, Y. Chen, and J. Wu, "Periodic solutions of a single species discrete population model with periodic harvest/stock," *Computers & Mathematics with Applications*, vol. 39, no. 1-2, pp. 77–90, 2000.
- [40] R. Xu, M. A. J. Chaplain, and F. A. Davidson, "Periodic solutions of a discrete nonautonomous Lotka-Volterra predator-prey model with time delays," *Discrete and Continuous Dynamical Systems*, vol. 4, no. 3, pp. 823–831, 2004.
- [41] Y. Muroya, "Persistence and global stability in discrete models of Lotka-Volterra type," *Journal of Mathematical Analysis and Applications*, vol. 330, no. 1, pp. 24–33, 2007.
- [42] Y. Xia, J. Cao, and M. Lin, "Discrete-time analogues of predator-prey models with monotonic or nonmonotonic functional responses," *Nonlinear Analysis: Real World Applications*, vol. 8, no. 4, pp. 1079–1095, 2007.
- [43] F. Chen, "Permanence in a discrete Lotka-Volterra competition model with deviating arguments," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 5, pp. 2150–2155, 2008.