Research Article

Periodic Solutions for a System of Difference Equations

Shugui Kang^{1,2} and Bao Shi²

¹ College of Mathematics and Computer Science, Shanxi Datong University, Datong, Shanxi 037009, China

² Department of Basic Sciences, Naval Aeronautical Engineering Institute, Yantai, Shandong 264001, China

Correspondence should be addressed to Shugui Kang, dtkangshugui@126.com

Received 9 January 2009; Accepted 8 March 2009

Recommended by Guang Zhang

This paper deals with the second-order nonlinear systems of difference equations, we obtain the existence theorems of periodic solutions. The theorems are proved by using critical point theory.

Copyright © 2009 S. Kang and B. Shi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let $\mathbb{N}, \mathbb{Z}, \mathbb{R}$ be the set of all natural numbers, integers, and real numbers, respectively. For $a, b \in \mathbb{Z}$, note that $\mathbb{Z}[a, b] = \{a, a+1, \dots, b\}$, where $a \leq b$.

In this paper, we consider the existence of periodic solutions for the system of difference equations of the form

$$\begin{aligned} \Delta(p_{n1}(\Delta x_{(n-1)1})^\delta) + q_{n1}(x_{n1})^\delta &= f_1(n, X_n), \\ \Delta(p_{n2}(\Delta x_{(n-1)2})^\delta) + q_{n2}(x_{n2})^\delta &= f_2(n, X_n), \\ &\vdots \\ \Delta(p_{nk}(\Delta x_{(n-1)k})^\delta) + q_{nk}(x_{nk})^\delta &= f_k(n, X_n), \end{aligned} \quad (1.1)$$

which can be recorded as

$$\Delta(\overline{P}_n(\Delta X_{n-1}^T)^\delta) + \overline{Q}_n(X_n^T)^\delta = f(n, X_n), \quad n \in \mathbb{Z}, \quad (1.2)$$

where k is a positive integer,

$$\bar{P}_n = \begin{pmatrix} p_{n1} & 0 & \cdots & 0 \\ 0 & p_{n2} & \cdots & 0 \\ \cdots & & & \\ 0 & 0 & \cdots & p_{nk} \end{pmatrix}, \quad \bar{Q}_n = \begin{pmatrix} q_{n1} & 0 & \cdots & 0 \\ 0 & q_{n2} & \cdots & 0 \\ \cdots & & & \\ 0 & 0 & \cdots & q_{nk} \end{pmatrix}, \quad (1.3)$$

and $\bar{P}_{n+\omega} = \bar{P}_n > 0$ (i.e., $p_{n1} > 0, p_{n2} > 0, \dots, p_{nk} > 0$), $\bar{Q}_{n+\omega} = \bar{Q}_n$, $f = (f_1, f_2, \dots, f_k)^T$, $f_i = f_i(n, X_n) = f_i(n, x_{n1}, x_{n2}, \dots, x_{nk})$, $f(n + \omega, U) = f(n, U)$ for any $(n, U) \in \mathbb{Z} \times \mathbb{R}^k$, $\omega > 0$ is a positive integer, $(-1)^\delta = -1$, δ is the ratio of odd positive integers, $\Delta X_n^T = X_{n+1}^T - X_n^T = (x_{(n+1)1} - x_{n1}, x_{(n+1)2} - x_{n2}, \dots, x_{(n+1)k} - x_{nk})^T$, $\Delta^2 X_{n-1}^T = \Delta(\Delta X_{n-1}^T) = \Delta X_n^T - \Delta X_{n-1}^T$. For $U = (u_1, u_2, \dots, u_k) \in \mathbb{R}^k$, define $U^\delta = (u_1^\delta, u_2^\delta, \dots, u_k^\delta)$. $|U| = (|u_1|, |u_2|, \dots, |u_k|)$, $|U|^\delta = (|u_1|^\delta, |u_2|^\delta, \dots, |u_k|^\delta)$. A sequence $X = \{X_n\}_{n \in \mathbb{Z}}$ is a ω -periodic solution of (1.2) if substitution of it into (1.2) yields an identity for all $n \in \mathbb{Z}$.

In [1, 2], the qualitative behavior of linear difference equations

$$\Delta(p_n \Delta x_n) + q_n x_n = 0 \quad (1.4)$$

has been investigated. In [3], the nonlinear difference equation

$$\Delta(p_n \Delta x_{n-1}) + q_n x_n = f(n, x_n) \quad (1.5)$$

has been considered. In [4], by critical point method, the existence of periodic and subharmonic solutions of equation

$$\Delta^2 x_{n-1} + f(n, x_n) = 0, \quad n \in \mathbb{Z} \quad (1.6)$$

has been studied. Other interesting results can be found in [5–8]. In [9], the authors consider the existence of periodic solutions for second-order nonlinear difference equation

$$\Delta(p_n (\Delta x_{n-1})^\delta) + q_n x_n^\delta = f(n, x_n), \quad n \in \mathbb{Z}, \quad (1.7)$$

using critical point theory, obtaining some new results. It is a discrete analogues of differential equation

$$(p(t)\phi(u'))' + f(t, u) = 0. \quad (1.8)$$

They do have physical applications in the study of nuclear physics, gas aerodynamics, and so on (see [10, 11]). In this paper, we obtain some new results of existence of periodic solution for the second-order nonlinear system of difference equations by using critical point theory. We remark, however, the result in [9] is only good for (1.7) which is much less general than our results in what follows.

2. Some Basic Lemmas

Let E be a real Hilbert space, $I \in C^1(E, \mathbb{R})$ mean that I is continuously Fréchet differentiable functional defined on E . I is said to be satisfying Palais-Smale condition (P-S condition) if any bounded sequence $\{I(u_n)\}$ and $I'(u_n) \rightarrow 0$ ($n \rightarrow \infty$) possess a convergent subsequence in E . Let B_ρ be the open ball in E with radius ρ and centered at θ , and let ∂B_ρ denote its boundary, θ is null element of E .

Lemma 2.1 (see [12]). *Let E be a real Hilbert space, and assume that $I \in C^1(E, \mathbb{R})$ satisfies the P-S condition and the following conditions:*

- (I₁) *there exist constants $\rho > 0$ and $a > 0$ such that $I(x) \geq a$ for all $x \in \partial B_\rho$, where $B_\rho = \{x \in E : \|x\| < \rho\}$;*
- (I₂) *$I(0) \leq 0$ and there exists $x_0 \notin B_\rho$ such that $I(x_0) \leq 0$.*

Then $c = \inf_{h \in \Gamma} \sup_{s \in [0,1]} I(h(s))$ is a positive critical value of I , where

$$\Gamma = \{h \in C([0,1], X) : h(0) = \theta, h(1) = x_0\}. \quad (2.1)$$

Let Ω_* be the set of sequences

$$X = \{X_n\}_{n \in \mathbb{Z}} = \{\dots, X_{-n}, \dots, X_{-1}, X_0, X_1, \dots, X_n, \dots\}, \quad (2.2)$$

where $X_n = (x_{n1}, x_{n2}, \dots, x_{nk}) \in \mathbb{R}^k$, that is,

$$\Omega_* = \{X = \{X_n\}_{n \in \mathbb{Z}} : X_n \in \mathbb{R}^k, n \in \mathbb{Z}\}. \quad (2.3)$$

For any $X, Y \in \Omega_*$, $a, b \in \mathbb{R}$, $aX + bY$ is defined by

$$aX + bY = \{aX_n + bY_n\}_{n=-\infty}^{+\infty}, \quad (2.4)$$

then Ω_* is a vector space. For given positive integer ω , E_ω is defined as a subspace of Ω_* by

$$E_\omega = \{X = \{X_n\} \in \Omega_* : X_{n+\omega} = X_n, n \in \mathbb{Z}\}. \quad (2.5)$$

Obviously, E_ω is isomorphic to $\mathbb{R}^{k\omega}$, for any $X, Y \in E_\omega$, defined inner product

$$\langle X, Y \rangle = \sum_{i=1}^{\omega} \langle X_i, Y_i \rangle, \quad (2.6)$$

by which the norm $\|\cdot\|$ can be induced by

$$\|X\| = \left(\sum_{i=1}^{\omega} \|X_i\|^2 \right)^{1/2}, \quad X \in E_\omega. \quad (2.7)$$

where $\|X_i\| = (\sum_{j=1}^k |x_{ij}|^2)^{1/2}$. It is obvious that E_ω with the inner product defined by (2.6) is a finite-dimensional Hilbert space and linearly homeomorphic to $\mathbb{R}^{k\omega}$. Define the functional J on E_ω as follows:

$$J(X) = \frac{1}{\delta+1} \sum_{n=1}^{\omega} \langle P_n, (\Delta X_{n-1})^{\delta+1} \rangle - \frac{1}{\delta+1} \sum_{n=1}^{\omega} \langle Q_n, X_n^{\delta+1} \rangle + \sum_{n=1}^{\omega} F(n, X_n), \quad X \in E_\omega, \quad (2.8)$$

where $F(n, X_n)$ such that $\nabla_U F(n, U) = f(n, U)$, that is,

$$f_i(n, U) = f_i(n, u_1, u_2, \dots, u_k) = \frac{\partial}{\partial u_i} F(n, u_1, u_2, \dots, u_k) \quad (2.9)$$

for any $(n, U) \in \mathbb{Z}[1, \omega] \times \mathbb{R}^k$, $P_n = (p_{n1}, p_{n2}, \dots, p_{nk})$, $Q_n = (q_{n1}, q_{n2}, \dots, q_{nk})$. Clearly $J \in C^1(E_\omega, \mathbb{R})$, and for any $X = \{X_n\}_{n \in \mathbb{Z}} \in E_\omega$, by $X_0 = X_\omega$ and $X_1 = X_{\omega+1}$, we have

$$\frac{\partial J(X)}{\partial x_{nl}} = -\Delta(p_{nl}(\Delta x_{(n-1)l})^\delta) - q_{nl}(x_{nl})^\delta + f_l(n, X_n), \quad l \in \mathbb{Z}[1, k], \quad n \in \mathbb{Z}[1, \omega]. \quad (2.10)$$

Thus $X = \{X_n\}_{n \in \mathbb{Z}}$ is a critical point of J on E_ω ($J'(X) = 0$) if and only if

$$\Delta(p_{nl}(\Delta x_{(n-1)l})^\delta) + q_{nl}(x_{nl})^\delta = f_l(n, X_n), \quad l \in \mathbb{Z}[1, k], \quad n \in \mathbb{Z}[1, \omega]. \quad (2.11)$$

That is,

$$\Delta(\bar{P}_n(\Delta X_{n-1}^T)^\delta) + \bar{Q}_n(X_n^T)^\delta = f(n, X_n), \quad n \in \mathbb{Z}. \quad (2.12)$$

By the periodicity of X_n and $f(n, X_n)$ in the first variable n , we know that if $X = \{X_n\}_{n \in \mathbb{Z}} \in E_\omega$ is a critical point of the real functional J defined by (2.8), then it is a periodic solution of (1.2).

For $X = \{X_n\}_{n \in \mathbb{Z}} \in E_\omega$, $X_n = (x_{n1}, x_{n2}, \dots, x_{nk}) \in \mathbb{R}^k$, $r > 1$, denote

$$\|X\|_r = \left(\sum_{i=1}^{\omega} \|X_i\|^r \right)^{1/r}, \quad \|X_n\|_r = \left(\sum_{i=1}^k \|x_{ni}\|^r \right)^{1/r}. \quad (2.13)$$

Clearly, $\|X\|_2 = \|X\|$, $\|X_n\|_2 = \|X_n\|$. Because of $\|\cdot\|_{r_1}$ and $\|\cdot\|_{r_2}$ being equivalent when $r_1, r_2 > 1$, so there exist constants $c_1, c_2, c_3, c_4, \hbar_1, \hbar_2, \hbar_3$, and $\hbar_4$ such that $c_2 \geq c_1 > 0$, $c_4 \geq c_3 > 0$, $\hbar_2 \geq \hbar_1 > 0$, and $\hbar_4 \geq \hbar_3 > 0$,

$$\begin{aligned} c_1 \|X\| &\leq \|X\|_{\delta+1} \leq c_2 \|X\|, \\ c_3 \|X\| &\leq \|X\|_\beta \leq c_4 \|X\|, \\ \hbar_1 \|X_n\| &\leq \|X_n\|_{\delta+1} \leq \hbar_2 \|X_n\|, \\ \hbar_3 \|X_n\| &\leq \|X_n\|_\beta \leq \hbar_4 \|X_n\|, \end{aligned} \quad (2.14)$$

for all $X \in E_\omega$, $\delta > 0$, and $\beta > 1$.

Lemma 2.2. Suppose that

(F₁) there exist constants $a_1 > 0$, $a_2 > 0$, $\beta > \delta + 1$ such that

$$F(n, U) \leq -a_1 \|U\|^\beta + a_2 \quad (2.15)$$

for any $(n, U) \in \mathbb{Z}[1, \omega] \times R^k$;

(F₂)

$$q_{ni} \leq 0, \quad n \in \mathbb{Z}, i \in \mathbb{Z}[1, k]. \quad (2.16)$$

Then

$$J(X) = \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \langle P_n, (\Delta X_{n-1})^{\delta+1} \rangle - \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \langle Q_n, X_n^{\delta+1} \rangle + \sum_{n=1}^{\omega} F(n, X_n) \quad (2.17)$$

satisfies P-S condition.

Proof. For any sequence $\{X^{(l)}\} = \{\dots, X_{-n}^{(l)}, \dots, X_{-1}^{(l)}, X_0^{(l)}, X_1^{(l)}, \dots, X_n^{(l)}, \dots\} \in E_\omega$, $J(X^{(l)})$ is bounded and $J'(X^{(l)}) \rightarrow 0$ ($l \rightarrow \infty$). Then there exists a positive constant $M > 0$, such that $|J(X^{(l)})| \leq M$. From (F₁), we have

$$\begin{aligned} -M &\leq J(X^{(l)}) \\ &= \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \left[\langle P_n, (X_n^{(l)} - X_{n-1}^{(l)})^{\delta+1} \rangle - \langle Q_n, (X_n^{(l)})^{\delta+1} \rangle \right] \\ &\quad + \sum_{n=1}^{\omega} F(n, X_n^{(l)}) \\ &\leq \frac{1}{\delta + 1} \sum_{n=1}^{\omega} 2^{\delta+1} \langle P_n, (|X_n^{(l)}|^{\delta+1} + |X_{n-1}^{(l)}|^{\delta+1}) \rangle \\ &\quad - \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \langle Q_n, |X_n^{(l)}|^{\delta+1} \rangle + \sum_{n=1}^{\omega} F(n, X_n^{(l)}) \\ &\leq \frac{2^{\delta+1}}{\delta + 1} \sum_{n=1}^{\omega} \langle P_n + P_{n+1}, |X_n^{(l)}|^{\delta+1} \rangle - \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \langle Q_n, |X_n^{(l)}|^{\delta+1} \rangle \\ &\quad + \sum_{n=1}^{\omega} F(n, X_n^{(l)}) \\ &\leq \frac{2^{\delta+1}}{\delta + 1} \sum_{n=1}^{\omega} \langle P_n + P_{n+1}, |X_n^{(l)}|^{\delta+1} \rangle - \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \langle Q_n, |X_n^{(l)}|^{\delta+1} \rangle \\ &\quad - a_1 \sum_{n=1}^{\omega} \|X_n^{(l)}\|^\beta + a_2 \omega \\ &= \frac{1}{\delta + 1} \sum_{n=1}^{\omega} \langle 2^{\delta+1} (P_n + P_{n+1}) - Q_n, |X_n^{(l)}|^{\delta+1} \rangle - a_1 \|X^{(l)}\|_\beta^\beta + a_2 \omega. \end{aligned} \quad (2.18)$$

Set

$$A_0 = \max_{n \in \mathbb{Z}[1, \omega], i \in \mathbb{Z}[1, k]} \{2^{\delta+1}(p_{ni} + p_{(n+1)i}) - q_{ni}\}. \quad (2.19)$$

Then $A_0 > 0$, and

$$\begin{aligned} -M &\leq J(X^{(l)}) \\ &\leq \frac{A_0}{\delta+1} \sum_{n=1}^{\omega} \|X_n^{(l)}\|_{\delta+1}^{\delta+1} - a_1 \|X^{(l)}\|_{\beta}^{\beta} + a_2 \omega \\ &\leq \frac{A_0 \hbar_2^{\delta+1}}{\delta+1} \sum_{n=1}^{\omega} \|X_n^{(l)}\|^{\delta+1} - a_1 \|X^{(l)}\|_{\beta}^{\beta} + a_2 \omega \\ &= \frac{A_0 \hbar_2^{\delta+1}}{\delta+1} \|X^{(l)}\|_{\delta+1}^{\delta+1} - a_1 \|X^{(l)}\|_{\beta}^{\beta} + a_2 \omega. \end{aligned} \quad (2.20)$$

Because of $\beta > \delta + 1$, and $(\beta - \delta - 1)/\beta + (\delta + 1)/\beta = 1$, in view of Hölder inequality, we have

$$\sum_{n=1}^{\omega} \|X_n^{(l)}\|^{\delta+1} \leq \omega^{(\beta-\delta-1)/\beta} \left(\sum_{n=1}^{\omega} \|X_n^{(l)}\|_{\beta}^{\beta} \right)^{(\delta+1)/\beta}. \quad (2.21)$$

Thus

$$\|X^{(l)}\|_{\beta}^{\beta} \geq \omega^{(\delta+1-\beta)/(\delta+1)} \|X^{(l)}\|_{\delta+1}^{\beta}. \quad (2.22)$$

Then we have

$$\begin{aligned} -M &\leq J(X^{(l)}) \\ &\leq \frac{A_0 \hbar_2^{\delta+1}}{\delta+1} \|X^{(l)}\|_{\delta+1}^{\delta+1} - a_1 \|X^{(l)}\|_{\beta}^{\beta} + a_2 \omega \\ &\leq \frac{A_0 \hbar_2^{\delta+1}}{\delta+1} \|X^{(l)}\|_{\delta+1}^{\delta+1} - a_1 \omega^{(\delta+1-\beta)/(\delta+1)} \|X^{(l)}\|_{\delta+1}^{\beta} + a_2 \omega. \end{aligned} \quad (2.23)$$

Thus, for any $l \in \mathbb{N}$,

$$a_1 \omega^{(\delta+1-\beta)/(\delta+1)} \|X^{(l)}\|_{\delta+1}^{\beta} - \frac{A_0 \hbar_2^{\delta+1}}{\delta+1} \|X^{(l)}\|_{\delta+1}^{\delta+1} \leq M + a_2 \omega. \quad (2.24)$$

Because of $\beta > \delta + 1$, it is easily seen that the inequality (2.24) implies that $\{X^{(l)}\}$ is a bounded sequence in E_{ω} . Thus $\{X^{(l)}\}$ possesses convergent subsequences. The proof is complete. $\square$

3. Main Result

Theorem 3.1. *Suppose that condition (F₁) holds, and*

(F₃) for each $n \in \mathbb{Z}$,

$$\lim_{\|U\| \rightarrow 0} \frac{F(n, U)}{\|U\|^{\delta+1}} = 0; \quad (3.1)$$

(F₄) for any $i \in \mathbb{Z}[1, k], n \in \mathbb{Z}[1, \omega]$,

$$q_{ni} < 0; \quad (3.2)$$

(F₅) $F(n, \theta) = 0$.

Then (1.2) has at least two nontrivial ω -periodic solutions.

Proof. By Lemma 2.2, J satisfies P-S condition. Next, we will verify the conditions (I₁) and (I₂) of Lemma 2.1. By (F₃), there exists $\rho > 0$, such that

$$|F(n, U)| \leq -\frac{q_{\max} h_1^{\delta+1}}{2(\delta+1)} \|U\|^{\delta+1} \quad (3.3)$$

for any $\|U\| < \rho$ and $n \in \mathbb{Z}[1, \omega]$, where $q_{\max} = \max_{n \in \mathbb{Z}[1, \omega], i \in \mathbb{Z}[1, k]} q_{ni} < 0$. Thus

$$\begin{aligned} J(X) &\geq -\frac{1}{\delta+1} \sum_{n=1}^{\omega} \langle Q_n, X_n^{\delta+1} \rangle + \sum_{n=1}^{\omega} F(n, X_n) \\ &\geq -\frac{q_{\max}}{\delta+1} \sum_{n=1}^{\omega} \|X_n\|_{\delta+1}^{\delta+1} + \frac{q_{\max} h_1^{\delta+1}}{2(\delta+1)} \sum_{n=1}^{\omega} \|X_n\|^{\delta+1} \\ &\geq -\frac{q_{\max} h_1^{\delta+1}}{\delta+1} \sum_{n=1}^{\omega} \|X_n\|^{\delta+1} + \frac{q_{\max} h_1^{\delta+1}}{2(\delta+1)} \sum_{n=1}^{\omega} \|X_n\|^{\delta+1} \\ &= -\frac{q_{\max} h_1^{\delta+1}}{2(\delta+1)} \|X\|_{\delta+1}^{\delta+1} \\ &\geq -\frac{q_{\max} h_1^{\delta+1} c_1^{\delta+1}}{2(\delta+1)} \|X\|^{\delta+1} \end{aligned} \quad (3.4)$$

for any $X \in E_{\omega}$ with $\|X\| \leq \rho$. We choose $a = -h_1^{\delta+1} c_1^{\delta+1} (q_{\max}/2(\delta+1)) \rho^{\delta+1}$, then we have

$$J(X)|_{\partial B_{\rho}} \geq a > 0, \quad (3.5)$$

that is, the condition (I₁) of Lemma 2.1 holds.

Obviously, $J(0) = 0$. For any given $V \in E_\omega$ with $\|V\| = 1$ and constant $\alpha > 0$,

$$\begin{aligned}
J(\alpha V) &= \frac{1}{\delta+1} \sum_{n=1}^{\omega} \langle P_n, (\alpha V_n - \alpha V_{n-1})^{\delta+1} \rangle - \frac{1}{\delta+1} \sum_{n=1}^{\omega} \langle Q_n, (\alpha V_n)^{\delta+1} \rangle + \sum_{n=1}^{\omega} F(n, \alpha V_n) \\
&= \frac{1}{\delta+1} \sum_{n=1}^{\omega} \left\{ p_{n1} (\alpha v_{n1} - \alpha v_{(n-1)1})^{\delta+1} + p_{n2} (\alpha v_{n2} - \alpha v_{(n-1)2})^{\delta+1} \right. \\
&\quad \left. + \cdots + p_{nk} (\alpha v_{nk} - \alpha v_{(n-1)k})^{\delta+1} \right\} \\
&\quad - \frac{1}{\delta+1} \sum_{n=1}^{\omega} \{ q_{n1} (\alpha v_{n1})^{\delta+1} + q_{n2} (\alpha v_{n2})^{\delta+1} + \cdots + q_{nk} (\alpha v_{nk})^{\delta+1} \} \\
&\quad + \sum_{n=1}^{\omega} F(n, \alpha V_n) \\
&\leq \frac{1}{\delta+1} \sum_{n=1}^{\omega} \{ p_{n1} (2\alpha)^{\delta+1} + p_{n2} (2\alpha)^{\delta+1} + \cdots + p_{nk} (2\alpha)^{\delta+1} \} \\
&\quad - \frac{1}{\delta+1} \sum_{n=1}^{\omega} \{ q_{n1} \alpha^{\delta+1} + q_{n2} \alpha^{\delta+1} + \cdots + q_{nk} \alpha^{\delta+1} \} \\
&\quad - a_1 \sum_{n=1}^{\omega} \|\alpha V_n\|^\beta + a_2 \omega \\
&\leq \frac{1}{\delta+1} \sum_{n=1}^{\omega} \{ 2^{\delta+1} \|P_n\|_1 + \|Q_n\|_1 \} \alpha^{\delta+1} - a_1 \alpha^\beta \sum_{n=1}^{\omega} \|V_n\|^\beta + a_2 \omega \\
&\leq \frac{1}{\delta+1} \sum_{n=1}^{\omega} \{ 2^{\delta+1} \|P_n\|_1 + \|Q_n\|_1 \} \alpha^{\delta+1} - a_1 \alpha^\beta \|V\|_\beta^\beta + a_2 \omega \\
&\leq \frac{1}{\delta+1} \sum_{n=1}^{\omega} \{ 2^{\delta+1} \|P_n\|_1 + \|Q_n\|_1 \} \alpha^{\delta+1} - a_1 c_3^\beta \alpha^\beta + a_2 \omega \\
&\longrightarrow -\infty, \quad (\alpha \longrightarrow +\infty).
\end{aligned} \tag{3.6}$$

Thus we can choose a sufficiently large α such that $\alpha > \rho$, and $\bar{X} = \alpha V \in E_\omega$, $J(\bar{X}) < 0$. According to Lemma 2.1, there exists at least one critical value $c \geq a > 0$. We suppose that X^* is a critical point corresponding to c , then $J(X^*) = c$ and $J'(X^*) = 0$.

By similar argument of Lemma 2.2, we know that $J(X)$ is bounded from above, so there exists $X^{**} \in E_\omega$ such that $J(X) \leq J(X^{**}) = c_{\max}$ for any $X \in E_\omega$. Obviously, $X^{**} \neq 0$. If $X^{**} \neq X^*$, then the proof is complete. Otherwise, $X^{**} = X^*$, $c = c_{\max}$. In view of Lemma 2.1,

$$c = \inf_{h \in \Gamma} \sup_{s \in [0,1]} J(h(s)), \tag{3.7}$$

where $\Gamma = \{h \in C([0,1], E_\omega) : h(0) = \theta, h(1) = \bar{X}\}$. Then $c_{\max} = \max_{s \in [0,1]} J(h(s))$ for any $h \in \Gamma$ holds. In view of the continuity of $J(h(s))$ in s , $J(\theta) \leq 0$, and $J(\bar{X}) < 0$, we know that there

exists some $s_0 \in (0, 1)$ such that $J(h(s_0)) = c_{\max}$. If we choose $h_1, h_2 \in \Gamma$ such that

$$\{h_1(s) : s \in (0, 1)\} \cap \{h_2(s) : s \in (0, 1)\} = \emptyset, \quad (3.8)$$

then there exist $s_1, s_2 \in (0, 1)$ such that $J(h_1(s_1)) = J(h_2(s_2)) = c_{\max}$. Then J possesses two different critical points $\tilde{Y} = h_1(s_1)$ and $\tilde{Z} = h_2(s_2)$ in E_ω , hence, we obtain at least two nontrivial critical points which correspond to the critical value $c_{\max}$. Thus (1.2) possesses at least two nontrivial ω -periodic solutions. The proof is complete. $\square$

Acknowledgments

This work is supported by Natural Science Foundation of Shanxi Province (2008011002-1) and Shanxi Datong University and by the Development Foundation of Higher Education Department of Shanxi Province.

References

- [1] C. D. Ahlbrandt and A. C. Peterson, *Discrete Hamiltonian Systems: Difference Equations, Continued Fractions, and Riccati Equations*, vol. 16 of *Kluwer Texts in the Mathematical Sciences*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1996.
- [2] S.-S. Cheng, H.-J. Li, and W. T. Patula, "Bounded and zero convergent solutions of second-order difference equations," *Journal of Mathematical Analysis and Applications*, vol. 141, no. 2, pp. 463–483, 1989.
- [3] T. Peil and A. Peterson, "Criteria for C-disfocality of a selfadjoint vector difference equation," *Journal of Mathematical Analysis and Applications*, vol. 179, no. 2, pp. 512–524, 1993.
- [4] Z. Guo and J. Yu, "Existence of periodic and subharmonic solutions for second-order superlinear difference equations," *Science in China Series A*, vol. 46, no. 4, pp. 506–515, 2003.
- [5] R. P. Agarwal and P. J. Y. Wong, *Advanced Topics in Difference Equations*, vol. 404 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [6] M. Cecchi, Z. Dořlá, and M. Marini, "Positive decreasing solutions of quasi-linear difference equations," *Computers & Mathematics with Applications*, vol. 42, no. 10-11, pp. 1401–1410, 2001.
- [7] P. J. Y. Wong and R. P. Agarwal, "Oscillations and nonoscillations of half-linear difference equations generated by deviating arguments," *Computers & Mathematics with Applications*, vol. 36, no. 10-12, pp. 11–26, 1998.
- [8] P. J. Y. Wong and R. P. Agarwal, "Oscillation and monotone solutions of second order quasilinear difference equations," *Funkcialaj Ekvacioj*, vol. 39, no. 3, pp. 491–517, 1996.
- [9] X. Cai and J. Yu, "Existence theorems of periodic solutions for second-order nonlinear difference equations," *Advances in Difference Equations*, vol. 2008, Article ID 247071, 11 pages, 2008.
- [10] R. P. Agarwal and P. Y. H. Pang, "On a generalized difference system," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 30, no. 1, pp. 365–376, 1997.
- [11] M. Marini, "On nonoscillatory solutions of a second-order nonlinear differential equation," *Bollettino della Unione Matematica Italiana*, vol. 3, no. 1, pp. 189–202, 1984.
- [12] J. Mawhin and M. Willem, *Critical Point Theory and Hamiltonian Systems*, vol. 74 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1989.