Research Article

Complete Convergence for Weighted Sums of ρ^* -Mixing Random Variables

Soo Hak Sung

Department of Applied Mathematics, Pai Chai University, Taejeon 302-735, South Korea

Correspondence should be addressed to Soo Hak Sung, sungsh@pcu.ac.kr

Received 2 August 2009; Accepted 24 March 2010

Academic Editor: Leonid Shaikhet

Copyright © 2010 Soo Hak Sung. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We obtain the complete convergence for weighted sums of ρ^* -mixing random variables. Our result extends the result of Peligrad and Gut (1999) on unweighted average to a weighted average under a mild condition of weights. Our result also generalizes and sharpens the result of An and Yuan (2008).

1. Introduction

In many stochastic models, the assumption that random variables are independent is not plausible. So it is of interest to extend the concept of independence to dependence cases. One of these dependence structures is ρ^* -mixing.

Let $\{X_n, n \geq 1\}$ be a sequence of random variables defined on a probability space $(\Omega, \mathcal{F}, P)$, and let $\mathcal{F}_n^m$ denote the σ -algebra generated by the random variables $X_n, X_{n+1}, \dots, X_m$. For any $S \subset N$, define $\mathcal{F}_S = \sigma(X_i, i \in S)$. Given two σ -algebras $\mathcal{A}, \mathcal{B}$ in $\mathcal{F}$, put

$$\rho(\mathcal{A}, \mathcal{B}) = \sup\{\text{corr}(X, Y); X \in L_2(\mathcal{A}), Y \in L_2(\mathcal{B})\}, \quad (1.1)$$

where $\text{corr}(X, Y) = (EXY - EXEY) / \sqrt{\text{var}(X)\text{var}(Y)}$. Define the ρ^* -mixing coefficients by

$$\rho_n^* = \sup\{\rho(\mathcal{F}_S, \mathcal{F}_T); S, T \subset N \text{ with } \text{dist}(S, T) \geq n\}. \quad (1.2)$$

Obviously, $0 \leq \rho_{n+1}^* \leq \rho_n^* \leq \rho_0^* = 1$. The sequence $\{X_n, n \geq 1\}$ is called ρ^* -mixing (or $\tilde{\rho}$ -mixing) if there exists $k \in \mathbb{N}$ such that $\rho_k^* < 1$. Note that if $\{X_n, n \geq 1\}$ is a sequence of independent random variables, then $\rho_n^* = 0$ for all $n \geq 1$.

A number of limit results for ρ^* -mixing sequences of random variables have been established by many authors. We refer to Bradley [1] for the central limit theorem, Bryc and Smoleński [2], Peligrad and Gut [3], and Utev and Peligrad [4] for moment inequalities, Gan [5], Kucmaszewska [6], and Wu and Jiang [7] for almost sure convergence, and An and Yuan [8], Cai [9], Gan [5], Kucmaszewska [10], Peligrad and Gut [3], and Zhu [11] for complete convergence.

The concept of complete convergence of a sequence of random variables was introduced by Hsu and Robbins [12]. A sequence $\{X_n, n \geq 1\}$ of random variables converges completely to the constant θ if

$$\sum_{n=1}^{\infty} P(|X_n - \theta| > \epsilon) < \infty \quad \forall \epsilon > 0. \quad (1.3)$$

In view of the Borel-Cantelli lemma, this implies that $X_n \rightarrow \theta$ almost surely. Therefore, the complete convergence is a very important tool in establishing almost sure convergence of summation of random variables as well as weighted sums of random variables. Hsu and Robbins [12] proved that the sequence of arithmetic means of independent and identically distributed random variables converges completely to the expected value if the variance of the summands is finite. Erdős [13] proved the converse. The result of Hsu-Robbins-Erdős is a fundamental theorem in probability theory and has been generalized and extended in several directions by many authors. One of the most important generalizations is Baum and Katz [14] strong law of large numbers.

Theorem 1.1 (Baum and Katz [14]). *Let $p \geq 1/\alpha$ and $1/2 < \alpha \leq 1$. Let $\{X_n, n \geq 1\}$ be a sequence of independent and identically distributed random variables with $EX_1 = 0$. Then the following statements are equivalent:*

- (i) $E|X_1|^p < \infty$;
- (ii) $\sum_{n=1}^{\infty} n^{p\alpha-2} P(\max_{1 \leq j \leq n} |\sum_{i=1}^j X_i| > \epsilon n^\alpha) < \infty$ for all $\epsilon > 0$,

Peligrad and Gut [3] extended the result of Baum and Katz [14] to ρ^ -mixing random variables.*

Theorem 1.2 (Peligrad and Gut [3]). *Let $p > 1/\alpha$ and $1/2 < \alpha \leq 1$. Let $\{X_n, n \geq 1\}$ be a sequence of identically distributed ρ^* -mixing random variables with $EX_1 = 0$. Then the following statements are equivalent:*

- (i) $E|X_1|^p < \infty$;
- (ii) $\sum_{n=1}^{\infty} n^{p\alpha-2} P(\max_{1 \leq j \leq n} |\sum_{i=1}^j X_i| > \epsilon n^\alpha) < \infty$ for all $\epsilon > 0$.

Cai [9] complemented Theorem 1.2 when $p = 1/\alpha$.

Recently, An and Yuan [8] obtained a complete convergence result for weighted sums of identically distributed ρ^ -mixing random variables.*

Theorem 1.3 (An and Yuan [8]). *Let $p > 1/\alpha$ and $1/2 < \alpha \leq 1$. Let $\{X_n, n \geq 1\}$ be a sequence of identically distributed ρ^* -mixing random variables with $EX_1 = 0$. Assume that $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$ is an array of real numbers satisfying*

$$\sum_{i=1}^n |a_{ni}|^p = O(n^\delta) \quad \text{for some } 0 < \delta < 1, \quad (1.4)$$

$$\#A_{nk} = \#\left\{1 \leq i \leq n : |a_{ni}|^p > (k+1)^{-1}\right\} \geq ne^{-1/k} \quad \forall k \geq 1, n \geq 1. \quad (1.5)$$

Then the following statements are equivalent:

(i) $E|X_1|^p < \infty$;

(ii) $\sum_{n=1}^{\infty} n^{p\alpha-2} P(\max_{1 \leq j \leq n} |\sum_{i=1}^j a_{ni} X_i| > \epsilon n^\alpha) < \infty$ for all $\epsilon > 0$.

Note that the result of An and Yuan [8] is not an extension of Peligrad and Gut's [3] result, since condition (1.4) does not hold for the array with $a_{ni} = 1, 1 \leq i \leq n, n \geq 1$. An and Yuan [8] proved the implication (i) $\Rightarrow$ (ii) under condition (1.4), and proved the converse under conditions (1.4) and (1.5). However, the array satisfying both (1.4) and (1.5) does not exist. Noting that $\#A_{nk}/(k+1) \leq \sum_{i=1}^n |a_{ni}|^p \leq O(n^\delta)$, we have that $ne^{-1/k} \leq \#A_{nk} \leq (k+1)O(n^\delta)$. But, this does not hold when k is fixed and n is large enough.

In this paper, we obtain a new complete convergence result for weighted sums of identically distributed ρ^* -mixing random variables. Our result extends the result of Peligrad and Gut [3], and generalizes and sharpens the result of An and Yuan [8].

Throughout this paper, the symbol C denotes a positive constant which is not necessarily the same one in each appearance, $[x]$ denotes the integer part of x , and $a \wedge b = \min\{a, b\}$.

2. Main Result

To prove our main result, we need the following lemma which is a Rosenthal-type inequality for ρ^* -mixing random variables.

Lemma 2.1 (Utev and Peligrad [4]). *Let $\{X_n, n \geq 1\}$ be a sequence of ρ^* -mixing random variables with $EX_n = 0$ and $E|X_n|^r < \infty$ for some $r \geq 2$ and all $n \geq 1$. Then there exists a constant $D = D(r, k, \rho_k^*)$ depending only on r, k , and ρ_k^* such that for any $n \geq 1$,*

$$E\left(\max_{1 \leq j \leq n} \left|\sum_{i=1}^j X_i\right|^r\right) \leq D \left\{ \sum_{i=1}^n E|X_i|^r + \left(\sum_{i=1}^n EX_i^2\right)^{r/2} \right\}, \quad (2.1)$$

where $\rho_k^* < 1$.

Now we state the main result of this paper.

Theorem 2.2. Let $p > 1/\alpha$ and $1/2 < \alpha \leq 1$. Let $\{X_n, n \geq 1\}$ be a sequence of identically distributed ρ^* -mixing random variables with $EX_1 = 0$. Assume that $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$ is an array of real numbers satisfying

$$\sum_{i=1}^n |a_{ni}|^q = O(n) \quad \text{for some } q > p. \quad (2.2)$$

If $E|X_1|^p < \infty$, then

$$\sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| > \epsilon n^\alpha\right) < \infty \quad \forall \epsilon > 0. \quad (2.3)$$

Conversely, if (2.3) holds for any array $\{a_{ni}\}$ satisfying (2.2), then $E|X_1|^p < \infty$.

To prove Theorem 2.2, we first prove the following lemma which is the sufficiency of Theorem 2.2 when the array is bounded.

Lemma 2.3. Let $\{X_n, n \geq 1\}$ be a sequence of identically distributed ρ^* -mixing random variables with $EX_1 = 0$ and $E|X_1|^p < \infty$ for some $p > 1/\alpha$ and $1/2 < \alpha \leq 1$. Assume that $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$ is an array of real numbers satisfying $|a_{ni}| \leq 1$ for $1 \leq i \leq n$ and $n \geq 1$. Then (2.3) holds.

Proof. For $1 \leq i \leq n$ and $n \geq 1$, define $X'_{ni} = X_i I(|X_i| \leq n^\alpha)$. Since $EX_i = 0$ and $\sum_{i=1}^n |a_{ni}| \leq n$, we have that

$$\begin{aligned} n^{-\alpha} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} EX'_{ni} \right| &= n^{-\alpha} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} EX_i I(|X_i| > n^\alpha) \right| \\ &\leq n^{-\alpha} \sum_{i=1}^n |a_{ni}| E|X_i| I(|X_i| > n^\alpha) \\ &\leq n^{1-\alpha} E|X_1| I(|X_1| > n^\alpha) \\ &\leq n^{1-p\alpha} E|X_1|^p I(|X_1| > n^\alpha) \longrightarrow 0 \end{aligned} \quad (2.4)$$

as $n \rightarrow \infty$. Hence for n large enough, we have

$$n^{-\alpha} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} EX'_{ni} \right| < \frac{\epsilon}{2}. \quad (2.5)$$

It follows that

$$\begin{aligned}
& \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| > \epsilon n^{\alpha}\right) \\
& \leq \sum_{n=1}^{\infty} n^{p\alpha-2} \sum_{i=1}^n P(|X_i| > n^{\alpha}) + \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X'_{ni} \right| > \epsilon n^{\alpha}\right) \\
& \leq \sum_{n=1}^{\infty} n^{p\alpha-1} P(|X_1| > n^{\alpha}) + C \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} (X'_{ni} - EX'_{ni}) \right| > \frac{\epsilon n^{\alpha}}{2}\right) \\
& =: I + CJ.
\end{aligned} \tag{2.6}$$

Noting that $\sum_{n=1}^{\infty} n^{p\alpha-1} P(|X_1| > n^{\alpha}) \leq CE|X_1|^p < \infty$, we have $I < \infty$. Thus, it remains to show that $J < \infty$.

We have by Markov's inequality and Lemma 2.1 that for any $r \geq 2$,

$$\begin{aligned}
J & \leq \left(\frac{2}{\epsilon}\right)^r \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} E \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} (X'_{ni} - EX'_{ni}) \right|^r \\
& \leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} \left\{ \left(\sum_{i=1}^n a_{ni}^2 E|X'_{ni}|^2 \right)^{r/2} + \sum_{i=1}^n |a_{ni}|^r E|X'_{ni}|^r \right\} \\
& \leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2+r/2} \left(E|X_1|^2 I(|X_1| \leq n^{\alpha}) \right)^{r/2} + C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-1} E|X_1|^r I(|X_1| \leq n^{\alpha}) \\
& =: CJ_1 + CJ_2.
\end{aligned} \tag{2.7}$$

In the last inequality, we used the fact that $|a_{ni}| \leq 1$ for $1 \leq i \leq n$ and $n \geq 1$.

If $p \geq 2$, then we take large enough r such that $r > \max\{(p\alpha - 1)/(\alpha - 1/2), p\}$. Since $r > (p\alpha - 1)/(\alpha - 1/2)$, we get

$$J_1 \leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2+r/2} < \infty. \tag{2.8}$$

Since $r > p$, we also get

$$\begin{aligned}
 J_2 &= \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-1} \sum_{i=1}^n E|X_1|^r I((i-1)^\alpha < |X_1| \leq i^\alpha) \\
 &= \sum_{i=1}^{\infty} E|X_1|^r I((i-1)^\alpha < |X_1| \leq i^\alpha) \sum_{n=i}^{\infty} n^{p\alpha-r\alpha-1} \\
 &\leq C \sum_{i=1}^{\infty} E|X_1|^r I((i-1)^\alpha < |X_1| \leq i^\alpha) i^{p\alpha-r\alpha} \\
 &\leq CE|X_1|^p < \infty.
 \end{aligned} \tag{2.9}$$

If $p < 2$, then we take $r = 2$. Since $r > p$, (2.9) still holds, and so $J_1 = J_2 < \infty$. $\square$

We next prove the sufficiency of Theorem 2.2 when the array is unbounded.

Lemma 2.4. *Let $\{X_n, n \geq 1\}$ be a sequence of identically distributed p^* -mixing random variables with $EX_1 = 0$ and $E|X_1|^p < \infty$ for some $p > 1/\alpha$ and $1/2 < \alpha \leq 1$. Assume that $\{a_{ni}, 1 \leq i \leq n, n \geq 1\}$ is an array of real numbers satisfying $a_{ni} = 0$ or $|a_{ni}| > 1$, and*

$$\sum_{i=1}^n |a_{ni}|^q \leq n \quad \text{for some } q > p. \tag{2.10}$$

Then (2.3) holds.

Proof. If $p < 2$, then we can take $\delta > 0$ such that $p < p + \delta < \min\{2, q\}$. Since $a_{ni} = 0$ or $|a_{ni}| > 1$, we have that $\sum_{i=1}^n |a_{ni}|^{p+\delta} \leq \sum_{i=1}^n |a_{ni}|^q \leq n$. Thus we may assume that (2.10) holds for some $p < q < 2$ when $p < 2$.

Let $S'_{nj} = \sum_{i=1}^j a_{ni} X_i I(|a_{ni} X_i| \leq n^\alpha)$ for $1 \leq j \leq n$ and $n \geq 1$. In view of $EX_i = 0$, we get

$$\begin{aligned}
 n^{-\alpha} \max_{1 \leq j \leq n} |ES'_{nj}| &= n^{-\alpha} \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} EX_i I(|a_{ni} X_i| > n^\alpha) \right| \\
 &\leq n^{-\alpha} \sum_{i=1}^n E|a_{ni} X_i| I(|a_{ni} X_i| > n^\alpha) \\
 &\leq n^{-p\alpha} \sum_{i=1}^n E|a_{ni} X_i|^p I(|a_{ni} X_i| > n^\alpha) \\
 &\leq n^{-p\alpha} \sum_{i=1}^n |a_{ni}|^p E|X_1|^p \\
 &\leq n^{-p\alpha} \left(\sum_{i=1}^n |a_{ni}|^q \right)^{p/q} n^{1-p/q} E|X_1|^p \\
 &\leq n^{1-p\alpha} E|X_1|^p \longrightarrow 0,
 \end{aligned} \tag{2.11}$$

since $p\alpha > 1$. Hence for n large enough, we have that $n^{-\alpha} \max_{1 \leq j \leq n} |ES'_{nj}| < \epsilon/2$. It follows that

$$\begin{aligned}
& \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| > \epsilon n^{\alpha}\right) \\
& \leq \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq i \leq n} |a_{ni} X_i| > n^{\alpha}\right) + \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} |S'_{nj}| > \epsilon n^{\alpha}\right) \\
& \leq \sum_{n=1}^{\infty} n^{p\alpha-2} \sum_{i=1}^n P(|a_{ni} X_i| > n^{\alpha}) + C \sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} |S'_{nj} - ES'_{nj}| > \frac{\epsilon n^{\alpha}}{2}\right) \\
& =: I + CJ.
\end{aligned} \tag{2.12}$$

For $1 \leq j \leq n-1$ and $n \geq 2$, let

$$I_{nj} = \left\{ 1 \leq i \leq n : n^{1/q}(j+1)^{-1/q} < |a_{ni}| \leq n^{1/q}j^{-1/q} \right\}. \tag{2.13}$$

Then $\{I_{nj}, 1 \leq j \leq n-1\}$ are disjoint, $\bigcup_{j=1}^{n-1} I_{nj} = \{1 \leq i \leq n : a_{ni} \neq 0\}$, and $\sum_{j=1}^k \#I_{nj} \leq k+1$ for $1 \leq k \leq n-1$, since

$$n \geq \sum_{\{1 \leq i \leq n : a_{ni} \neq 0\}} |a_{ni}|^q = \sum_{j=1}^{n-1} \sum_{i \in I_{nj}} |a_{ni}|^q \geq n \sum_{j=1}^k \frac{1}{j+1} \#I_{nj} \geq \frac{n}{(k+1)} \sum_{j=1}^k \#I_{nj}. \tag{2.14}$$

For convenience of notation, let $t = 1/(\alpha - 1/q)$. Since $a_{ni} = 0$ or $|a_{ni}| > 1$, and $\sum_{i=1}^n |a_{ni}|^q \leq n$, we have $a_{11} = 0$. It follows that

$$\begin{aligned}
I &= \sum_{n=2}^{\infty} n^{p\alpha-2} \sum_{j=1}^{n-1} \sum_{i \in I_{nj}} P(|a_{ni} X_i| > n^{\alpha}) \\
&\leq \sum_{n=2}^{\infty} n^{p\alpha-2} \sum_{j=1}^{n-1} P(|X_1|^t > nj^{t/q}) \#I_{nj} \\
&\leq \sum_{n=2}^{\infty} n^{p\alpha-2} \sum_{j=1}^{n-1} \#I_{nj} \sum_{k \geq \lceil nj^{t/q} \rceil} P(k < |X_1|^t \leq k+1)
\end{aligned}$$

$$\begin{aligned}
&\leq \sum_{n=2}^{\infty} n^{p\alpha-2} \sum_{k=n}^{\infty} P(k < |X_1|^t \leq k+1) \sum_{j=1}^{(n-1) \wedge [(k+1)/n]^{q/t}} \#I_{nj} \\
&\leq \sum_{n=2}^{\infty} n^{p\alpha-2} \sum_{k=n}^{\infty} P(k < |X_1|^t \leq k+1) n \wedge \left(\left[\left(\frac{k+1}{n} \right)^{q/t} \right] + 1 \right) \\
&\leq \sum_{n=1}^{\infty} n^{p\alpha-2} \sum_{k=n}^{[n^{1+t/q}]} P(k < |X_1|^t \leq k+1) \left(\left[\left(\frac{k+1}{n} \right)^{q/t} \right] + 1 \right) \\
&\quad + \sum_{n=1}^{\infty} n^{p\alpha-1} \sum_{k=[n^{1+t/q}]+1}^{\infty} P(k < |X_1|^t \leq k+1) \\
&=: I_1 + I_2.
\end{aligned} \tag{2.15}$$

Since $p\alpha - 2 - q/t = -\alpha(q-p) - 1 < -1$, we obtain

$$\begin{aligned}
I_1 &\leq C \sum_{n=1}^{\infty} n^{p\alpha-2-q/t} \sum_{k=n}^{[n^{1+t/q}]} P(k < |X_1|^t \leq k+1) k^{q/t} \\
&\leq C \sum_{k=1}^{\infty} P(k < |X_1|^t \leq k+1) k^{q/t} \sum_{n=[k^{q/(q+t)}]}^k n^{p\alpha-2-q/t} \\
&\leq C \sum_{k=1}^{\infty} P(k < |X_1|^t \leq k+1) k^{q/t - q\alpha(q-p)/(q+t)} \\
&\leq CE|X_1|^p < \infty.
\end{aligned} \tag{2.16}$$

We also obtain

$$\begin{aligned}
I_2 &\leq \sum_{k=1}^{\infty} P(k < |X_1|^t \leq k+1) \sum_{n=1}^{[k^{q/(t+q)}]} n^{p\alpha-1} \\
&\leq C \sum_{k=1}^{\infty} P(k < |X_1|^t \leq k+1) k^{p(\alpha-1/q)} \leq CE|X_1|^p < \infty.
\end{aligned} \tag{2.17}$$

From $I_1 < \infty$ and $I_2 < \infty$, we have $I < \infty$. Thus, it remains to show that $J < \infty$.

We have by Markov's inequality and Lemma 2.1 that for any $r \geq 2$,

$$\begin{aligned}
 J &\leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} E \max_{1 \leq j \leq n} |S'_{nj} - ES'_{nj}|^r \\
 &\leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} \left(\sum_{i=1}^n E |a_{ni} X_i|^2 I(|a_{ni} X_i| \leq n^\alpha) \right)^{r/2} \\
 &\quad + C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} \sum_{i=1}^n E |a_{ni} X_i|^r I(|a_{ni} X_i| \leq n^\alpha) \\
 &=: J_1 + J_2.
 \end{aligned} \tag{2.18}$$

Observe that for $r \geq q$ and $n > m$,

$$n \geq \sum_{j=1}^{n-1} \sum_{i \in I_{nj}} |a_{ni}|^q \geq n \sum_{j=1}^{n-1} \frac{1}{j+1} \#I_{nj} \geq n(m+1)^{r/q-1} \sum_{j=m}^{n-1} (j+1)^{-r/q} \#I_{nj}. \tag{2.19}$$

So $\sum_{j=m}^{n-1} j^{-r/q} \#I_{nj} \leq C m^{-(r/q-1)}$ for $r \geq q$ and $n > m$.

For J_1 and J_2 , we proceed with two cases.

(i) If $p \geq 2$, then we take r large enough such that $r > \max\{(p\alpha-1)/(\alpha-1/2), q\}$. Then we obtain that

$$\begin{aligned}
 J_1 &\leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} \left(\sum_{i=1}^n |a_{ni}|^2 \right)^{r/2} \\
 &\leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2} \left(\sum_{i=1}^n |a_{ni}|^q \right)^{r/2} \\
 &\leq C \sum_{n=1}^{\infty} n^{p\alpha-r\alpha-2+r/2} < \infty.
 \end{aligned} \tag{2.20}$$

The second inequality follows by the fact that $a_{ni} = 0$ or $|a_{ni}| > 1$.

Noting that $a_{11} = 0$, we also obtain that

$$\begin{aligned}
J_2 &= \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2} \sum_{j=1}^{n-1} \sum_{i \in I_{nj}} E|a_{ni}X_i|^r I(|a_{ni}X_i| \leq n^\alpha) \\
&\leq \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2+r/q} \sum_{j=1}^{n-1} j^{-r/q} \#I_{nj} E|X_1|^r I(|X_1|^t \leq n(j+1)^{t/q}) \\
&\leq \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2+r/q} \sum_{j=1}^{n-1} j^{-r/q} \#I_{nj} \sum_{0 \leq k \leq [n(j+1)^{t/q}]} E|X_1|^r I(k < |X_1|^t \leq k+1) \\
&= \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2+r/q} \sum_{j=1}^{n-1} j^{-r/q} \#I_{nj} \sum_{k=0}^{2n} E|X_1|^r I(k < |X_1|^t \leq k+1) \\
&\quad + \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2+r/q} \sum_{j=1}^{n-1} j^{-r/q} \#I_{nj} \sum_{k=2n+1}^{[n(j+1)^{t/q}]} E|X_1|^r I(k < |X_1|^t \leq k+1) \\
&=: J_3 + J_4.
\end{aligned} \tag{2.21}$$

Since $p\alpha - r\alpha - 2 + r/q < q\alpha - r\alpha - 2 + r/q = -(r-q)(\alpha - 1/q) - 1 < -1$ and $q > p$, we have that

$$\begin{aligned}
J_3 &= \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2+r/q} \sum_{k=0}^{2n} E|X_1|^r I(k < |X_1|^t \leq k+1) \sum_{j=1}^{n-1} j^{-r/q} \#I_{nj} \\
&\leq C \sum_{n=2}^{\infty} n^{p\alpha-r\alpha-2+r/q} \sum_{k=0}^{2n} E|X_1|^r I(k < |X_1|^t \leq k+1) \\
&\leq C \sum_{k=1}^{\infty} E|X_1|^r I(k < |X_1|^t \leq k+1) \sum_{n=[k/2]}^{\infty} n^{p\alpha-r\alpha-2+r/q} \\
&\leq C \sum_{k=1}^{\infty} E|X_1|^r I(k < |X_1|^t \leq k+1) k^{p\alpha-r\alpha-1+r/q} \\
&\leq C \sum_{k=1}^{\infty} P(k < |X_1|^t \leq k+1) k^{p\alpha-1} \\
&\leq CE|X_1|^p < \infty.
\end{aligned} \tag{2.22}$$

Since $1/t + 1/q - \alpha = 0$ and $p\alpha - 2 - q/t = -\alpha(q - p) - 1 < -1$, we also have that

$$\begin{aligned}
J_4 &\leq \sum_{n=2}^{\infty} n^{p\alpha - r\alpha - 2 + r/q} \sum_{k=2n+1}^{\lfloor n^{(q+t)/q} \rfloor} E|X_1|^r I(k < |X_1|^t \leq k+1) \sum_{j=\lfloor (k/n)^{q/t} \rfloor - 1}^{n-1} j^{-r/q} \#I_{nj} \\
&\leq C \sum_{n=2}^{\infty} n^{p\alpha - r\alpha - 2 + r/q} \sum_{k=2n+1}^{\lfloor n^{(q+t)/q} \rfloor} E|X_1|^r I(k < |X_1|^t \leq k+1) \left(\left\lfloor \left(\frac{k}{n} \right)^{q/t} \right\rfloor - 1 \right)^{-(r/q-1)} \\
&\leq C \sum_{k=5}^{\infty} E|X_1|^r I(k < |X_1|^t \leq k+1) k^{-(r-q)/t} \sum_{n=\lfloor k^{q/(q+t)} \rfloor}^{\lfloor k/2 \rfloor} n^{p\alpha - 2 - q/t} \\
&\leq C \sum_{k=5}^{\infty} E|X_1|^r I(k < |X_1|^t \leq k+1) k^{-(r-q)/t - (\alpha-1/q)(q-p)} \\
&\leq CE|X_1|^p < \infty.
\end{aligned} \tag{2.23}$$

From $J_3 < \infty$ and $J_4 < \infty$, we have $J_2 < \infty$.

(ii) If $p < 2$, then we take $r = 2$. As noted above, we may assume that $p < q < 2$. Since $r > q$, as in the case $p \geq 2$, we have $J_1 = J_2 \leq CE|X_1|^p < \infty$. $\square$

We now prove Theorem 2.2 by using Lemmas 2.3 and 2.4.

Proof of Theorem 2.2.

Sufficiency. Without loss of generality, we may assume that $\sum_{i=1}^n |a_{ni}|^q \leq n$ for some $q > p$. For $n \geq 1$, let

$$A_n = \{1 \leq i \leq n : |a_{ni}| \leq 1\}, \quad B_n = \{1 \leq i \leq n : |a_{ni}| > 1\}, \tag{2.24}$$

and let $a'_{ni} = a_{ni}$ if $i \in A_n$, $a'_{ni} = 0$ otherwise, and $a''_{ni} = a_{ni}$ if $i \in B_n$, $a''_{ni} = 0$ otherwise. Then

$$\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| \leq \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a'_{ni} X_i \right| + \max_{1 \leq j \leq n} \left| \sum_{i=1}^j a''_{ni} X_i \right|. \tag{2.25}$$

It follows that

$$\begin{aligned}
\sum_{n=1}^{\infty} n^{p\alpha-2} P \left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a_{ni} X_i \right| > \epsilon n^\alpha \right) &\leq \sum_{n=1}^{\infty} n^{p\alpha-2} P \left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a'_{ni} X_i \right| > \frac{\epsilon n^\alpha}{2} \right) \\
&\quad + \sum_{n=1}^{\infty} n^{p\alpha-2} P \left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j a''_{ni} X_i \right| > \frac{\epsilon n^\alpha}{2} \right) \\
&=: I + J.
\end{aligned} \tag{2.26}$$

By Lemma 2.3, we have $I < \infty$. By Lemma 2.4, we have $J < \infty$. Hence (2.3) holds.

Necessity. Choose, for each $n \geq 1$, $a_{n1} = \dots = a_{nn} = 1$. Then $\{a_{ni}\}$ satisfies (2.2). By (2.3), we obtain that

$$\sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} \left| \sum_{i=1}^j X_i \right| > \epsilon n^\alpha\right) < \infty \quad \forall \epsilon > 0, \quad (2.27)$$

which implies that

$$\sum_{n=1}^{\infty} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} |X_j| > \epsilon n^\alpha\right) < \infty \quad \forall \epsilon > 0. \quad (2.28)$$

Observe that

$$\begin{aligned} \infty &> \sum_{i=1}^{\infty} \sum_{n=2^{i-1}+1}^{2^i} n^{p\alpha-2} P\left(\max_{1 \leq j \leq n} |X_j| > \epsilon n^\alpha\right) \\ &\geq \begin{cases} \sum_{i=1}^{\infty} (2^{i-1})^{p\alpha-2} 2^{i-1} P\left(\max_{1 \leq j \leq 2^{i-1}} |X_j| > \epsilon (2^i)^\alpha\right) & \text{if } p\alpha \geq 2, \\ \sum_{i=1}^{\infty} (2^i)^{p\alpha-2} 2^{i-1} P\left(\max_{1 \leq j \leq 2^{i-1}} |X_j| > \epsilon (2^i)^\alpha\right) & \text{if } 1 < p\alpha < 2, \end{cases} \quad (2.29) \\ &\geq \begin{cases} \sum_{i=1}^{\infty} P\left(\max_{1 \leq j \leq 2^{i-1}} |X_j| > \epsilon (2^i)^\alpha\right) & \text{if } p\alpha \geq 2, \\ 2^{p\alpha-2} \sum_{i=1}^{\infty} P\left(\max_{1 \leq j \leq 2^{i-1}} |X_j| > \epsilon (2^i)^\alpha\right) & \text{if } 1 < p\alpha < 2. \end{cases} \end{aligned}$$

Hence we have that for any $\epsilon > 0$, $P(\max_{1 \leq j \leq 2^{i-1}} |X_j| > \epsilon (2^i)^\alpha) \rightarrow 0$ as $i \rightarrow \infty$, and so $P(\max_{1 \leq j \leq n} |X_j| > n^\alpha) \rightarrow 0$ as $n \rightarrow \infty$. The rest of the proof is same as that of Peligrad and Gut [3] and is omitted.

□

Remark 2.5. Taking $a_{ni} = 1$ for $1 \leq i \leq n$ and $n \geq 1$, we can immediately get Theorem 1.2 from Theorem 2.2. If the array $\{a_{ni}\}$ satisfies (1.4), then it satisfies (2.2): taking q such that $p < q < p/\delta$, we have

$$\sum_{i=1}^n |a_{ni}|^q \leq \max_{1 \leq i \leq n} |a_{ni}|^{q-p} \sum_{i=1}^n |a_{ni}|^p \leq C n^{\delta(q-p)/p} n^\delta \leq C n. \quad (2.30)$$

So the implication (i) $\Rightarrow$ (ii) of Theorem 1.3 follows from Theorem 2.2. As noted after Theorem 1.3, the implication (ii) $\Rightarrow$ (i) of Theorem 1.3 is not true. Therefore, our result extends the result of Peligrad and Gut [3] to a weighted average, and generalizes and sharpens the result of An and Yuan [8].

Acknowledgments

The author is grateful to the editor Leonid Shaikhet and the referees for the helpful comments and suggestions that considerably improved the presentation of this paper. This work was supported by the Korea Science and Engineering Foundation (KOSEF) Grant funded by the Korea government (MOST) (no. R01-2007-000-20053-0).

References

- [1] R. C. Bradley, "On the spectral density and asymptotic normality of weakly dependent random fields," *Journal of Theoretical Probability*, vol. 5, no. 2, pp. 355–373, 1992.
- [2] W. Bryc and W. Smoleński, "Moment conditions for almost sure convergence of weakly correlated random variables," *Proceedings of the American Mathematical Society*, vol. 119, no. 2, pp. 629–635, 1993.
- [3] M. Peligrad and A. Gut, "Almost-sure results for a class of dependent random variables," *Journal of Theoretical Probability*, vol. 12, no. 1, pp. 87–104, 1999.
- [4] S. Utev and M. Peligrad, "Maximal inequalities and an invariance principle for a class of weakly dependent random variables," *Journal of Theoretical Probability*, vol. 16, no. 1, pp. 101–115, 2003.
- [5] S. Gan, "Almost sure convergence for $\tilde{\rho}$ -mixing random variable sequences," *Statistics & Probability Letters*, vol. 67, no. 4, pp. 289–298, 2004.
- [6] A. Kucmaszewska, "On Chung-Teicher type strong law of large numbers for ρ^* -mixing random variables," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 140548, 10 pages, 2008.
- [7] Q. Wu and Y. Jiang, "Some strong limit theorems for $\tilde{\rho}$ -mixing sequences of random variables," *Statistics & Probability Letters*, vol. 78, no. 8, pp. 1017–1023, 2008.
- [8] J. An and D. Yuan, "Complete convergence of weighted sums for ρ^* -mixing sequence of random variables," *Statistics & Probability Letters*, vol. 78, no. 12, pp. 1466–1472, 2008.
- [9] G.-H. Cai, "Strong law of large numbers for ρ^* -mixing sequences with different distributions," *Discrete Dynamics in Nature and Society*, vol. 2006, Article ID 27648, 7 pages, 2006.
- [10] A. Kucmaszewska, "On complete convergence for arrays of rowwise dependent random variables," *Statistics & Probability Letters*, vol. 77, no. 11, pp. 1050–1060, 2007.
- [11] M.-H. Zhu, "Strong laws of large numbers for arrays of rowwise ρ^* -mixing random variables," *Discrete Dynamics in Nature and Society*, vol. 2007, Article ID 74296, 6 pages, 2007.
- [12] P. L. Hsu and H. Robbins, "Complete convergence and the law of large numbers," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 33, pp. 25–31, 1947.
- [13] P. Erdős, "On a theorem of Hsu and Robbins," *Annals of Mathematical Statistics*, vol. 20, pp. 286–291, 1949.
- [14] L. E. Baum and M. Katz, "Convergence rates in the law of large numbers," *Transactions of the American Mathematical Society*, vol. 120, pp. 108–123, 1965.