Research Article

Global Behavior of a Higher-Order Difference Equation

Tuo Li and Xiu-Mei Jia

Department of Mathematics, Hexi University, Zhangye, Gansu 734000, China

Correspondence should be addressed to Tuo Li, lituo1963@163.com

Received 18 March 2010; Revised 28 April 2010; Accepted 4 May 2010

Academic Editor: Elena Braverman

Copyright © 2010 T. Li and X.-M. Jia. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the global behavior of higher-order difference equation of the form $y_{n+1} = (y_n \exp(\beta(1 - 2 \sum_{i=0}^k a_i y_{n-i}))) / (1 - y_n + y_n \exp(\beta(1 - 2 \sum_{i=0}^k a_i y_{n-i})))$, $n = 0, 1, 2, \dots$, $y_{-k}, y_{-k+1}, \dots, y_0 \in (0, 1)$. Under some certain assumptions, it is proved that the positive equilibrium is globally asymptotical stable.

1. Introduction and Preliminaries

Nonlinear difference equations of order greater than one are of paramount importance in applications where the $(n + 1)$ th generation of the system depends on the previous k generations. Such equations appear naturally as discrete analogues and as numerical solutions of differential and delay differential equations which model various diverse phenomena in biology, ecology, physiology, physics, engineering, and economics [1–8]. The global character of difference equations is a most important topic and there have been many recent investigations and interest in the topic [1, 5–7, 9–14]. In particular, many researchers have paid attention to the global attractivity and convergence of the k th-order recursive sequence [2, 10, 14–19] and several approaches have been developed for finding the global character of difference equations; see [2, 6, 7, 9–15, 17, 19–21]. Moreover, we refer to [3, 4, 6, 7, 16] and the references therein for the oscillation and nonoscillation of difference equations. However, a large number of the literatures concerned with the *rational* difference equations and it is not enough to understand the global dynamics of a general difference equations, particularly irrational difference equation.

In this paper we study the global behavior of higher-order difference equation of some genotype selection model:

$$y_{n+1} = \frac{y_n \exp\left(\beta\left(1 - 2 \sum_{i=0}^k a_i y_{n-i}\right)\right)}{1 - y_n + y_n \exp\left(\beta\left(1 - 2 \sum_{i=0}^k a_i y_{n-i}\right)\right)}, \quad n = 1, 2, \dots, \quad (1.1)$$

where $\beta \in (0, \infty)$, $k \in \{0, 1, 2, \dots\}$, $a_0, a_1, \dots, a_k \in [0, 1]$, $\sum_{i=0}^k a_i = 1$, and initial conditions are $y_{-k}, y_{-k+1}, \dots, y_0 \in (0, 1)$. When $a_0 = a_1 = \dots = a_{k-1} = 0$, $a_k = 1$, (1.1) reduces to

$$y_{n+1} = \frac{y_n \exp(\beta(1 - 2y_{n-k}))}{1 - y_n + y_n \exp(\beta(1 - 2y_{n-k}))}, \quad n = 1, 2, \dots, \quad (1.2)$$

which was introduced by [6] as an example of a map generation by a simple mode for frequency dependent natural selection. The local stability of positive equilibrium $\bar{y} = 1/2$ of (1.2) was investigated by [6].

We note that the appearance of y_{n-i} ($i = 0, 1, \dots, k$) in the selection coefficient reflects the fact that the environment at the present time depends upon the activity of the population at some time in the past and that this in turn depends upon the gene frequency at that time. The points 0, 1/2, and 1 are the only equilibrium solutions of (1.1). One can easily see that $y_n \in [0, 1]$ for all $n = 1, 2, \dots$. If $y_N = 0$ for some $N \in \mathbb{N}$, then $y_n = 0$ for all $n \geq N$ and if $y_N = 1$ for some $N \in \mathbb{N}$, then $y_n = 1$ for all $n \geq N$. So in the following, we will restrict our attention to the difference equation:

$$y_{n+1} = \frac{y_n \exp\left(\beta\left(1 - 2 \sum_{i=0}^k a_i y_{n-i}\right)\right)}{1 - y_n + y_n \exp\left(\beta\left(1 - 2 \sum_{i=0}^k a_i y_{n-i}\right)\right)}, \quad y_{-k}, y_{-k+1}, \dots, y_0 \in (0, 1). \quad (1.3)$$

By introducing the substitution

$$y_n = \frac{x_n}{1 + x_n}, \quad (1.4)$$

then (1.3) becomes

$$x_{n+1} = x_n \exp\left(\beta\left(2 \sum_{i=0}^k \frac{a_i}{1 + x_{n-i}} - 1\right)\right), \quad n = 1, 2, \dots, \quad (1.5)$$

where $\beta \in (0, \infty)$, $k \in \{0, 1, 2, \dots\}$, $\sum_{i=0}^k a_i = 1$, and $x_{-k}, x_{-k+1}, \dots, x_0 \in (0, \infty)$ are arbitrary initial conditions.

In the sequel we will consider (1.5). It is clear that (1.5) has unique positive equilibrium point $\bar{x} = 1$.

In the following, we give some results which will be useful in our investigation of the behavior of solutions of (1.5), and the proof of lemmas can be found in [6].

Definition 1.1. The equilibrium point $\bar{x}$ of the equation $x_{n+1} = F(x_n, x_{n-1}, \dots, x_{n-k})$, $n = 0, 1, \dots$, is the point that satisfies the condition $\bar{x} = F(\bar{x}, \bar{x}, \dots, \bar{x})$.

Definition 1.2. (a) A sequence $\{x_n\}$ is said to be oscillate about zero or simply oscillate if the terms x_n are neither eventually all positive nor eventually all negative. Otherwise the sequence is called nonoscillatory. A sequence $\{x_n\}$ is called strictly oscillatory if for every $n_0 \geq 0$, there exist $n_1, n_2 \geq n_0$ such that $x_{n_1}x_{n_2} < 0$.

(b) A sequence $\{x_n\}$ is said to be oscillate about $\bar{x}$ if the sequence $\{x_n - \bar{x}\}$ oscillates. The sequence is called strictly oscillatory about $\bar{x}$ if the sequence $\{x_n - \bar{x}\}$ is strictly oscillatory.

Definition 1.3. Let $\bar{y}$ be an equilibrium point of equation $y_{n+1} = f(y_n, y_{n-1}, \dots, y_{n-k})$; then the equilibrium point $\bar{y}$ is called

- (a) locally stable if for every $\varepsilon > 0$ there exists $\delta > 0$ such that for all $y_k, y_{-k+1}, \dots, y_0 \in I$ with $|y_{-k} - \bar{y}| + |y_{-k+1} - \bar{y}| + \dots + |y_0 - \bar{y}| < \delta$, we have $|y_n - \bar{y}| < \varepsilon$, for all $n > -1$;
- (b) locally asymptotically stable if it is locally stable and if there exists $\gamma > 0$ such that for all $y_{-k}, y_{-k+1}, \dots, y_0 \in I$ with $|y_{-k} - \bar{y}| + |y_{-k+1} - \bar{y}| + \dots + |y_0 - \bar{y}| < \gamma$, we have $\lim_{n \rightarrow \infty} y_n = \bar{y}$;
- (c) a global attractor if for all $y_{-k}, y_{-k+1}, \dots, y_0 \in I$, we have $\lim_{n \rightarrow \infty} y_n = \bar{y}$.
- (d) globally asymptotically stable if $\bar{y}$ is locally stable and $\bar{y}$ is a global attractor.

Lemma 1.4. Assume that β is a positive real number and k is a nonnegative integer; then the following statements are true.

- (a) If $k = 0$, then every solution of (1.2) oscillates about $\bar{y} = 1/2$ if and only if $\beta > 2$.
- (b) If $k \geq 1$, then every solution of (1.2) oscillates about $\bar{y} = 1/2$ if and only if

$$\beta > 2 \frac{k^k}{(k+1)^{k+1}}. \quad (1.6)$$

Lemma 1.5. The linear difference equation

$$x_{n+k} + \sum_{i=1}^k p_i x_{n+k-i} = 0, \quad n = 0, 1, 2, \dots, \quad (1.7)$$

where $p_1, p_2, \dots, p_m \in \mathbb{R}$, is asymptotically stable provided that

$$\sum_{i=1}^k |p_i| < 1. \quad (1.8)$$

Lemma 1.6. The linear difference equation

$$x_{n+1} - x_n + \sum_{i=1}^m p_i x_{n-k_i} = 0, \quad n = 1, 2, \dots, \quad (1.9)$$

where $p_1, \dots, p_m \in (0, \infty)$ and $k_1, k_2, \dots, k_m$ are positive integers, is asymptotically stable provided that

$$\sum_{i=1}^m k_i p_i < 1. \quad (1.10)$$

Lemma 1.7. Consider the difference equation

$$x_{n+1} = x_n f(x_n, x_{n-k_1}, \dots, x_{n-k_r}); \quad (1.11)$$

one assumes that the function f satisfies the following hypotheses.

(H₁) $f \in C[[0, \infty) \times [0, \infty)^r, (0, \infty)]$, and $g \in C[[0, \infty)^{r+1}, (0, \infty)]$, where

$$\begin{aligned} g(u_0, u_1, \dots, u_r) &= u_0 f(u_0, u_1, \dots, u_r) \quad \text{for } u_0 \in (0, \infty), u_1, u_2, \dots, u_r \in [0, \infty), \\ g(0, u_1, \dots, u_r) &= \lim_{u_0 \rightarrow 0_+} g(u_0, u_1, \dots, u_r). \end{aligned} \quad (1.12)$$

(H₂) $f(u_0, u_1, \dots, u_r)$ is nonincreasing in $u_1, \dots, u_r$.

(H₃) The equation

$$f(x, x, \dots, x) = 1 \quad (1.13)$$

has a unique positive solution $\bar{x}$.

(H₄) Either the function $f(u_0, u_1, \dots, u_r)$ does not depend on u_0 or for every $x > 0$ and $u \geq 0$

$$[f(x, u, \dots, u) - f(\bar{x}, u, \dots, u)](x - \bar{x}) \leq 0 \quad (1.14)$$

with

$$[f(x, u, \dots, u) - f(\bar{x}, u, \dots, u)](x - \bar{x}) < 0 \quad \text{for } x \neq \bar{x}. \quad (1.15)$$

(H₅) The function $f(u_0, u_1, \dots, u_r)$ does not depend on u_0 or that for every $x, y \in (0, \infty)$ with $x \neq \bar{x}$

$$[xf(x, y, \dots, y) - \bar{x}f(\bar{x}, y, \dots, y)] > 0. \quad (1.16)$$

Furthermore assume that the function $F(x)$ is given by

$$F(x) = \bar{x}f(\bar{x}, x, \dots, x)^{k+1} \quad (1.17)$$

and has no periodic orbits of prime period 2; then $\bar{x}$ is a global attractor of all positive solutions of (1.11).

Lemma 1.8. Let $F \in C[[0, \infty), (0, \infty)]$ be a nonincreasing function, and let $\bar{x}$ denote the (unique) fixed point of F . Then the following statements are equivalent:

- (a) $\bar{x}$ is only fixed point of F^2 in $(0, \infty)$;
- (b) $F^2(x) > x$ for $0 < x < \bar{x}$.

2. Main Results

In this section, we will investigate the asymptotic stability and global behavior of the positive equilibrium point of (1.5).

Theorem 2.1. Assume that β is a positive real number and k is a nonnegative integer; then the following statements are true.

- (a) If $k = 0$, then every solution of (1.5) oscillates about $\bar{x} = 1$ if and only if $\beta > 2$.
- (b) If $k \geq 1$, $0 < \beta < \frac{4}{k(k+1)}$, $a_0 = 0$, then $\bar{x} = 1$ is an asymptotically stable solution of (1.5).
- (c) If $k \geq 1$, $0 < \beta \leq 2$, $1/2 < a_0 \leq 1$, then $\bar{x} = 1$ is an asymptotically stable solution of (1.5).

Proof. When $k = 0$, then $a_0 = 1$, $a_1 = a_2 = \dots = a_k = 0$, (1.5) becomes (1.2), and case (a) follows from Lemma 1.4(a).

When $k \neq 0$, the linearized equation of (1.5) about equilibrium point $\bar{x} = 1$ is

$$z_{n+1} - \left(1 - \frac{\beta a_0}{2}\right) z_n + \sum_{i=1}^k \frac{\beta a_i}{2} z_{n-i} = 0. \quad (2.1)$$

If $a_0 = 0$, by applying the Lemma 1.6, (2.1) is asymptotically stable provided that

$$\sum_{i=1}^k i \frac{\beta a_i}{2} < 1, \quad (2.2)$$

so for

$$0 < \beta < \frac{4}{k(k+1)}, \quad (2.3)$$

and $\bar{x} = 1$ is an asymptotically stable solution of (1.5). The proof of (b) is completed.

If $1/2 < a_0 \leq 1$, by applying Lemma 1.5, (2.1) is asymptotically stable provided that

$$\left| \left(1 - \frac{\beta a_0}{2}\right) \right| + \sum_{i=1}^k \left| \frac{\beta a_i}{2} \right| < 1, \quad (2.4)$$

so for $0 < \beta < 2$, and $\bar{x} = 1$ is an asymptotically stable solution of (1.5). The proof of (c) is completed. $\square$

Theorem 2.2. Assume that β is a positive real number and k is a positive integer. Then the equilibrium $\bar{x} = 1$ of (1.5) is globally asymptotically stable, when one of the following three cases holds:

- (a) $1 \leq k \leq 7$, $0 < \beta < 1/2(k+1)$, $a_0 = 0$;
- (b) $k \geq 8$, $0 < \beta < 4/k(k+1)$, $a_0 = 0$;
- (c) $k \geq 1$, $0 < \beta < 2c_0/(k+1)$, $1/2 < a_0 < 1$, where c_0 is a constant with $1/2 < c_0 < 1$.

Proof. If one of the three conditions (a), (b), (c) is satisfied, by applying Theorem 2.1, $\bar{x} = 1$ is an asymptotically stable solution of (1.5). To complete the proof it remains to show that $\bar{x}$ is a global attractor of all positive solutions of (1.5). We will employ Lemma 1.7; set

$$f(u_0, u_1, \dots, u_k) = \exp \left[\beta \left(2 \sum_{i=0}^k \frac{a_i}{1+u_i} - 1 \right) \right]; \quad (2.5)$$

clearly equation $x_{n+1} = x_n f(x_n, x_{n-1}, \dots, x_{n-k})$ satisfies the hypotheses (H₁)–(H₄) of Lemma 1.7. Set

$$h(x, y) = x f(x, y, \dots, y) = x \exp \left(\beta \left(\frac{1-y-2a_0}{1+y} + \frac{2a_0}{1+x} \right) \right). \quad (2.6)$$

If $0 \leq a_0 \leq 1$, $x \neq \bar{x}$, $0 < \beta < 2$, then

$$\begin{aligned} \frac{\partial h}{\partial x} &= \exp \left(\beta \left(\frac{1-y-2a_0}{1+y} + \frac{2a_0}{1+x} \right) \right) \left(1 - \frac{2\beta a_0 x}{(1+x)^2} \right) \\ &= \exp \left(\beta \left(\frac{1-y-2a_0}{1+y} + \frac{2a_0}{1+x} \right) \right) \left(\frac{x^2 + (2-2\beta a_0)x + 1}{(1+x)^2} \right) > 0; \end{aligned} \quad (2.7)$$

hence, the function $f(u_0, u_1, \dots, u_r)$ does not depend on u_0 or that for every $x, y \in (0, \infty)$ with $x \neq \bar{x}$,

$$[x f(x, y, \dots, y) - \bar{x} f(\bar{x}, y, \dots, y)] > 0. \quad (2.8)$$

Furthermore assume that the function $F(x)$ is given by

$$F(x) = \bar{x} f(\bar{x}, x, \dots, x)^{k+1} = \exp \left((k+1) \beta \left(\frac{2(1-a_0)}{1+x} + a_0 - 1 \right) \right). \quad (2.9)$$

Set

$$\Phi(x) = x - F(F(x)); \quad (2.10)$$

then

$$\begin{aligned}
 \Phi'(x) &= 1 - F'(F(x))F'(x) \\
 &= 1 - \left[\exp\left((k+1)\beta\left(\frac{2(1-a_0)}{1+x} + a_0 - 1\right)\right)(k+1)\beta\frac{2(1-a_0)}{(1+x)^2} \right] \Bigg|_{F(x)} \\
 &\quad \times \exp\left((k+1)\beta\left(\frac{2(1-a_0)}{1+x} + a_0 - 1\right)\right)(k+1)\beta\frac{2(1-a_0)}{(1+x)^2}.
 \end{aligned} \tag{2.11}$$

When $0 < \beta(k+1)(1-a_0) < c_0$, $0 < x < 1$, then $1 < F(x) < \exp((k+1)\beta(1-a_0)) < e^{c_0}$. Hence,

$$\begin{aligned}
 \Phi'(x) &= 1 - \left[\exp\left((k+1)\beta\left(\frac{2(1-a_0)}{1+x} + a_0 - 1\right)\right)(k+1)\beta\frac{2(1-a_0)}{(1+x)^2} \right] \Bigg|_{F(x)} \\
 &\quad \times \left(\exp\left((k+1)\beta\left(\frac{2(1-a_0)}{1+x} + a_0 - 1\right)\right)(k+1)\beta\frac{2(1-a_0)}{(1+x)^2} \right. \\
 &\quad \left. \geq 1 - \left[\exp\left((k+1)\beta\left(\frac{2(1-a_0)}{1+x} + a_0 - 1\right)\right)(k+1)\beta\frac{2(1-a_0)}{(1+x)^2} \right] \Bigg|_{F(x)=1} \right. \\
 &\quad \left. \times \exp\left((k+1)\beta\left(\frac{2(1-a_0)}{1+x} + a_0 - 1\right)\right)(k+1)\beta\frac{2(1-a_0)}{(1+x)^2} \right] \Bigg|_{x=0} \\
 &\geq 1 - (k+1)^2\beta^2(1-a_0)^2e^{c_0} \geq 1 - c_0^2e^{c_0} = 0,
 \end{aligned} \tag{2.12}$$

where c_0 is a constant with $1 - c_0^2e^{c_0} = 0$, and clearly $1/2 < c_0 < 1$. So $\Phi(x)$ is an increasing function, hence

$$\Phi(x) = x - F^2(x) < \Phi(1) = 0, \tag{2.13}$$

and so, $F^2(x) > x$ for $0 < x < 1$, by Lemma 1.8, we know that $\bar{x}$ is only fixed point of F^2 in $(0, \infty)$, and by Lemma 1.7, $\bar{x}$ is a global attractor of all positive solutions of (1.5). The proof is complete. $\square$

References

- [1] R. P. Agarwal, *Difference Equations and Inequalities*, vol. 228 of *Monographs and Textbooks in Pure and Applied Mathematics*, Marcel Dekker, New York, NY, USA, 2nd edition, 2000.
- [2] M. J. Douraki, M. Dehghan, and A. Razavi, "On the global behavior of higher order recursive sequences," *Applied Mathematics and Computation*, vol. 169, no. 2, pp. 819–831, 2005.
- [3] L. Erbe, J. Baoguo, and A. Peterson, "Nonoscillation for second order sublinear dynamic equations on time scales," *Journal of Computational and Applied Mathematics*, vol. 232, no. 2, pp. 594–599, 2009.
- [4] L. Erbe and A. Peterson, "Oscillation criteria for second-order matrix dynamic equations on a time scale," *Journal of Computational and Applied Mathematics*, vol. 141, no. 1-2, pp. 169–185, 2002.
- [5] W. G. Kelley and A. C. Peterson, *Difference Equations*, Academic Press, Boston, Mass, USA, 1991, An introduction with application.

- [6] V. L. Kocić and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, vol. 256 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1993.
- [7] M. R. S. Kulenović and G. Ladas, *Dynamics of Second Order Rational Difference Equations*, Chapman and Hall/CRC, Boca Raton, Fla, USA, 2002, With open problems and conjecture.
- [8] V. Tkachenko and S. Trofimchuk, "Global stability in difference equations satisfying the generalized Yorke condition," *Journal of Mathematical Analysis and Applications*, vol. 303, no. 1, pp. 173–187, 2005.
- [9] R. P. Agarwal, W. T. Li, and P. Y. H. Pang, "Asymptotic behavior of a class of nonlinear delay difference equations," *Journal of Difference Equations and Applications*, vol. 8, no. 8, pp. 719–728, 2002.
- [10] E. Camouzis and G. Ladas, *Dynamics of Third-Order Rational Difference Equations with Open Problems and Conjectures*, vol. 5 of *Advances in Discrete Mathematics and Applications*, Chapman and Hall/CRC, Boca Raton, Fla, USA, 2008.
- [11] H. El-Owaidy and H. Y. Mohamed, "On the global attractivity of systems of nonlinear difference equations," *Applied Mathematics and Computation*, vol. 135, no. 2-3, pp. 377–382, 2003.
- [12] V. L. Kocić and G. Ladas, "Global attractivity in a nonlinear second-order difference equation," *Communications on Pure and Applied Mathematics*, vol. 48, no. 9-10, pp. 1115–1122, 1995.
- [13] G. Ladas, "On the recursive sequence $x_{n+1} = (\alpha + \beta x_n + \gamma x_{n-1}) / (A + Bx_n + Cx_{n-1})$," *Journal of Difference Equations and Applications*, vol. 1, no. 3, pp. 317–321, 1995.
- [14] G. Papaschinopoulos and C. J. Schinas, "Global asymptotic stability and oscillation of a family of difference equations," *Journal of Mathematical Analysis and Applications*, vol. 294, no. 2, pp. 614–620, 2004.
- [15] M. Aloqeili, "Global stability of a rational symmetric difference equation," *Applied Mathematics and Computation*, vol. 215, no. 3, pp. 950–953, 2009.
- [16] M. Dehghan and M. J. Douraki, "The oscillatory character of the recursive sequence $x_{n+1} = \alpha + \beta x_{n-k+1} / A + Bx_{n-2k+1}$," *Applied Mathematics and Computation*, vol. 175, no. 1, pp. 38–48, 2006.
- [17] A. E. Hamza and S. G. Barbary, "Attractivity of the recursive sequence $x_{n+1} = (\alpha + \beta x_n F(x_{n-1}, \dots, x_{n-k}))$," *Mathematical and Computer Modelling*, vol. 48, no. 11-12, pp. 1744–1749, 2008.
- [18] L. X. Hu, W. T. Li, and S. Stević, "Global asymptotic stability of a second order rational difference equation," *Journal of Difference Equations and Applications*, vol. 14, no. 8, pp. 779–797, 2008.
- [19] M. Saleh and S. Abu-Baha, "Dynamics of a higher order rational difference equation," *Applied Mathematics and Computation*, vol. 181, no. 1, pp. 84–102, 2006.
- [20] M. Dehghan and R. Mazrooei-Sebdani, "The characteristics of a higher-order rational difference equation," *Applied Mathematics and Computation*, vol. 182, no. 1, pp. 521–528, 2006.
- [21] X. Yang, H. Lai, D. J. Evans, and G. M. Megson, "Global asymptotic stability in a rational recursive sequence," *Applied Mathematics and Computation*, vol. 158, no. 3, pp. 703–716, 2004.