Research Article

On the Basic k -nacci Sequences in Finite Groups

Ömür Deveci^{1,2} and Erdal Karaduman^{1,2}

¹ Department of Mathematics, Faculty of Science and Letters, Kafkas University, 36100 Kars, Turkey

² Department of Mathematics, Faculty of Science, Atatürk University, 25240 Erzurum, Turkey

Correspondence should be addressed to Ömür Deveci, odeveci36@yahoo.com.tr

Received 13 December 2010; Accepted 16 March 2011

Academic Editor: Carlo Piccardi

Copyright © 2011 Ö. Deveci and E. Karaduman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We define the basic k -nacci sequences and the basic periods of these sequences in finite groups, then we obtain the basic periods of the basic k -nacci sequences and the periods of the k -nacci sequences in symmetric group S_4 , its subgroups, and binary polyhedral groups which related with these groups.

1. Introduction

The study of Fibonacci sequences in groups began with the earlier work of Wall [1], where the ordinary Fibonacci sequences in cyclic groups were investigated. In the mid-eighties, Wilcox extended the problem to Abelian groups [2]. The theory is expanded to some finite simple groups by Campbell et al. [3]. There, they defined the Fibonacci length of the Fibonacci orbit and the basic Fibonacci length of the basic Fibonacci orbit in a 2-generator group. The concept of Fibonacci length for more than two generators has also been considered; see, for example, [4, 5]. Also, the theory has been expanded to the nilpotent groups; see, for example, [6, 7]. Other works on Fibonacci length are discussed in, for example, [8–10]. Knox proved that the periods of k -nacci (k -step Fibonacci) sequences in dihedral groups were equal to $2k + 2$ [11]. Deveci, Karaduman, and Campbell examined the period of the k -nacci sequences in some finite binary polyhedral groups in [12]. Recently, k -nacci sequences have been investigated; see, for example, [13, 14].

This paper defines the basic k -nacci sequences and the periods of these sequences in finite groups and discusses the basic periods of the basic k -nacci sequences and the periods of the k -nacci sequences in the symmetric group S_4 , alternating group A_4 , D_2 four-group, and binary polyhedral groups $\langle 2, 3, 4 \rangle$ and $\langle 2, 3, 3 \rangle$ with related S_4 and A_4 , respectively. We

consider the groups S_4 , A_4 , $\langle 2, 3, 4 \rangle$, and $\langle 2, 3, 3 \rangle$ both as 2-generator and as 3-generator groups.

A *k-nacci sequence* in a finite group is a sequence of group elements $x_0, x_1, x_2, \dots, x_n, \dots$ for which, given an initial (seed) set $x_0, x_1, x_2, \dots, x_{j-1}$, each element is defined by

$$x_n = \begin{cases} x_0 x_1 \cdots x_{n-1} & \text{for } j \leq n < k, \\ x_{n-k} x_{n-k+1} \cdots x_{n-1} & \text{for } n \geq k. \end{cases} \quad (1.1)$$

We also require that the initial elements of the sequence $x_0, x_1, x_2, \dots, x_{j-1}$ generate the group, thus forcing the *k-nacci* sequence to reflect the structure of the group. The *k-nacci* sequence of a group G generated by $x_0, x_1, x_2, \dots, x_{j-1}$ is denoted by $F_k(G; x_0, x_1, \dots, x_{j-1})$ [11].

A sequence of group elements is *periodic* if, after a certain point, it consists only of repetitions of a fixed subsequence. The number of elements in the repeating subsequence is called the *period of the sequence*. For example, the sequence $a, b, c, d, e, b, c, d, e, b, c, d, e, \dots$ is periodic after the initial element a and has period 4. A sequence of group elements is *simply periodic* with period k if the first k elements in the sequence form a repeating subsequence. For example, the sequence $a, b, c, d, e, f, a, b, c, d, e, f, a, b, c, d, e, f, \dots$ is simply periodic with period 6. In [11], Knox had denoted the period of a *k-nacci* sequence $F_k(G; x_0, x_1, \dots, x_{j-1})$ by $P_k(G; x_0, x_1, \dots, x_{j-1})$.

Definition 1.1. For a finitely generated group $G = \langle A \rangle$, where $A = \{a_1, a_2, \dots, a_n\}$, the sequence $x_i = a_{i+1}$, $0 \leq i \leq n-1$, $x_{i+n} = \prod_{j=1}^n x_{i+j-1}$, $i \geq 0$ is called the *Fibonacci orbit* of G with respect to the generating set A , denoted as $F_A(G)$ [4].

Definition 1.2. If $F_A(G)$ is simply periodic, then the period of the sequence is called the *Fibonacci length* of G with respect to generating set A , written, $LEN_A(G)$ [4].

Notice that the orbit of a *k-generated* group is a *k-nacci* sequence.

Let G be a finite *j-generator* group, and let X be the subset of $\underbrace{G \times G \times G \cdots \times G}_j$ such that $(x_0, x_1, \dots, x_{j-1}) \in X$ if and only if G is generated by $x_0, x_1, \dots, x_{j-1}$. We call $(x_0, x_1, \dots, x_{j-1})$ a *generating j-tuple* for G .

2. Basic Period of Basic *k-nacci* Sequence

To examine the concept more fully, we study the action of automorphism group $\text{Aut}G$ of G on X and on the *k-nacci* sequences $F_k(G; x_0, x_1, \dots, x_{j-1})$, $(x_0, x_1, \dots, x_{j-1}) \in X$. Now, $\text{Aut}G$ consists of all isomorphism $\theta : G \rightarrow G$ and if $\theta \in \text{Aut}G$ and $(x_0, x_1, \dots, x_{j-1}) \in X$, then $(x_0\theta, x_1\theta, \dots, x_{j-1}\theta) \in X$.

For a subset $A \subseteq G$ and $\theta \in \text{Aut}G$, the image of A under θ is

$$A\theta = \{a\theta : a \in A\}. \quad (2.1)$$

Definition 2.1. For a generating pair $(x, y) \in X$, the basic *Fibonacci orbit* $\overline{F}_{x,y}$ of the *basic length* m is defined by the sequence $\{b_i\}$ of elements of G such that

$$b_0 = x, \quad b_1 = y, \quad b_{i+2} = b_i b_{i+1}, \quad i \geq 0, \quad (2.2)$$

where $m \geq 1$ is the least integer with

$$b_0 = b_m\theta, \quad b_1 = b_{m+1}\theta, \quad (2.3)$$

for some $\theta \in \text{Aut}G$. Since b_m, b_{m+1} generate G , it follows that θ is uniquely determined. For more information, see [3].

Lemma 2.2. *Let $(x_0, x_1, \dots, x_{j-1}) \in X$ and let $\theta \in \text{Aut}G$, then $(F_k(G : x_0, x_1, \dots, x_{j-1}))\theta = F_k(G : x_0\theta, x_1\theta, \dots, x_{j-1}\theta)$.*

Proof. Let $F_k(G : x_0, x_1, \dots, x_{j-1}) = \{b_i\}$. The result is obvious since $\{b_i\}\theta = \{b_i\theta\}$ and

$$b_{i+k}\theta = (b_i b_{i+1} \cdots b_{i+k-1})\theta = b_i\theta b_{i+1}\theta \cdots b_{i+k-1}\theta. \quad (2.4)$$

Each generating j -tuple $(x_0, x_1, \dots, x_{j-1}) \in X$ maps to $|\text{Aut}G|$ distinct elements of X under the action of elements of $\text{Aut}G$. Hence, there are

$$d_j(G) = |X|/|\text{Aut}G|, \quad (2.5)$$

(where $|X|$ means the number of elements of X) nonisomorphic generating j -tuples for G . The notation $d_j(G)$ was introduced in [15].

Suppose that ω elements of $\text{Aut}G$ map $F_k(G : x_0, x_1, \dots, x_{j-1})$ into itself, then there are $|\text{Aut}G|/\omega$ distinct k -nacci sequences $F_k(G : x_0\theta, x_1\theta, \dots, x_{j-1}\theta)$ for $\theta \in \text{Aut}G$. $\square$

Definition 2.3. For a j -tuple $(x_0, x_1, \dots, x_{j-1}) \in X$, the basic k -nacci sequence $\overline{F}_k(G : x_0, x_1, \dots, x_{j-1})$ of the basic period m is a sequence of group elements $b_0, b_1, b_2, \dots, b_n, \dots$ for which, given an initial (seed) set $b_0 = x_0, b_1 = x_1, b_2 = x_2, \dots, b_{j-1} = x_{j-1}$, each element is defined by

$$b_n = \begin{cases} b_0 b_1 \cdots b_{n-1} & \text{for } j \leq n < k, \\ b_{n-k} b_{n-k+1} \cdots b_{n-1} & \text{for } n \geq k, \end{cases} \quad (2.6)$$

where $m \geq 1$ is the least integer with

$$b_0 = b_m\theta, \quad b_1 = b_{m+1}\theta, \quad b_2 = b_{m+2}\theta, \quad \dots, \quad b_{k-1} = b_{m+k-1}\theta, \quad (2.7)$$

for some $\theta \in \text{Aut}G$. Since G is a finite j -generator group and $b_m, b_{m+1}, \dots, b_{m+j-1}$ generate G , it follows that θ is uniquely determined. The basic k -nacci sequence $\overline{F}_k(G : x_0, x_1, \dots, x_{j-1})$ is finite containing m element.

In this paper, we denote the basic period of the basic k -nacci sequence $\overline{F}_k(G : x_0, x_1, \dots, x_{j-1})$ by $BP_k(G; x_0, x_1, \dots, x_{j-1})$.

From the definitions, it is clear that the periods of the k -nacci sequences and the basic k -nacci sequences in a finite group depend on the chosen generating set and the order of the generating elements.

Theorem 2.4. Let G be a finite group and $(x_0, x_1, \dots, x_{j-1}) \in X$. If $P_k(G; x_0, x_1, \dots, x_{j-1}) = n$ and $BP_k(G; x_0, x_1, \dots, x_{j-1}) = m$, then m divides n , and there are n/m elements of $\text{Aut}G$ which map $F_k(G; x_0, x_1, \dots, x_{j-1})$ into itself.

Proof. We have $n = m\lambda$ where λ is the order of automorphism $\theta \in \text{Aut}G$ since

$$\begin{aligned} F_k(G; x_0, x_1, \dots, x_{j-1}) &= \overline{F}_k(G; x_0, x_1, \dots, x_{j-1}) \cup \overline{F}_k(G; x_0\theta, x_1\theta, \dots, x_{j-1}\theta) \\ &\cup \overline{F}_k(G; x_0\theta^2, x_1\theta^2, \dots, x_{j-1}\theta^2) \cup \dots \end{aligned} \quad (2.8)$$

and $BP_k(G; x_0, x_1, \dots, x_{j-1}) = BP_k(G; x_0\theta, x_1\theta, \dots, x_{j-1}\theta)$. Clearly, $1, \theta, \theta^2, \dots, \theta^{\lambda-1}$ map $F_k(G; x_0, x_1, \dots, x_{j-1})$ into itself. $\square$

3. Applications

Definition 3.1. The *polyhedral group* (l, m, n) for $l, m, n > 1$ is defined by the presentation

$$\langle x, y, z : x^l = y^m = z^n = xyz = e \rangle, \quad (3.1)$$

or

$$\langle x, y : x^l = y^m = (xy)^n = e \rangle. \quad (3.2)$$

The *polyhedral group* (l, m, n) is finite if and only if the number

$$\mu = lmn \left(\frac{1}{l} + \frac{1}{m} + \frac{1}{n} - 1 \right) = mn + nl + lm - lmn \quad (3.3)$$

is positive, that is, in the cases $(2, 2, n)$, $(2, 3, 3)$, $(2, 3, 4)$, and $(2, 3, 5)$. Its order is $2lmn/\mu$. A_4 , S_4 , and A_5 are the groups $(2, 3, 3)$, $(2, 3, 4)$, and $(2, 3, 5)$, respectively. Also, the groups A_4 , S_4 , and A_5 being isomorphic to the groups of rotations of the regular tetrahedron, octahedron, and icosahedron. Using Tietze transformations, we may show that $(l, m, n) \cong (m, n, l) \cong (n, l, m)$. For more information on these groups, see, [16, 17, pp. 67-68].

Definition 3.2. The *binary polyhedral group* $\langle l, m, n \rangle$, for $l, m, n > 1$, is defined by the presentation

$$\langle x, y, z : x^l = y^m = z^n = xyz \rangle, \quad (3.4)$$

or

$$\langle x, y : x^l = y^m = (xy)^n \rangle. \quad (3.5)$$

The *binary polyhedral group* $\langle l, m, n \rangle$ is finite if and only if the number $k = lmn(1/l + 1/m + 1/n - 1) = mn + nl + lm - lmn$ is positive. Its order is $4lmn/k$.

For more information on these groups, see [17, pp. 68–71].

Definition 3.3. Let $f_n^{(k)}$ denote the n th member of the k -step Fibonacci sequence defined as

$$f_n^{(k)} = \sum_{j=1}^k f_{n-j}^{(k)} \quad \text{for } n > k, \quad (3.6)$$

with boundary conditions $f_i^{(k)} = 0$ for $1 \leq i < k$ and $f_k^{(k)} = 1$. Reducing this sequence by a modulo m , we can get a repeating sequence, which we denote by

$$f(k, m) = (f_1^{(k,m)}, f_2^{(k,m)}, \dots, f_n^{(k,m)} \dots), \quad (3.7)$$

where $f_i^{(k,m)} = f_i^{(k)} \pmod{m}$. We then have that $(f_1^{(k,m)}, f_2^{(k,m)}, \dots, f_k^{(k,m)}) = (0, 0, \dots, 0, 1)$, and it has the same recurrence relation as in (3.6) [18].

Theorem 3.4 ($f(k, m)$ is a periodic sequence [18]). Let $h_k(m)$ denote the smallest period of $f(k, m)$, called the period of $f(k, m)$ or the wall number of the k -step Fibonacci sequence modulo m .

Theorem 3.5. The periods of the k -nacci sequences and the basic periods of the basic k -nacci sequences in the group S_4 are as follows.

if the group is defined by the presentation $S_4 = \langle x, y, z : x^2 = y^3 = z^4 = xyz = e \rangle$, then

- (i) if $k = 2$, $P_2(S_4; y, z, x) = 18$ and $BP_2(S_4; y, z, x) = 9$,
- (ii) if $k > 2$, $P_k(S_4; x, y, z) = 6k + 6$ and $BP_k(S_4; x, y, z) = 3k + 3$.

If S_4 has the presentation $S_4 = \langle x, y : x^2 = y^3 = (xy)^4 = e \rangle$, then

- (i') if $k = 2$, $P_2(S_4; x, y) = 18$ and $BP_2(S_4; x, y) = 9$,
- (ii') if $k > 2$, $P_k(S_4; x, y) = 6k + 6$ and $BP_k(S_4; x, y) = 3k + 3$.

Proof. Firstly, let us consider the 3-generator case. We first note that $|x| = 2$, $|y| = 3$, and $|z| = 4$ (where $|x|$ means the order of x).

- (i) If $k = 2$, we have the sequence for the generating triple (y, z, x) ,

$$\begin{aligned} & y, z, x, y^2, xy^2, y^2xy^2, z^2y, z^2yz^3y, yxy, xyx, \\ & xy^2, x, xy^2x, y^2x, yxy, yxz, zy, y^2xy^2, y, z, x, \dots, \end{aligned} \quad (3.8)$$

which has period 18 and the basic period 9 since $x\theta = x$, $y\theta = xyx$, and $z\theta = xy^2$, where θ is the inner automorphism induced by conjugation by x .

- (ii) If $k = 3$, we have the sequence for the generating triple (x, y, z) ,

$$\begin{aligned} & x, y, z, e, x, y^2, xy^2, xzy^2, x, y, yxy^2, xzy^2, x, \\ & y^2, yx, e, x, y, xy, z^2, x, y^2, zy, z^2, x, y, z, \dots, \end{aligned} \quad (3.9)$$

which has period 24 and the basic period 12 since $x\theta = x$, $y\theta = y^2$, and $z\theta = yx$ where θ is an outer automorphism of order 2.

If $k \geq 4$, the first k elements of sequence for the generating triple (x, y, z) are

$$x_0 = x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = xyz, \quad x_4 = (xyz)^2 \dots, \quad x_{k-1} = (xyz)^{2^{k-4}}. \quad (3.10)$$

Thus, using the above information, sequence reduces to

$$x_0 = x, \quad x_1 = y, \quad x_2 = z, \quad x_3 = e, \dots, e, \quad x_{k-1} = e, \quad (3.11)$$

where $x_j = e$ for $3 \leq j \leq k-1$. Thus,

$$\begin{aligned} x_k &= e, \quad x_{k+1} = x, \quad x_{k+2} = y^2, \quad x_{k+3} = xy^2, \quad x_{k+4} = xzy^2, \\ x_{k+5} &= e, \quad \dots, \quad e, \quad x_{2k+1} = e, \quad x_{2k+2} = x, \quad x_{2k+3} = y, \\ x_{2k+4} &= yxy^2, \quad x_{2k+5} = xzy^2, \quad x_{2k+6} = e, \dots, \quad e, \quad x_{3k+2} = e, \\ x_{3k+3} &= x, \quad x_{3k+4} = y^2, \quad x_{3k+5} = yx, \quad x_{3k+6} = e, \quad \dots, \quad e, \quad x_{4k+3} = e, \\ x_{4k+4} &= x, \quad x_{4k+5} = y, \quad x_{4k+6} = xy, \quad x_{4k+7} = z^2, \\ x_{4k+8} &= e, \quad \dots, \quad e, \quad x_{5k+4} = e, \quad x_{5k+5} = x, \quad x_{5k+6} = y^2, \\ x_{5k+7} &= zy, \quad x_{5k+8} = z^2, \quad x_{5k+9} = e, \dots, \quad e, \quad x_{6k+5} = e, \end{aligned} \quad (3.12)$$

where $x_j = e$ for $k+5 \leq j \leq 2k+1$, $2k+6 \leq j \leq 3k+2$, $3k+6 \leq j \leq 4k+3$, $4k+8 \leq j \leq 5k+4$, and $5k+9 \leq j \leq 6k+5$.

We also have

$$x_{6k+6} = \prod_{i=5k+6}^{6k+5} x_i = x, \quad x_{6k+7} = \prod_{i=5k+7}^{6k+6} x_i = y, \quad x_{6k+8} = \prod_{i=5k+8}^{6k+7} x_i = z. \quad (3.13)$$

Since the elements succeeding x_{6k+6} , x_{6k+7} , and x_{6k+8} depend on x , y , and z for their values, the cycle begins again with the $6k+6^{\text{th}}$ element, that is, $x_0 = x_{6k+6}$, $x_1 = x_{6k+7}$, $x_2 = x_{6k+8}$, $\dots$. Thus, $P_k(S_4; x, y, z) = 6k+6$.

It is easy to see from the above sequence that

$$x_{3k+3} = x, \quad x_{3k+4} = y^2, \quad x_{3k+5} = yx, \quad x_{3k+6} = e, \quad \dots, e, \quad x_{4k+2} = e. \quad (3.14)$$

$BP_k(S_4; x, y, z) = 3k+3$ since $x\theta = x$, $y\theta = y^2$, and $z\theta = yx$ where θ is an outer automorphism of order 2.

Secondly, let us consider the 2-generator case. We first note that $|x| = 2$, $|y| = 3$, and $|xy| = 4$.

(i') If $k = 2$, $P_2(S_4; x, y) = 18$ and $BP_2(S_4; x, y) = 9$ since $x\theta = x$ and $y\theta = xyx$ where θ is the inner automorphism induced by conjugation by x .

- (ii') If $k > 2$, $P_k(S_4; x, y) = 6k + 6$ and $BP_k(S_4; x, y) = 3k + 3$ since $x\theta = x$ and $y\theta = y^2$ where θ is an outer automorphism of order 2.

The proofs are similar to above and are omitted. $\square$

Theorem 3.6. *The periods of the k -nacci sequences and the basic periods of the basic k -nacci sequences in the binary polyhedral group $\langle 2, 3, 4 \rangle$ are as follows.*

If the group is defined by the presentation $\langle 2, 3, 4 \rangle = \langle x, y, z : x^2 = y^3 = z^4 = xyz \rangle$, then

- (i) if $k = 2$, $P_2(\langle 2, 3, 4 \rangle; y, z, x) = 18$ and $BP_2(\langle 2, 3, 4 \rangle; y, z, x) = 9$,
(ii) if $k > 2$, $P_k(\langle 2, 3, 4 \rangle; x, y, z) = 6k + 6$ and $BP_k(\langle 2, 3, 4 \rangle; x, y, z) = 6k + 6$.

If the group is defined by the presentation $\langle 2, 3, 4 \rangle = \langle x, y : x^2 = y^3 = (xy)^4 \rangle$, then

- (i') if $k = 2$, $P_2(\langle 2, 3, 4 \rangle; x, y) = 18$ and $BP_2(\langle 2, 3, 4 \rangle; x, y) = 9$,
(ii') if $k > 2$, $P_k(\langle 2, 3, 4 \rangle; x, y) = 6k + 6$ and $BP_k(\langle 2, 3, 4 \rangle; x, y) = 6k + 6$.

Proof. Firstly, let us consider the 2-generator case. We first note that $|x| = 4$, $|y| = 6$, and $|xy| = 8$.

- (i') If $k = 2$, we have the sequence for the generating pair (x, y) ,

$$\begin{aligned} x, y, xy, yxy, xy^2xy, xyxy^2x, y^2xy^2, xy^5x, xy, x^3, \\ xyx^3, yx^3, y^2xy^2, y^2xyx, yxy^2, yxy, y^2, y^4x, x, y, \dots, \end{aligned} \quad (3.15)$$

which has period 18 and the basic period 9 since $x\theta = x^3$ and $y\theta = x^3yx$ where θ is a outer automorphism of order 2.

- (ii') If $k = 3$, we have the sequence for the generating pair (x, y) ,

$$\begin{aligned} x, y, xy, (xy)^2, x, y^2, y^5xy, (xy)^2, x, y, (xy)^3, (xy)^4, x^3, \\ y^2, xy^2, (yx)^2, x^3, y, yxy^2, (yx)^2, x^3, y^2, y^4x, e, x, y, xy, \dots, \end{aligned} \quad (3.16)$$

which has period 24 and the basic period 24 since $x\theta = x$ and $y\theta = y$ where θ is an inner automorphism induced by conjugation by x^2 .

If $k = 4$, we have the sequence for the generating pair (x, y) ,

$$\begin{aligned} x, y, xy, (xy)^2, (xy)^4, x^3, y^2, y^5xy, (xy)^2, e, x, \\ y, (xy)^3, (xy)^4, e, x^3, y^2, xy^2, (yx)^2, x^2, x, y, \\ yxy^2, (yx)^2, e, x^3, y^2, y^4x, e, e, x, y, xy, (xy)^2, \dots, \end{aligned} \quad (3.17)$$

which has period 30 and the basic period 30 since $x\theta = x$ and $y\theta = y$ where θ is an inner automorphism induced by conjugation by x^2 .

If $k \geq 5$, the first k elements of sequence for the generating pair (x, y) are

$$x_0 = x, x_1 = y, x_2 = xy, x_3 = (xy)^2, x_4 = (xy)^4, x_5 = (xy)^8 \dots, x_{k-1} = (xy)^{2^{k-3}}. \quad (3.18)$$

Thus, using the above information, sequence reduces to

$$x_0 = x, x_1 = y, x_2 = xy, x_3 = (xy)^2, x_4 = (xy)^4, x_5 = e, \dots, e, x_{k-1} = e, \quad (3.19)$$

where $x_j = e$ for $5 \leq j \leq k-1$. Thus,

$$\begin{aligned} x_k &= e, x_{k+1} = x^3, x_{k+2} = y^2, x_{k+3} = y^5 xy, \\ x_{k+4} &= (xy)^2, x_{k+5} = e, \dots, e, x_{2k+1} = e, x_{2k+2} = x, \end{aligned} \quad (3.20)$$

$$\begin{aligned} x_{2k+3} &= y, x_{2k+4} = (xy)^3, x_{2k+5} = (xy)^4, x_{2k+6} = e, \dots, e, \\ x_{3k+2} &= e, x_{3k+3} = x^3, x_{3k+4} = y^2, x_{3k+5} = xy^2, \\ x_{3k+6} &= (yx)^2, x_{3k+7} = x^2, x_{3k+8} = e \dots, e, x_{4k+3} = e, \\ x_{4k+4} &= x, x_{4k+5} = y, x_{4k+6} = yxy^2 x_{4k+7} = (yx)^2, \end{aligned} \quad (3.21)$$

$$\begin{aligned} x_{4k+8} &= e, \dots, e, x_{5k+4} = e, x_{5k+5} = x^3, x_{5k+6} = y^2, \\ x_{5k+7} &= y^4 x, x_{5k+8} = e, \dots, e, x_{6k+5} = e, \end{aligned}$$

where $x_j = e$ for $k+5 \leq j \leq 2k+1, 2k+6 \leq j \leq 3k+2, 3k+8 \leq j \leq 4k+3, 4k+8 \leq j \leq 5k+4$, and $5k+8 \leq j \leq 6k+5$.

We also have

$$x_{6k+6} = \prod_{i=5k+6}^{6k+5} x_i = x, \quad x_{6k+7} = \prod_{i=5k+7}^{6k+6} x_i = y. \quad (3.22)$$

Since the elements succeeding x_{6k+6}, x_{6k+7} depend on x and y for their values, the cycle begins again with the $6k+6$ th element, that is, $x_0 = x_{6k+6}, x_1 = x_{6k+7}, \dots$. Thus, $P_k(\langle 2, 3, 4 \rangle; x, y) = 6k+6$ and $BP_k(\langle 2, 3, 4 \rangle; x, y) = 6k+6$ since $x\theta = x$ and $y\theta = y$ where θ is an inner automorphism induced by conjugation by x^2 .

Secondly, let us consider the 3-generator case. We first note that $|x| = 4, |y| = 6$, and $|z| = 8$.

- (i) If $k = 2, P_2(\langle 2, 3, 4 \rangle; y, z, x) = 18$ and $BP_2(\langle 2, 3, 4 \rangle; y, z, x) = 9$ since $x\theta = x^3, y\theta = x^3 yx$, and $z\theta = xy^2$ where θ is an outer automorphism of order 2.
- (ii) If $k > 2, P_k(\langle 2, 3, 4 \rangle; x, y, z) = 6k+6$ and $BP_k(\langle 2, 3, 4 \rangle; x, y, z) = 6k+6$ since $x\theta = x$ and $y\theta = y$ where θ is an inner automorphism induced by conjugation by x^2 .

The proofs are similar to the proofs of Theorems 3.5.(i) and 3.5.(ii) and are omitted. $\square$

Theorem 3.7. *The periods of the k -nacci sequences and the basic periods of the basic k -nacci sequences in the group A_4 are as follows.*

If the group is defined by the presentation $A_4 = \langle x, y, z : x^2 = y^3 = z^3 = xyz = e \rangle$, then

(i) *if $k = 2$, $P_2(A_4; y, z, x) = 16$ and $BP_2(A_4; y, z, x) = 4$,*

(ii) *if $k > 2$,*

$$P_k(A_4; x, y, z) = \begin{cases} 3BP_k(A_4; x, y, z), & k \equiv 0 \pmod{4}, \\ 2BP_k(A_4; x, y, z), & k \equiv 2 \pmod{4}, \\ 2BP_k(A_4; x, y, z), & \text{otherwise}, \end{cases} \quad (3.23)$$

$$BP_k(A_4; x, y, z) = \begin{cases} u_1 h_k(3), & k \equiv 0 \pmod{4}, \\ u_2 h_k(3), & k \equiv 2 \pmod{4}, \\ u_3 h_k(3), & \text{otherwise}, \end{cases}$$

where $u_1, u_2, u_3 \in \mathbb{N}$, and $h_k(3)$ denote the wall number of the k -step Fibonacci sequence modulo 3.

If the group is defined by the presentation $A_4 = \langle x, y : x^2 = y^3 = (xy)^3 = e \rangle$, then

(i') *if $k = 2$, $P_2(A_4; x, y) = 16$ and $BP_2(A_4; x, y) = 4$,*

(ii') *if $k > 2$,*

$$P_k(A_4; x, y) = \begin{cases} 3BP_k(A_4; x, y), & k \equiv 0 \pmod{4}, \\ 2BP_k(A_4; x, y), & k \equiv 2 \pmod{4}, \\ 2BP_k(A_4; x, y), & \text{otherwise}, \end{cases} \quad (3.24)$$

$$BP_k(A_4; x, y) = \begin{cases} u_1 h_k(3), & k \equiv 0 \pmod{4}, \\ u_2 h_k(3), & k \equiv 2 \pmod{4}, \\ u_3 h_k(3), & \text{otherwise}, \end{cases}$$

where $u_1, u_2, u_3 \in \mathbb{N}$.

Proof. Firstly, let us consider the 2-generator case. We process as similar to the proof of Theorem 3.6 We first note that $|x| = 2$, $|y| = 3$, and $|xy| = 3$.

(i') If $k = 2$, we have the sequence for the generating pair (x, y) ,

$$\begin{aligned} & x, y, xy, yxy, yxy^2, (xy)^2, xy^2, y, x, \\ & yx, xyx, y^2x, yxy^2, yxy, y^2, yx, x, y, \dots, \end{aligned} \quad (3.25)$$

which has period 16 and the basic period 4 since $x\theta = yxy^2$ and $y\theta = yxy$ where θ is an outer automorphism of order 4.

(ii') If $k > 2$,

let k be even, then the first k elements of sequence for the generating pair (x, y) are

$$x_0 = x, x_1 = y, x_2 = xy, x_3 = (xy)^2, x_4 = xy, x_5 = (xy)^2, \dots, x_{k-2} = xy, x_{k-1} = (xy)^2. \quad (3.26)$$

If $k \equiv 0 \pmod{4}$,

$$\begin{aligned} x_{u_1 h_k(3)-(k-2)} &= e, \quad x_{u_1 h_k(3)-(k-1)} = e, \dots, e, \\ x_{u_1 h_k(3)-1} &= e, \quad x_{u_1 h_k(3)} = y^2 xy, \quad x_{u_1 h_k(3)+1} = yx, \dots \end{aligned} \quad (3.27)$$

$P_k(A_4; x, y) = 3BP_k(A_4; x, y)$ and $BP_k(A_4; x, y) = u_1 h_k(3)$ since $x\theta = yxy^2$ and $y\theta = xyx$ where θ is an outer automorphism of order 3.

If $k \equiv 2 \pmod{4}$,

$$\begin{aligned} x_{u_2 h_k(3)-(k-2)} &= e, \quad x_{u_2 h_k(3)-(k-1)} = e, \dots, e, \\ x_{u_2 h_k(3)-1} &= e, \quad x_{u_2 h_k(3)} = x, \quad x_{u_2 h_k(3)+1} = xy, \dots \end{aligned} \quad (3.28)$$

$P_k(A_4; x, y) = 2BP_k(A_4; x, y)$ and $BP_k(A_4; x, y) = u_2 h_k(3)$ since $x\theta = x$ and $y\theta = xy$ where θ is a outer automorphism of order 2.

Let k be odd, then the first k elements of sequence are for the generating pair (x, y) ,

$$x_0 = x, x_1 = y, x_2 = xy, x_3 = (xy)^2, x_4 = xy, x_5 = (xy)^2, \dots, x_{k-2} = (xy)^2, x_{k-1} = xy. \quad (3.29)$$

Also,

$$\begin{aligned} x_{u_3 h_k(3)-(k-2)} &= e, \quad x_{u_3 h_k(3)-(k-1)} = e, \dots, e, \\ x_{u_3 h_k(3)-1} &= e, \quad x_{u_3 h_k(3)} = x, \quad x_{u_3 h_k(3)+1} = yx, \dots \end{aligned} \quad (3.30)$$

$P_k(A_4; x, y) = 2BP_k(A_4; x, y)$ and $BP_k(A_4; x, y) = u_3 h_k(3)$ since $x\theta = x$ and $y\theta = yx$ where θ is an outer automorphism of order 2.

Secondly, let us consider the 3-generator case. We first note that $|x| = 2$, $|y| = 3$, and $|z| = 3$.

(i) If $k = 2$, $P_2(A_4; y, z, x) = 16$ and $BP_2(A_4; y, z, x) = 4$ since $x\theta = y^2 xy$, $y\theta = yxy$, and $z\theta = yx$ where θ is an outer automorphism of order 4.

(ii) If $k > 2$,

let $k \equiv 0 \pmod{4}$, then $P_k(A_4; x, y, z) = 3BP_k(A_4; x, y, z)$ and $BP_k(A_4; x, y, z) = u_1 h_k(3)$ since $x\theta = y^2 xy$, $y\theta = xyx$, and $z\theta = zx$ where θ is an outer

automorphism of order 3; let $k \equiv 2 \pmod{4}$, then $P_k(A_4; x, y, z) = 2BP_k(A_4; x, y, z)$ and $BP_k(A_4; x, y, z) = u_2 h_k(3)$ since $x\theta = x$, $y\theta = yx$, and $z\theta = yz^2$ where θ is an outer automorphism of order 2; let k be odd; then $P_k(A_4; x, y, z) = 2BP_k(A_4; x, y, z)$ and $BP_k(A_4; x, y, z) = u_3 h_k(3)$ since $x\theta = x$, $y\theta = xy$, and $z\theta = zx$ where θ is an outer automorphism of order 2.

The proofs are similar to the proofs of Theorems 3.5.(i) and 3.5.(i.i) and are omitted. $\square$

Theorem 3.8. *The periods of the k -nacci sequences and the basic periods of the basic k -nacci sequences in the binary polyhedral group $\langle 2, 3, 3 \rangle$ are as follows.*

If the group is defined by the presentation $\langle 2, 3, 3 \rangle = \langle x, y, z : x^2 = y^3 = z^3 = xyz \rangle$, then

- (i) if $k = 2$, $P_2(\langle 2, 3, 3 \rangle; y, z, x) = 48$ and $BP_2(\langle 2, 3, 3 \rangle; y, z, x) = 12$,
- (ii) if $k > 2$,

$$P_k(\langle 2, 3, 3 \rangle; x, y, z) = \begin{cases} 3BP_k(\langle 2, 3, 3 \rangle; x, y, z), & k \equiv 0 \pmod{4}, \\ BP_k(\langle 2, 3, 3 \rangle; x, y, z), & k \not\equiv 0 \pmod{4}, \end{cases} \quad (3.31)$$

$$BP_k(\langle 2, 3, 3 \rangle; x, y, z) = \begin{cases} v_1 h_k(6), & k \equiv 0 \pmod{4}, \\ v_2 h_k(6), & k \not\equiv 0 \pmod{4}, \end{cases} \quad (3.32)$$

where $v_1, v_2 \in \mathbb{N}$, and $h_k(6)$ denote the wall number of the k -step Fibonacci sequence modulo 6.

If the group is defined by the presentation $\langle 2, 3, 3 \rangle = \langle x, y : x^2 = y^3 = (xy)^3 \rangle$, then

- (i') if $k = 2$, $P_2(\langle 2, 3, 3 \rangle; x, y) = 48$ and $BP_2(\langle 2, 3, 3 \rangle; x, y) = 12$,
- (ii') if $k > 2$,

$$P_k(\langle 2, 3, 3 \rangle; x, y) = \begin{cases} 3BP_k(\langle 2, 3, 3 \rangle; x, y), & k \equiv 0 \pmod{4}, \\ BP_k(\langle 2, 3, 3 \rangle; x, y), & k \not\equiv 0 \pmod{4}, \end{cases} \quad (3.33)$$

$$BP_k(\langle 2, 3, 3 \rangle; x, y) = \begin{cases} v_1 h_k(6), & k \equiv 0 \pmod{4}, \\ v_2 h_k(6), & k \not\equiv 0 \pmod{4}, \end{cases} \quad (3.34)$$

where $v_1, v_2 \in \mathbb{N}$.

Proof. Firstly, let us consider the 3-generator case. We first note that $|x| = 4$, $|y| = 6$, and $|z| = 6$.

- (i) If $k = 2$, $P_2(\langle 2, 3, 3 \rangle; y, z, x) = 48$ and $BP_2(\langle 2, 3, 3 \rangle; y, z, x) = 12$ since $x\theta = y^2xy$, $y\theta = xz^4x$, and $z\theta = y^2xy^2$ where θ is an outer automorphism of order 4.

(ii) If $k > 2$,

let $k \equiv 0 \pmod{4}$, then $P_k(\langle 2, 3, 3 \rangle; x, y, z) = 3BP_k(\langle 2, 3, 3 \rangle; x, y, z)$ and $BP_k(\langle 2, 3, 3 \rangle; x, y, z) = v_1 h_k(6)$ since $x\theta = yxy^5$, $y\theta = z^3xy$, and $z\theta = xy^2x$ where θ is an inner automorphism induced by conjugation by z^3yx ;

let $k \not\equiv 0 \pmod{4}$, then $P_k(\langle 2, 3, 3 \rangle; x, y, z) = BP_k(\langle 2, 3, 3 \rangle; x, y, z)$ and $BP_k(\langle 2, 3, 3 \rangle; x, y, z) = v_2 h_k(6)$ since $x\theta = x$, $y\theta = y$, and $z\theta = z$ where θ is an inner automorphism induced by conjugation by x^2 .

The proofs are similar to the proofs of Theorems 3.5.(i) and 3.5.(ii) and are omitted.

Secondly, let us consider the 2-generator case. We first note that $|x| = 4$, $|y| = 6$, and $|xy| = 6$.

(i') If $k = 2$, $P_2(\langle 2, 3, 3 \rangle; x, y) = 48$ and $BP_2(\langle 2, 3, 3 \rangle; x, y) = 12$ since $x\theta = yxy^2$ and $y\theta = y^2x$ where θ is an outer automorphism of order 4.

(ii') If $k > 2$,

let $k \equiv 0 \pmod{4}$, then $P_k(\langle 2, 3, 3 \rangle; x, y) = 3BP_k(\langle 2, 3, 3 \rangle; x, y)$ and $BP_k(\langle 2, 3, 3 \rangle; x, y) = v_1 h_k(6)$ since $x\theta = y^5xy$, $y\theta = yx$, and $z\theta = xy^2x$ where θ is an inner automorphism induced by conjugation by y^5x ,

let $k \not\equiv 0 \pmod{4}$, then $P_k(\langle 2, 3, 3 \rangle; x, y) = BP_k(\langle 2, 3, 3 \rangle; x, y)$ and $BP_k(\langle 2, 3, 3 \rangle; x, y) = v_2 h_k(6)$ since $x\theta = x$ and $y\theta = y$ where θ is an inner automorphism induced by conjugation by x^2 .

The proofs are similar to the proofs of Theorem 3.6.(i') and Theorem 3.6.(ii') and are omitted. $\square$

Theorem 3.9. *The periods of the k -nacci sequences are $k + 1$, and the basic period of the basic k -nacci sequences is $k + 1$ in D_2 four-group.*

Proof. We have the presentation $D_2 = \langle x, y : x^2 = y^2 = e, xy = yx \rangle$. $P_k(D_2; x, y) = k + 1$; see [14] for a proof and $BP_k(D_2; x, y) = k + 1$ since $x\theta = x$ and $y\theta = y$ where θ is an inner automorphism induced by conjugation by x . $\square$

Acknowledgments

The authors thank the referees for their valuable suggestions which improved the presentation of the paper. This paper was supported by the Commission for the Scientific Research Projects of Kafkas University, Project no. 2010-FEF-61.

References

- [1] D. D. Wall, "Fibonacci series modulo m ," *The American Mathematical Monthly*, vol. 67, pp. 525–532, 1960.
- [2] H. J. Wilcox, "Fibonacci sequences of period n in groups," *The Fibonacci Quarterly*, vol. 24, no. 4, pp. 356–361, 1986.
- [3] C. M. Campbell, H. Doostie, and E. F. Robertson, "Fibonacci length of generating pairs in groups," in *Applications of Fibonacci Numbers, Vol. 3*, pp. 27–35, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1990.
- [4] C. M. Campbell and P. P. Campbell, "The Fibonacci lengths of binary polyhedral groups and related groups," *Congressus Numerantium*, vol. 194, pp. 95–102, 2009.

- [5] C. M. Campbell and P. P. Campbell, "The Fibonacci length of certain centro-polyhedral groups," *Journal of Applied Mathematics & Computing*, vol. 19, no. 1-2, pp. 231–240, 2005.
- [6] H. Aydın and R. Dikici, "General Fibonacci sequences in finite groups," *The Fibonacci Quarterly*, vol. 36, no. 3, pp. 216–221, 1998.
- [7] H. Aydın and G. C. Smith, "Finite p -quotients of some cyclically presented groups," *Journal of the London Mathematical Society. Second Series*, vol. 49, no. 1, pp. 83–92, 1994.
- [8] H. Doostie and C. M. Campbell, "Fibonacci length of automorphism groups involving Tribonacci numbers," *Vietnam Journal of Mathematics*, vol. 28, no. 1, pp. 57–65, 2000.
- [9] H. Doostie and M. Hashemi, "Fibonacci lengths involving the Wall number $k(n)$," *Journal of Applied Mathematics & Computing*, vol. 20, no. 1-2, pp. 171–180, 2006.
- [10] E. Özkan, "On truncated Fibonacci sequences," *Indian Journal of Pure and Applied Mathematics*, vol. 38, no. 4, pp. 241–251, 2007.
- [11] S. W. Knox, "Fibonacci sequences in finite groups," *The Fibonacci Quarterly*, vol. 30, no. 2, pp. 116–120, 1992.
- [12] Ö. Deveci, E. Karaduman, and C. M. Campbell, "On The k -nacci sequences in finite binary polyhedral groups," to appear in *Algebra Colloquium*.
- [13] E. Karaduman and H. Aydın, " k -nacci sequences in some special groups of finite order," *Mathematical and Computer Modelling*, vol. 50, no. 1-2, pp. 53–58, 2009.
- [14] E. Karaduman and Ö. Deveci, " k -nacci sequences in finite triangle groups," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 453750, 10 pages, 2009.
- [15] P. Hall, "The Eulerian functions of a group," *The Quarterly Journal of Mathematics*, vol. 7, pp. 134–151, 1936.
- [16] J. H. Conway, H. S. M. Coxeter, and G. C. Shephard, "The centre of a finitely generated group," *The Tensor Society*, vol. 25, pp. 405–418, 1972, Erratum in *IBID Journal*, vol. 26, pp. 477, 1972.
- [17] H. S. M. Coxeter and W. O. J. Moser, *Generators and Relations for Discrete Groups*, Springer, New York, NY, USA, 3rd edition, 1972.
- [18] K. Lü and W. Jun, " k -step Fibonacci sequence modulo m ," *Utilitas Mathematica*, vol. 71, pp. 169–177, 2006.