Research Article

Generalized Systems of Variational Inequalities and Projection Methods for Inverse-Strongly Monotone Mappings

Wiyada Kumam,^{1,2,3} Prapairat Junlouchai,¹ and Poom Kumam^{2,3}

¹ Department of Mathematics, Faculty of Science and Technology, Rajamangala University of Technology Thanyaburi (RMUTT), Thanyaburi, Pathumthani 12110, Thailand

² Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi (KMUTT), Bangmod, Bangkok 10140, Thailand

³ Centre of Excellence in Mathematics, CHE, Si Ayutthaya Road, Bangkok 10400, Thailand

Correspondence should be addressed to Poom Kumam, poom.kum@kmutt.ac.th

Received 20 February 2011; Accepted 3 May 2011

Academic Editor: Jianshe Yu

Copyright © 2011 Wiyada Kumam et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce an iterative sequence for finding a common element of the set of fixed points of a nonexpansive mapping and the solutions of the variational inequality problem for three inverse-strongly monotone mappings. Under suitable conditions, some strong convergence theorems for approximating a common element of the above two sets are obtained. Moreover, using the above theorem, we also apply to find solutions of a general system of variational inequality and a zero of a maximal monotone operator in a real Hilbert space. As applications, at the end of the paper we utilize our results to study some convergence problem for strictly pseudocontractive mappings. Our results include the previous results as special cases extend and improve the results of Ceng et al., (2008) and many others.

1. Introduction

Variational inequalities are known to play a crucial role in mathematics as a unified framework for studying a large variety of problems arising, for instance, in structural analysis, engineering sciences and others. Roughly speaking, they can be recast as fixed-point problems, and most of the numerical methods related to this topic are based on projection methods. Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$, and let E be a nonempty, closed, convex subset of H . A mapping $A : E \rightarrow H$ is called α -inverse-strongly monotone if there exists a positive real number $\alpha > 0$ such that

$$\langle Ax - Ay, x - y \rangle \geq \alpha \|Ax - Ay\|^2, \quad \forall x, y \in E \quad (1.1)$$

(see [1, 2]). It is obvious that every α -inverse-strongly monotone mapping A is monotone and Lipschitz continuous. A mapping $S : E \rightarrow E$ is called nonexpansive if

$$\|Sx - Sy\| \leq \|x - y\|, \quad \forall x, y \in E. \quad (1.2)$$

We denote by $F(S)$ the set of fixed points of S and by P_E the metric projection of H onto E . Recall that the classical variational inequality, denoted by $VI(A, E)$, is to find an $x^* \in E$ such that

$$\langle Ax^*, x - x^* \rangle \geq 0, \quad \forall x \in E. \quad (1.3)$$

The set of solutions of $VI(A, E)$ is denoted by Γ . The variational inequality has been widely studied in the literature; see, for example, [3–6] and the references therein.

For finding an element of $F(S) \cap \Gamma$, Takahashi and Toyoda [7] introduced the following iterative scheme:

$$x_{n+1} = \alpha_n x_n + (1 - \alpha_n) SP_E(x_n - \lambda_n Ax_n), \quad (1.4)$$

for every $n = 0, 1, 2, \dots$, where $x_0 = x \in E$, $\{\alpha_n\}$ is a sequence in $(0, 1)$, and $\{\lambda_n\}$ is a sequence in $(0, 2\alpha)$. On the other hand, for solving the variational inequality problem in the finite-dimensional Euclidean space $\mathbf{R}^n$, Korpelevich (1976) [8] introduced the following so-called extragradient method:

$$\begin{aligned} x_0 &= x \in E, \\ y_n &= P_E(x_n - \lambda_n Ax_n), \\ x_{n+1} &= P_E(x_n - \lambda_n Ay_n), \end{aligned} \quad (1.5)$$

for every $n = 0, 1, 2, \dots$, where $\lambda_n \in (0, 1/k)$. Many authors using extragradient method for approximating common fixed points and variational inequality problems (see also [9, 10]). Recently, Nadezhkina and Takahashi [11] and Zeng et al. [12] proposed some iterative schemes for finding elements in $F(S) \cap \Gamma$ by combining (1.4) and (1.5). Further, these iterative schemes are extended in Y. Yao and J. C. Yao [13] to develop a new iterative scheme for finding elements in $F(S) \cap \Gamma$.

Consider the following problem of finding $(x^*, y^*) \in E \times E$ such that (see cf. Ceng et al. [14]):

$$\begin{aligned} \langle \lambda Ay^* + x^* - y^*, x - x^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \mu Bx^* + y^* - x^*, x - y^* \rangle &\geq 0, \quad \forall x \in E, \end{aligned} \quad (1.6)$$

which is called *general system of variational inequalities* (GSVI), where $\lambda > 0$ and $\mu > 0$ are two constants. In particular, if $A = B$, then problem (1.6) reduces to finding $(x^*, y^*) \in E \times E$ such that

$$\begin{aligned} \langle \lambda Ay^* + x^* - y^*, x - x^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \mu Ax^* + y^* - x^*, x - y^* \rangle &\geq 0, \quad \forall x \in E, \end{aligned} \quad (1.7)$$

which is defined by [15, 16], and is called the new system of variational inequalities. Further, if $x^* = y^*$, then problem (1.7) reduces to the *classical variational inequality* $VI(A, E)$, that is, find $x^* \in E$ such that $\langle Ax^*, x - x^* \rangle \geq 0$, for all $x \in E$.

We can characterize problem, if $x^* \in F(S) \cap VI(A, E)$, then it follows that $x^* = Sx^* = P_E[x^* - \rho Ax^*]$, where $\rho > 0$ is a constant.

In 2008, Ceng et al. [14] introduced a relaxed extragradient method for finding solutions of problem (1.6). Let the mappings $A, B : E \rightarrow H$ be α -inverse-strongly monotone and β -inverse-strongly monotone, respectively. Let $S : E \rightarrow E$ be a nonexpansive mapping. Suppose $x_1 = u \in E$ and $\{x_n\}$ is generated by

$$\begin{aligned} y_n &= P_E(x_n - \mu Bx_n), \\ x_{n+1} &= \alpha_n u + \beta_n x_n + \gamma_n SP_E(y_n - \lambda_n Ay_n), \end{aligned} \quad (1.8)$$

where $\lambda \in (0, 2\alpha)$, $\mu \in (0, 2\beta)$, and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ are three sequences in $[0, 1]$ such that $\alpha_n + \beta_n + \gamma_n = 1$, for all $n \geq 1$. First, problem (1.6) is proven to be equivalent to a fixed point problem of a nonexpansive mapping.

In this paper, motivated by what is mentioned above, we consider generalized system of variational inequalities as follows.

Let E be a nonempty, closed, convex subset of a real Hilbert space H . Let $A, B, C : E \rightarrow H$ be three mappings. We consider the following problem of finding $(x^*, y^*, z^*) \in E \times E \times E$ such that

$$\begin{aligned} \langle \lambda Ay^* + x^* - y^*, x - x^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \mu Bz^* + y^* - z^*, x - y^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \tau Cx^* + z^* - x^*, x - z^* \rangle &\geq 0, \quad \forall x \in E, \end{aligned} \quad (1.9)$$

which is called a *general system of variational inequalities* where $\lambda > 0$, $\mu > 0$ and $\tau > 0$ are three constants.

In particular, if $A = B = C$, then problem (1.9) reduces to finding $(x^*, y^*, z^*) \in E \times E \times E$ such that

$$\begin{aligned} \langle \lambda Ay^* + x^* - y^*, x - x^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \mu Az^* + y^* - z^*, x - y^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \tau Ax^* + z^* - x^*, x - z^* \rangle &\geq 0, \quad \forall x \in E. \end{aligned} \quad (1.10)$$

Next, we consider some special classes of the GSVI problem (1.9) reduce to the following GSVI.

- (i) If $\tau = 0$, then the GSVI problems (1.9) reduce to GSVI problem (1.6).
- (ii) If $\tau = \mu = 0$, then the GSVI problems (1.9) reduce to classical variational inequality $VI(A, E)$ problem.

The above system enters a class of more general problems which originated mainly from the Nash equilibrium points and was treated from a theoretical viewpoint in [17, 18]. Observe at the same time that, to construct a mathematical model which is as close as possible to a real complex problem, we often have to use constraints which can be expressed as one several subproblems of a general problem. These constraints can be given, for instance, by variational inequalities, by fixed point problems, or by problems of different types.

This paper deals with a relaxed extragradient approximation method for solving a system of variational inequalities over the fixed-point sets of nonexpansive mapping. Under classical conditions, we prove a strong convergence theorem for this method. Moreover, the proposed algorithm can be applied for instance to solving the classical variational inequality problems.

2. Preliminaries

Let E be a nonempty, closed, convex subset of a real Hilbert space H . For every point $x \in H$, there exists a unique nearest point in E , denoted by $P_E x$, such that

$$\|x - P_E x\| \leq \|x - y\|, \quad \forall y \in E. \quad (2.1)$$

P_E is call the metric projection of H onto E .

Recall that, $P_E x$ is characterized by following properties: $P_E x \in E$ and

$$\begin{aligned} \langle x - P_E x, y - P_E x \rangle &\leq 0, \\ \|x - y\|^2 &\geq \|x - P_E x\|^2 + \|P_E x - y\|^2, \end{aligned} \quad (2.2)$$

for all $x \in H$ and $y \in E$.

Lemma 2.1 (see cf. Zhang et al. [19]). *The metric projection P_E has the following properties:*

- (i) $P_E : H \rightarrow E$ is nonexpansive;
- (ii) $P_E : H \rightarrow E$ is firmly nonexpansive, that is,

$$\|P_E x - P_E y\|^2 \leq \langle P_E x - P_E y, x - y \rangle, \quad \forall x, y \in H; \quad (2.3)$$

- (iii) for each $x \in H$,

$$z = P_E(x) \iff \langle x - z, z - y \rangle \geq 0, \quad \forall y \in E. \quad (2.4)$$

Lemma 2.2 (see Osilike and Igbokwe [20]). *Let $(E, \langle \cdot, \cdot \rangle)$ be an inner product space. Then for all $x, y, z \in E$ and $\alpha, \beta, \gamma \in [0, 1]$ with $\alpha + \beta + \gamma = 1$, one has*

$$\|\alpha x + \beta y + \gamma z\|^2 = \alpha\|x\|^2 + \beta\|y\|^2 + \gamma\|z\|^2 - \alpha\beta\|x - y\|^2 - \alpha\gamma\|x - z\|^2 - \beta\gamma\|y - z\|^2. \quad (2.5)$$

Lemma 2.3 (see Suzuki [21]). *Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space X and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$. Suppose $x_{n+1} = (1 - \beta_n)y_n + \beta_n x_n$ for all integers $n \geq 0$ and $\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$. Then, $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.*

Lemma 2.4 (see Xu [22]). *Assume $\{a_n\}$ is a sequence of nonnegative real numbers such that*

$$a_{n+1} \leq (1 - \alpha_n)a_n + \delta_n, \quad n \geq 0, \quad (2.6)$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence in $\mathbf{R}$ such that

- (i) $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (ii) $\limsup_{n \rightarrow \infty} (\delta_n / \alpha_n) \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n| < \infty$.

Then, $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 2.5 (Goebel and Kirk [23]). *Demiclosedness Principle. Assume that T is a nonexpansive self-mapping of a nonempty, closed, convex subset E of a real Hilbert space H . If T has a fixed point, then $I - T$ is demiclosed; that is, whenever $\{x_n\}$ is a sequence in E converging weakly to some $x \in E$ (for short, $x_n \rightharpoonup x \in E$), and the sequence $\{(I - T)x_n\}$ converges strongly to some y (for short, $(I - T)x_n \rightarrow y$), it follows that $(I - T)x = y$. Here, I is the identity operator of H .*

The following lemma is an immediate consequence of an inner product.

Lemma 2.6. *In a real Hilbert space H , there holds the inequality*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle, \quad \forall x, y \in H. \quad (2.7)$$

Remark 2.7. We also have that, for all $u, v \in E$ and $\lambda > 0$,

$$\begin{aligned} \|(I - \lambda A)u - (I - \lambda A)v\|^2 &= \|(u - v) - \lambda(Au - Av)\|^2 \\ &= \|u - v\|^2 - 2\lambda\langle u - v, Au - Av \rangle \\ &\quad + \lambda^2\|Au - Av\|^2 \\ &\leq \|u - v\|^2 + \lambda(\lambda - 2\alpha)\|Au - Av\|^2. \end{aligned} \quad (2.8)$$

So, if $\lambda \leq 2\alpha$, then $I - \lambda A$ is a nonexpansive mapping from E to H .

3. Main Results

In this section, we introduce an iterative process by the relaxed extragradient approximation method for finding a common element of the set of fixed points of a nonexpansive mapping and the solution set of the variational inequality problem for three inverse-strongly monotone mappings in a real Hilbert space. We prove that the iterative sequence converges strongly to a common element of the above two sets.

In order to prove our main result, the following lemmas are needed.

Lemma 3.1. *For given $x^*, y^*, z^* \in E \times E \times E$, (x^*, y^*, z^*) is a solution of problem (1.9) if and only if x^* is a fixed point of the mapping $G : E \rightarrow E$ defined by*

$$G(x) = P_E \{ P_E [P_E(x - \tau Cx) - \mu B P_E(x - \tau Cx)] - \lambda A P_E [P_E(x - \tau Cx) - \mu B P_E(x - \tau Cx)] \}, \quad \forall x \in E, \quad (3.1)$$

where $y^* = P_E(z^* - \mu B z^*)$ and $z^* = P_E(x^* - \tau C x^*)$.

Proof.

$$\begin{aligned} \langle \lambda A y^* + x^* - y^*, x - x^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \mu B z^* + y^* - z^*, x - y^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle \tau C x^* + z^* - x^*, x - z^* \rangle &\geq 0, \quad \forall x \in E, \end{aligned} \quad (3.2)$$

$\Leftrightarrow$

$$\begin{aligned} \langle (-y^* + \lambda A y^*) + x^*, x - x^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle (-z^* + \mu B z^*) + y^*, x - y^* \rangle &\geq 0, \quad \forall x \in E, \\ \langle (-x^* + \tau C x^*) + z^*, x - z^* \rangle &\geq 0, \quad \forall x \in E, \end{aligned} \quad (3.3)$$

$\Leftrightarrow$

$$\begin{aligned} \langle (y^* - \lambda A y^*) - x^*, x^* - x \rangle &\geq 0, \quad \forall x \in E, \\ \langle (z^* - \mu B z^*) - y^*, y^* - x \rangle &\geq 0, \quad \forall x \in E, \\ \langle (x^* - \tau C x^*) - z^*, z^* - x \rangle &\geq 0, \quad \forall x \in E, \end{aligned} \quad (3.4)$$

$\Leftrightarrow$

$$\begin{aligned} x^* &= P_E(y^* - \lambda A y^*), \\ y^* &= P_E(z^* - \mu B z^*), \\ z^* &= P_E(x^* - \tau C x^*), \end{aligned} \quad (3.5)$$

$$\Leftrightarrow x^* = P_E[P_E(z^* - \mu B z^*) - \lambda A P_E(z^* - \mu B z^*)].$$

Thus,

$$x^* = P_E \{ P_E [P_E(x^* - \tau Cx^*) - \mu BP_E(x^* - \tau Cx^*)] - \lambda AP_E [P_E(x^* - \tau Cx^*) - \mu BP_E(x^* - \tau Cx^*)] \}. \quad (3.6)$$

□

Lemma 3.2. *The mapping G defined by Lemma 3.1 is nonexpansive mappings.*

Proof. For all $x, y \in E$,

$$\begin{aligned} \|G(x) - G(y)\| &= \|P_E \{ P_E [P_E(x - \tau Cx) - \mu BP_E(x - \tau Cx)] \\ &\quad - \lambda AP_E [P_E(x - \tau Cx) - \mu BP_E(x - \tau Cx)] \} \\ &\quad - P_E \{ P_E [P_E(y - \tau Cy) - \mu BP_E(y - \tau Cy)] \\ &\quad - \lambda AP_E [P_E(y - \tau Cy) - \mu BP_E(y - \tau Cy)] \} \| \\ &\leq \| [P_E(x - \tau Cx) - \mu BP_E(x - \tau Cx)] \\ &\quad - \lambda AP_E [P_E(x - \tau Cx) - \mu BP_E(x - \tau Cx)] \\ &\quad - [P_E(y - \tau Cy) - \mu BP_E(y - \tau Cy)] \\ &\quad - \lambda AP_E [P_E(y - \tau Cy) - \mu BP_E(y - \tau Cy)] \| \\ &= \| (I - \lambda A) [P_E(x - \tau Cx) - \mu BP_E(x - \tau Cx)] \\ &\quad - (I - \lambda A) [P_E(y - \tau Cy) - \mu BP_E(y - \tau Cy)] \| \\ &\leq \| [P_E(x - \tau Cx) - \mu BP_E(x - \tau Cx)] - [P_E(y - \tau Cy) - \mu BP_E(y - \tau Cy)] \| \\ &= \| (I - \mu B) [P_E(x - \tau Cx)] - (I - \mu B) [P_E(y - \tau Cy)] \| \\ &\leq \| P_E(x - \tau Cx) - P_E(y - \tau Cy) \| \\ &\leq \| (x - \tau Cx) - (y - \tau Cy) \| \\ &= \| (I - \tau C)(x) - (I - \tau C)(y) \| \\ &\leq \| x - y \|. \end{aligned} \quad (3.7)$$

This shows that $G : E \rightarrow E$ is a nonexpansive mapping. □

Throughout this paper, the set of fixed points of the mapping G is denoted by Υ .
Now, we are ready to proof our main results in this paper.

Theorem 3.3. Let E be a nonempty, closed, convex subset of a real Hilbert space H . Let the mapping $A, B, C : E \rightarrow H$ be α -inverse-strongly monotone, β -inverse-strongly monotone, and γ -inverse-strongly monotone, respectively. Let S be a nonexpansive mapping of E into itself such that $F(S) \cap \Upsilon \neq \emptyset$. Let f be a contraction of H into itself and given $x_1 \in H$ arbitrarily and $\{x_n\}$ is generated by

$$\begin{aligned} z_n &= P_E(x_n - \tau Cx_n), \\ y_n &= P_E(z_n - \mu Bz_n), \\ x_{n+1} &= \alpha_n f(x_n) + \beta_n x_n + \gamma_n SP_E(y_n - \lambda Ay_n), \quad n \geq 0, \end{aligned} \tag{3.8}$$

where $\lambda \in (0, 2\alpha)$, $\mu \in (0, 2\beta)$, $\tau \in (0, 2\gamma)$, and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ are three sequences in $[0, 1]$ such that

- (i) $\alpha_n + \beta_n + \gamma_n = 1$,
- (ii) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (iii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$.

Then, $\{x_n\}$ converges strongly to $\bar{x} \in F(S) \cap \Gamma$, where $\bar{x} = P_{F(S) \cap \Gamma} f(\bar{x})$ and $(\bar{x}, \bar{y}, \bar{z})$ is a solution of problem (1.9), where

$$\begin{aligned} \bar{y} &= P_E(\bar{z} - \mu B\bar{z}), \\ \bar{z} &= P_E(\bar{x} - \tau C\bar{x}). \end{aligned} \tag{3.9}$$

Proof. Let $x^* \in F(S) \cap \Gamma$. Then, $x^* = Sx^*$ and $x^* = Gx^*$, that is,

$$x^* = P_E\{P_E[P_E(x^* - \tau Cx^*) - \mu BP_E(x^* - \tau Cx^*)] - \lambda AP_E[P_E(x^* - \tau Cx^*) - \mu BP_E(x^* - \tau Cx^*)]\}. \tag{3.10}$$

Put $x^* = P_E(y^* - \lambda Ay^*)$ and $t_n = P_E(y_n - \lambda Ay_n)$. Then, $x^* = P_E[P_E(z^* - \mu Bz^*) - \lambda AP_E(z^* - \mu Bz^*)]$ implies that $y^* = P_E(z^* - \mu Bz^*)$, where $z^* = P_E(x^* - \tau Cx^*)$. Since $I - \lambda A$, $I - \mu B$ and $I - \tau C$ are nonexpansive mappings. We obtain that

$$\begin{aligned} \|t_n - x^*\| &= \|P_E(y_n - \lambda Ay_n) - x^*\| \\ &= \|P_E(y_n - \lambda Ay_n) - P_E(y^* - \lambda Ay^*)\| \\ &\leq \|(y_n - \lambda Ay_n) - (y^* - \lambda Ay^*)\| \\ &= \|(I - \lambda A)y_n - (I - \lambda A)y^*\| \\ &\leq \|y_n - y^*\| \\ &= \|y_n - P_E(z^* - \mu Bz^*)\| \\ &= \|P_E(z_n - \mu Bz_n) - P_E(z^* - \mu Bz^*)\| \end{aligned} \tag{3.11}$$

$$\begin{aligned}
&\leq \|(I - \mu B)z_n - (I - \mu B)z^*\| \\
&\leq \|z_n - z^*\|,
\end{aligned} \tag{3.12}$$

$$\begin{aligned}
\|z_n - z^*\| &= \|P_E(x_n - \tau Cx_n) - P_E(x^* - \tau Cx^*)\| \\
&\leq \|(x_n - \tau Cx_n) - (x^* - \tau Cx^*)\| \\
&= \|(I - \tau C)x_n - (I - \tau C)x^*\| \\
&\leq \|x_n - x^*\|.
\end{aligned} \tag{3.13}$$

Substituting (3.13) into (3.12), we have

$$\|t_n - x^*\| \leq \|x_n - x^*\|, \tag{3.14}$$

and by (3.11) we also have

$$\|y_n - y^*\| \leq \|x_n - x^*\|. \tag{3.15}$$

Since $x_{n+1} = \alpha_n f(x_n) + \beta_n x_n + \gamma_n St_n$ and by Lemma 2.2, we compute

$$\begin{aligned}
\|x_{n+1} - x^*\| &= \|\alpha_n f(x_n) + \beta_n x_n + \gamma_n St_n - x^*\| \\
&= \|\alpha_n (f(x_n) - x^*) + \beta_n (x_n - x^*) + \gamma_n (St_n - x^*)\| \\
&\leq \alpha_n \|f(x_n) - x^*\| + \beta_n \|x_n - x^*\| + \gamma_n \|St_n - x^*\| \\
&\leq \alpha_n \|f(x_n) - x^*\| + \beta_n \|x_n - x^*\| + \gamma_n \|t_n - x^*\| \\
&\leq \alpha_n \|f(x_n) - x^*\| + \beta_n \|x_n - x^*\| + \gamma_n \|x_n - x^*\| \\
&= \alpha_n \|f(x_n) - x^*\| + (1 - \alpha_n) \|x_n - x^*\| \\
&= \alpha_n \|f(x_n) - f(x^*) + f(x^*) - x^*\| + (1 - \alpha_n) \|x_n - x^*\| \\
&\leq \alpha_n \|f(x_n) - f(x^*)\| + \alpha_n \|f(x^*) - x^*\| + (1 - \alpha_n) \|x_n - x^*\| \\
&\leq \alpha_n k \|x_n - x^*\| + \alpha_n \|f(x^*) - x^*\| + (1 - \alpha_n) \|x_n - x^*\|
\end{aligned}$$

$$\begin{aligned}
&= (\alpha_n k + (1 - \alpha_n)) \|x_n - x^*\| + \alpha_n \|f(x^*) - x^*\| \\
&= (1 - \alpha_n(1 - k)) \|x_n - x^*\| + \alpha_n \|f(x^*) - x^*\| \\
&= (1 - \alpha_n(1 - k)) \|x_n - x^*\| + \alpha_n(1 - k) \frac{\|f(x^*) - x^*\|}{(1 - k)}.
\end{aligned} \tag{3.16}$$

By induction, we get

$$\|x_{n+1} - x^*\| \leq M, \tag{3.17}$$

where $M = \max\{\|x_0 - x^*\| + (1/(1 - k))\|f(x^*) - x^*\|\}$, $n \geq 0$. Therefore, $\{x_n\}$ is bounded. Consequently, by (3.11), (3.12) and (3.13), the sequences $\{t_n\}$, $\{St_n\}$, $\{y_n\}$, $\{Ay_n\}$, $\{z_n\}$, $\{Bz_n\}$, $\{Cx_n\}$, and $\{f(x_n)\}$ are also bounded. Also, we observe that

$$\begin{aligned}
\|z_{n+1} - z_n\| &= \|P_E(x_{n+1} - \tau Cx_{n+1}) - P_E(x_n - \tau Cx_n)\| \\
&\leq \|(I - \tau C)x_{n+1} - (I - \tau C)x_n\|
\end{aligned} \tag{3.18}$$

$$\leq \|x_{n+1} - x_n\|,$$

$$\begin{aligned}
\|t_{n+1} - t_n\| &= \|P_E(y_{n+1} - \lambda Ay_{n+1}) - P_E(y_n - \lambda Ay_n)\| \\
&\leq \|(y_{n+1} - \lambda Ay_{n+1}) - (y_n - \lambda Ay_n)\| \\
&= \|(I - \lambda A)y_{n+1} - (I - \lambda A)y_n\| \\
&\leq \|y_{n+1} - y_n\|
\end{aligned} \tag{3.19}$$

$$\begin{aligned}
&= \|P_E(z_{n+1} - \mu Bz_{n+1}) - P_E(z_n - \mu Bz_n)\| \\
&\leq \|z_{n+1} - z_n\| \\
&\leq \|x_{n+1} - x_n\|.
\end{aligned}$$

Let $x_{n+1} = (1 - \beta_n)w_n + \beta_n x_n$. Thus, we get

$$w_n = \frac{x_{n+1} - \beta_n x_n}{1 - \beta_n} = \frac{\alpha_n f(x_n) + \gamma_n SP_C(y_n - \lambda_n Ay_n)}{1 - \beta_n} = \frac{\alpha_n u + \gamma_n St_n}{1 - \beta_n} \tag{3.20}$$

It follows that

$$\begin{aligned}
w_{n+1} - w_n &= \frac{\alpha_{n+1}f(x_{n+1}) + \gamma_{n+1}St_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n f(x_n) + \gamma_n St_n}{1 - \beta_n} \\
&= \frac{\alpha_{n+1}f(x_{n+1})}{1 - \beta_{n+1}} + \frac{\gamma_{n+1}St_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_{n+1}f(x_n)}{1 - \beta_{n+1}} + \frac{\alpha_{n+1}f(x_n)}{1 - \beta_{n+1}} - \frac{\alpha_n f(x_n)}{1 - \beta_n} - \frac{\gamma_n St_n}{1 - \beta_n} \\
&= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(f(x_{n+1}) - f(x_n)) + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) f(x_n) + \frac{\gamma_{n+1}St_{n+1}}{1 - \beta_{n+1}} - \frac{\gamma_n St_n}{1 - \beta_n} \\
&= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(f(x_{n+1}) - f(x_n)) + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) f(x_n) \\
&\quad + \frac{\gamma_{n+1}St_{n+1}}{1 - \beta_{n+1}} - \frac{\gamma_{n+1}St_n}{1 - \beta_{n+1}} + \frac{\gamma_{n+1}St_n}{1 - \beta_{n+1}} - \frac{\gamma_n St_n}{1 - \beta_n} \\
&= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(f(x_{n+1}) - f(x_n)) + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) f(x_n) \\
&\quad + \frac{\gamma_{n+1}}{1 - \beta_{n+1}}(St_{n+1} - St_n) + \left(\frac{\gamma_{n+1}}{1 - \beta_{n+1}} - \frac{\gamma_n}{1 - \beta_n} \right) St_n \\
&= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(f(x_{n+1}) - f(x_n)) + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) f(x_n) \\
&\quad + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) St_n + \frac{\gamma_{n+1}}{1 - \beta_{n+1}}(St_{n+1} - St_n) \\
&= \frac{\alpha_{n+1}}{1 - \beta_{n+1}}(f(x_{n+1}) - f(x_n)) + \left(\frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right) (f(x_n) + St_n) \\
&\quad + \frac{\gamma_{n+1}}{1 - \beta_{n+1}}(St_{n+1} - St_n).
\end{aligned} \tag{3.21}$$

Combining (3.19) and (3.21), we obtain

$$\begin{aligned}
\|w_{n+1} - w_n\| - \|x_{n+1} - x_n\| &\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \right| \|f(x_{n+1}) - f(x_n)\| + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| \|f(x_n) + St_n\| \\
&\quad + \left| \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \right| \|St_{n+1} - St_n\| - \|x_{n+1} - x_n\| \\
&\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \right| k \|x_{n+1} - x_n\| + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| \|f(x_n) + St_n\| \\
&\quad + \left| \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \right| \|t_{n+1} - t_n\| - \|x_{n+1} - x_n\| \\
&\leq \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \right| k \|x_{n+1} - x_n\| + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| \|f(x_n) + St_n\|
\end{aligned}$$

$$\begin{aligned}
& + \left| \frac{\gamma_{n+1}}{1 - \beta_{n+1}} \right| \|x_{n+1} - x_n\| - \|x_{n+1} - x_n\| \\
& = \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \right| k \|x_{n+1} - x_n\| + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| \|f(x_n) + St_n\| \\
& \quad + \left| \frac{\gamma_{n+1} - 1 + \beta_{n+1}}{1 - \beta_{n+1}} \right| \|x_{n+1} - x_n\| \\
& = \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \right| k \|x_{n+1} - x_n\| + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} - \frac{\alpha_n}{1 - \beta_n} \right| \|f(x_n) + St_n\| \\
& \quad + \left| \frac{\alpha_{n+1}}{1 - \beta_{n+1}} \right| \|x_{n+1} - x_n\|.
\end{aligned} \tag{3.22}$$

This together with (i), (ii), and (iii) implies that

$$\limsup_{n \rightarrow \infty} (\|w_{n+1} - w_n\| - \|x_{n+1} - x_n\|) \leq 0. \tag{3.23}$$

Hence, by Lemma 2.3, we have

$$\lim_{n \rightarrow \infty} \|w_n - x_n\| = 0. \tag{3.24}$$

Consequently,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} (1 - \beta_n) \|w_n - x_n\| = 0. \tag{3.25}$$

From (3.18) and (3.19), we also have $\|z_{n+1} - z_n\| \rightarrow 0$, $\|t_{n+1} - t_n\| \rightarrow 0$ and $\|y_{n+1} - y_n\| \rightarrow 0$ as $n \rightarrow \infty$. Since

$$x_{n+1} - x_n = \alpha_n f(x_n) + \beta_n x_n + \gamma_n St_n - x_n = \alpha_n (f(x_n) - x_n) + \gamma_n (St_n - x_n), \tag{3.26}$$

it follows by (ii) and (3.25) that

$$\lim_{n \rightarrow \infty} \|x_n - St_n\| = 0. \tag{3.27}$$

Since $x^* \in F(S) \cap \Gamma$, from (3.15) and Lemma 2.2, we get

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \|\alpha_n f(x_n) + \beta_n x_n + \gamma_n S t_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|S t_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|t_n - x^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|P_E(y_n - \lambda A y_n) - P_E(y^* - \lambda A y^*)\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|(y_n - \lambda A y_n) - (y^* - \lambda A y^*)\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|(y_n - y^*) - \lambda(A y_n - A y^*)\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 \\
&\quad + \gamma_n \left[\|y_n - y^*\|^2 - 2\lambda \langle y_n - y^*, A y_n - A y^* \rangle + \lambda^2 \|A y_n - A y^*\|^2 \right] \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 \\
&\quad + \gamma_n \left[\|y_n - y^*\|^2 - 2\lambda \alpha \|A y_n - A y^*\|^2 + \lambda^2 \|A y_n - A y^*\|^2 \right] \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \left[\|y_n - y^*\|^2 + \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2 \right] \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \left[\|x_n - x^*\|^2 + \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2 \right] \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|x_n - x^*\|^2 + \gamma_n \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + (\beta_n + \gamma_n) \|x_n - x^*\|^2 + \gamma_n \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + (1 - \alpha_n) \|x_n - x^*\|^2 + \gamma_n \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \|x_n - x^*\|^2 + \gamma_n \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2.
\end{aligned} \tag{3.28}$$

Therefore, we have

$$\begin{aligned}
-\gamma_n \lambda(\lambda - 2\alpha) \|A y_n - A y^*\|^2 &\leq \alpha_n \|f(x_n) - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \\
&\quad \times (\|x_n - x^*\| - \|x_{n+1} - x^*\|) \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \|x_n - x_{n+1}\|.
\end{aligned} \tag{3.29}$$

From (ii), (iii), and $\|x_{n+1} - x_n\| \rightarrow 0$, as $n \rightarrow \infty$, we get $\|A y_n - A y^*\| \rightarrow 0$ as $n \rightarrow \infty$.

Since $x^* \in F(S) \cap Y$, from (3.11) and Lemma 2.2, we get

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \|\alpha_n f(x_n) + \beta_n x_n + \gamma_n S t_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|t_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|y_n - y^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|P_E(z_n - \mu B z_n) - P_E(z^* - \mu B z^*)\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|(z_n - \mu B z_n) - (z^* - \mu B z^*)\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|(z_n - z^*) - (\mu B z_n - \mu B z^*)\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n [\|z_n - z^*\|^2 + \mu(\mu - 2\beta) \|B z_n - B z^*\|^2] \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \|x_n - x^*\|^2 + \gamma_n \mu(\mu - 2\beta) \|B z_n - B z^*\|^2.
\end{aligned} \tag{3.30}$$

Thus, we also have

$$\begin{aligned}
-\gamma_n \mu(\mu - 2\beta) \|B z_n - B z^*\|^2 &\leq \alpha_n \|f(x_n) - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \\
&\quad \times (\|x_n - x^*\| - \|x_{n+1} - x^*\|) \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \|x_n - x_{n+1}\|.
\end{aligned} \tag{3.31}$$

By again (ii), (iii), and (3.25), we also get $\|B z_n - B z^*\| \rightarrow 0$ as $n \rightarrow \infty$.

Let $x^* \in F(S) \cap Y$; again from (3.12), (3.13) and Lemma 2.2, we get

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \|\alpha_n f(x_n) + \beta_n x_n + \gamma_n S t_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|t_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|z_n - z^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|(x_n - \tau C x_n) - (x^* - \tau C x^*)\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n [\|x_n - x^*\|^2 + \tau(\tau - 2\gamma) \|C x_n - C x^*\|^2] \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \|x_n - x^*\|^2 + \gamma_n \tau(\tau - 2\gamma) \|C x_n - C x^*\|^2.
\end{aligned} \tag{3.32}$$

Again, we have

$$\begin{aligned}
-\gamma_n \tau (\tau - 2\gamma) \|Cx_n - Cx^*\|^2 &\leq \alpha_n \|f(x_n) - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 \\
&= \alpha_n \|f(x_n) - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \\
&\quad \times (\|x_n - x^*\| - \|x_{n+1} - x^*\|) \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + (\|x_n - x^*\| + \|x_{n+1} - x^*\|) \|x_n - x_{n+1}\|.
\end{aligned} \tag{3.33}$$

Similarly again by (ii), (iii), and $\|x_n - x_{n+1}\| \rightarrow 0$ as $n \rightarrow \infty$, and from (3.33), we also that $\|Cx_n - Cx^*\| \rightarrow 0$.

On the other hand, we compute that

$$\begin{aligned}
\|z_n - z^*\|^2 &= \|P_E(x_n - \tau Cx_n) - P_E(x^* - \tau Cx^*)\|^2 \\
&\leq \langle (x_n - \tau Cx_n) - (x^* - \tau Cx^*), P_E(x_n - \tau Cx_n) - P_E(x^* - \tau Cx^*) \rangle \\
&= \langle (x_n - \tau Cx_n) - (x^* - \tau Cx^*), z_n - z^* \rangle \\
&= \frac{1}{2} \left[\|(x_n - \tau Cx_n) - (x^* - \tau Cx^*)\|^2 + \|z_n - z^*\|^2 \right. \\
&\quad \left. - \|(x_n - \tau Cx_n) - (x^* - \tau Cx^*) - (z_n - z^*)\|^2 \right] \\
&= \frac{1}{2} \left[\|(I - \tau C)x_n - (I - \tau C)x^*\|^2 + \|z_n - z^*\|^2 \right. \\
&\quad \left. - \|(x_n - \tau Cx_n) - (x^* - \tau Cx^*) - (z_n - z^*)\|^2 \right] \\
&\leq \frac{1}{2} \left[\|x_n - x^*\|^2 + \|z_n - z^*\|^2 - \|(x_n - z_n) - \tau(Cx_n - Cx^*) - (x^* - z^*)\|^2 \right] \\
&= \frac{1}{2} \left[\|x_n - x^*\|^2 + \|z_n - z^*\|^2 - \|(x_n - z_n) - (x^* - z^*) - \tau(Cx_n - Cx^*)\|^2 \right] \\
&= \frac{1}{2} \left[\|x_n - x^*\|^2 + \|z_n - z^*\|^2 - \|(x_n - z_n) - (x^* - z^*)\|^2 \right. \\
&\quad \left. + 2\tau \langle (x_n - z_n) - (x^* - z^*), Cx_n - Cx^* \rangle - \tau^2 \|Cx_n - Cx^*\|^2 \right].
\end{aligned} \tag{3.34}$$

So, we obtain

$$\begin{aligned}
\|z_n - z^*\|^2 &\leq \|x_n - x^*\|^2 - \|(x_n - z_n) - (x^* - z^*)\|^2 \\
&\quad + 2\tau \langle (x_n - z_n) - (x^* - z^*), Cx_n - Cx^* \rangle - \tau^2 \|Cx_n - Cx^*\|^2.
\end{aligned} \tag{3.35}$$

Hence, by (3.12), it follows that

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 &= \|\alpha_n f(x_n) + \beta_n x_n + \gamma_n St_n - x^*\|^2 \\
&\leq \alpha_n \|f(x_n) - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|St_n - x^*\|^2
\end{aligned}$$

$$\begin{aligned}
&\leq \alpha_n k \|x_n - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|t_n - x^*\|^2 \\
&\leq \alpha_n k \|x_n - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|z_n - z^*\|^2 \\
&\leq \alpha_n k \|x_n - x^*\|^2 + \beta_n \|x_n - x^*\|^2 + \gamma_n \|x_n - x^*\|^2 - \gamma_n \|(x_n - z_n) - (x^* - z^*)\|^2 \\
&\quad + 2\tau\gamma_n \langle (x_n - z_n) - (x^* - z^*), Cx_n - Cx^* \rangle - \tau^2 \gamma_n \|Cx_n - Cx^*\|^2 \\
&= \alpha_n k \|x_n - x^*\|^2 + (1 - \alpha_n) \|x_n - x^*\|^2 - \gamma_n \|(x_n - z_n) - (x^* - z^*)\|^2 \\
&\quad + 2\tau\gamma_n \langle (x_n - z_n) - (x^* - z^*), Cx_n - Cx^* \rangle - \tau^2 \gamma_n \|Cx_n - Cx^*\|^2 \\
&\leq \alpha_n k \|x_n - x^*\|^2 + \|x_n - x^*\|^2 - \gamma_n \|(x_n - z_n) - (x^* - z^*)\|^2 \\
&\quad + 2\tau\gamma_n \langle (x_n - z_n) - (x^* - z^*), Cx_n - Cx^* \rangle \\
&\leq \alpha_n k \|x_n - x^*\|^2 + \|x_n - x^*\|^2 - \gamma_n \|(x_n - z_n) - (x^* - z^*)\|^2 \\
&\quad + 2\tau\gamma_n \|(x_n - z_n) - (x^* - z^*)\| \|Cx_n - Cx^*\|,
\end{aligned} \tag{3.36}$$

which implies that

$$\begin{aligned}
\gamma_n \|(x_n - z_n) - (x^* - z^*)\|^2 &\leq \alpha_n k \|x_n - x^*\|^2 + \|x_n - x^*\|^2 - \|x_{n+1} - x^*\|^2 \\
&\quad + 2\tau\gamma_n \|(x_n - z_n) - (x^* - z^*)\| \|Cx_n - Cx^*\| \\
&\leq \alpha_n k \|x_{n+1} - x^*\|^2 + 2\gamma_n \tau \|(x_n - z_n) - (x^* - z^*)\| \|Cx_n - Cx^*\| \\
&\quad + \|x_n - x_{n+1}\| (\|x_n - x^*\| + \|x_{n+1} - x^*\|).
\end{aligned} \tag{3.37}$$

By (ii), (iii), $\|x_n - x_{n+1}\| \rightarrow 0$, and $\|Cx_n - Cx^*\| \rightarrow 0$ as $n \rightarrow \infty$, from (3.37) and we get $\|(x_n - z_n) - (x^* - z^*)\| \rightarrow 0$ as $n \rightarrow \infty$. Now, observe that

$$\begin{aligned}
\|(z_n - t_n) + (x^* - z^*)\|^2 &= \|z_n - P_E(y_n - \lambda A y_n) + P_E(y^* - \lambda A y^*) - z^*\|^2 \\
&= \|z_n - P_E(y_n - \lambda A y_n) + P_E(y^* - \lambda A y^*) \\
&\quad - z^* + \mu B z_n - \mu B z_n + \mu B z^* - \mu B z^*\|^2 \\
&= \|z_n - \mu B z_n - (z^* - \mu B z^*) \\
&\quad - [P_E(y_n - \lambda A y_n) - P_E(y^* - \lambda A y^*)] + \mu(B z_n - B z^*)\|^2
\end{aligned}$$

$$\begin{aligned}
&\leq \|z_n - \mu Bz_n - (z^* - \mu Bz^*) - [P_E(y_n - \lambda Ay_n) - P_E(y^* - \lambda Ay^*)]\|^2 \\
&\quad + 2\mu \langle Bz_n - Bz^*, z_n - \mu Bz_n - (z^* - \mu Bz^*) \\
&\quad \quad - [P_E(y_n - \lambda Ay_n) - P_E(y^* - \lambda Ay^*)] + \mu(Bz_n - Bz^*) \rangle \\
&= \|z_n - \mu Bz_n - (z^* - \mu Bz^*) - [P_E(y_n - \lambda Ay_n) - P_E(y^* - \lambda Ay^*)]\|^2 \\
&\quad + 2\mu \langle Bz_n - Bz^*, (z_n - t_n) + (x^* - z^*) \rangle \\
&\leq \|z_n - \mu Bz_n - (z^* - \mu Bz^*)\|^2 \\
&\quad - \|P_E(y_n - \lambda_n Ay_n) - P_E(y^* - \lambda_n Ay^*)\|^2 \\
&\quad + 2\mu \|Bz_n - Bz^*\| \|(z_n - t_n) + (x^* - z^*)\| \\
&\leq \|z_n - \mu Bz_n - (z^* - \mu Bz^*)\|^2 \\
&\quad - \|SP_E(y_n - \lambda_n Ay_n) - SP_E(y^* - \lambda_n Ay^*)\|^2 \\
&\quad + 2\mu \|Bz_n - Bz^*\| \|(z_n - t_n) + (x^* - z^*)\| \\
&= \|z_n - \mu Bz_n - (z^* - \mu Bz^*)\|^2 - \|St_n - Sx^*\|^2 \\
&\quad + 2\mu \|Bz_n - Bz^*\| \|(z_n - t_n) + (x^* - z^*)\| \\
&\leq \|z_n - \mu Bz_n - (z^* - \mu Bz^*) - (St_n - x^*)\| \\
&\quad \times (\|z_n - \mu Bz_n - (z^* - \mu Bz^*)\| + \|St_n - x^*\|) \\
&\quad + 2\mu \|Bz_n - Bz^*\| \|(z_n - t_n) + (x^* - z^*)\|.
\end{aligned} \tag{3.38}$$

Since $\|St_n - x_n\| \rightarrow 0$, $\|(x_n - z_n) - (x^* - z^*)\| \rightarrow 0$, and $\|Bz_n - Bz^*\| \rightarrow 0$, as $n \rightarrow \infty$, it follows that

$$\|(z_n - t_n) + (x^* - z^*)\| \rightarrow 0, \quad \text{as } n \rightarrow \infty. \tag{3.39}$$

Since

$$\|St_n - t_n\| \leq \|St_n - x_n\| + \|(x_n - z_n) - (x^* - z^*)\| + \|(z_n - t_n) + (x^* - z^*)\|, \tag{3.40}$$

we obtain

$$\lim_{n \rightarrow \infty} \|St_n - t_n\| = 0. \tag{3.41}$$

Next, we show that

$$\limsup_{n \rightarrow \infty} \langle f(\bar{x}) - \bar{x}, x_n - \bar{x} \rangle \leq 0, \quad (3.42)$$

where $\bar{x} = P_{F(S) \cap Y} f(\bar{x})$.

Indeed, since $\{t_n\}$ and $\{St_n\}$ are two bounded sequence in E , we can choose a subsequence $\{t_{n_i}\}$ of $\{t_n\}$ such that $t_{n_i} \rightarrow z \in E$ and

$$\limsup_{n \rightarrow \infty} \langle f(\bar{x}) - \bar{x}, St_n - \bar{x} \rangle = \lim_{i \rightarrow \infty} \langle f(\bar{x}) - \bar{x}, St_{n_i} - \bar{x} \rangle. \quad (3.43)$$

Since $\lim_{n \rightarrow \infty} \|St_n - t_n\| = 0$, we obtain $St_{n_i} \rightarrow z$ as $i \rightarrow \infty$. Now, we claim that $z \in F(S) \cap Y$. First by Lemma 2.5, it is easy to see that $z \in F(S)$.

Since $\|St_n - t_n\| \rightarrow 0$, $\|St_n - x_n\| \rightarrow 0$, and

$$\begin{aligned} \|t_n - x_n\| &= \|t_n - St_n + St_n - x_n\| \\ &\leq \|t_n - St_n\| + \|St_n - x_n\| \\ &= \|St_n - t_n\| + \|St_n - x_n\|, \end{aligned} \quad (3.44)$$

we conclude that $\|t_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty$. Furthermore, by Lemma 3.2 that G is nonexpansive, then

$$\begin{aligned} \|t_n - G(t_n)\| &= \|G(x_n) - G(t_n)\| \\ &\leq \|x_n - t_n\|. \end{aligned} \quad (3.45)$$

Thus $\lim_{n \rightarrow \infty} \|t_n - G(t_n)\| = 0$. According to Lemma 2.5, we obtain $z \in Y$. Therefore, there holds $z \in F(S) \cap Y$.

On the other hand, it follows from (2.2) that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle f(\bar{x}) - \bar{x}, x_n - \bar{x} \rangle &= \limsup_{n \rightarrow \infty} \langle f(\bar{x}) - \bar{x}, St_n - \bar{x} \rangle \\ &= \lim_{n \rightarrow \infty} \langle f(\bar{x}) - \bar{x}, St_{n_i} - \bar{x} \rangle \\ &= \langle f(\bar{x}) - \bar{x}, z - \bar{x} \rangle \\ &\leq 0. \end{aligned} \quad (3.46)$$

Finally, we show that $x_n \rightarrow \bar{x}$, and by (3.14) that

$$\begin{aligned}
\|x_{n+1} - \bar{x}\|^2 &= \|\alpha_n f(x_n) + \beta_n x_n + \gamma_n S t_n - \bar{x}\|^2 \\
&\leq \|\beta_n(x_n - \bar{x}) + \gamma_n(S t_n - \bar{x})\|^2 + 2\alpha_n \langle f(x_n) - \bar{x}, x_{n+1} - \bar{x} \rangle \\
&\leq \|\beta_n(x_n - \bar{x}) + \gamma_n(S t_n - \bar{x})\|^2 + 2\alpha_n \langle f(x_n) - f(\bar{x}), x_{n+1} - \bar{x} \rangle \\
&\quad + 2\alpha_n \langle f(\bar{x}) - \bar{x}, x_{n+1} - \bar{x} \rangle \\
&\leq \left[\beta_n \|x_n - \bar{x}\|^2 + \gamma_n \|S t_n - \bar{x}\|^2 \right] + 2\alpha_n \|f(x_n) - f(\bar{x})\| \|x_{n+1} - \bar{x}\| \\
&\quad + 2\alpha_n \langle f(\bar{x}) - \bar{x}, x_{n+1} - \bar{x} \rangle \\
&\leq \left[\beta_n \|x_n - \bar{x}\|^2 + \gamma_n \|t_n - \bar{x}\|^2 \right] + 2\alpha_n \alpha \|x_n - \bar{x}\| \|x_{n+1} - \bar{x}\| \\
&\quad + 2\alpha_n \langle f(\bar{x}) - \bar{x}, x_{n+1} - \bar{x} \rangle \\
&\leq (1 - \alpha_n)^2 \|x_n - \bar{x}\|^2 + \alpha_n \alpha \left(\|x_n - \bar{x}\|^2 + \|x_{n+1} - \bar{x}\|^2 \right) \\
&\quad + 2\alpha_n \langle f(\bar{x}) - \bar{x}, x_{n+1} - \bar{x} \rangle
\end{aligned} \tag{3.47}$$

which implies that

$$\begin{aligned}
\|x_{n+1} - \bar{x}\|^2 &\leq \left(1 - \frac{2(1-\alpha)\alpha_n}{1-\alpha\alpha_n} \right) \|x_n - \bar{x}\|^2 + \frac{\alpha_n^2}{1-\alpha\alpha_n} \|x_n - \bar{x}\|^2 \\
&\quad + \frac{2\alpha_n}{1-\alpha\alpha_n} \langle f(\bar{x}) - \bar{x}, x_{n+1} - \bar{x} \rangle \\
&:= (1 - \sigma_n) \|x_n - \bar{x}\|^2 + \delta_n, \quad n \geq 0,
\end{aligned} \tag{3.48}$$

where $\sigma_n = (2(1-\alpha)\alpha_n)/(1-\alpha\alpha_n)$ and $\delta_n = \alpha_n^2/(1-\alpha\alpha_n) \|x_n - \bar{x}\|^2 + 2\alpha_n/(1-\alpha\alpha_n) \langle f(\bar{x}) - \bar{x}, x_{n+1} - \bar{x} \rangle$. Therefore, by (3.46) and Lemma 2.4, we get that $\{x_n\}$ converges to $\bar{x}$, where $\bar{x} = P_{F(S) \cap Y} f(\bar{x})$. This completes the proof. $\square$

Setting $A = B = C$, we obtain the following corollary.

Corollary 3.4. *Let E be a nonempty, closed, convex subset of a real Hilbert space H . Let the mapping $A : E \rightarrow H$ be α -inverse-strongly monotone. Let S be a nonexpansive mapping of E into itself such that $F(S) \cap Y \neq \emptyset$. Let f be a contraction of H into itself and given $x_0 \in H$ arbitrarily and $\{x_n\}$ is generated by*

$$\begin{aligned}
z_n &= P_E(x_n - \tau A x_n), \\
y_n &= P_E(z_n - \mu A z_n), \\
x_{n+1} &= \alpha_n f(x_n) + \beta_n x_n + \gamma_n S P_E(y_n - \lambda A y_n), \quad n \geq 1,
\end{aligned} \tag{3.49}$$

where $\lambda, \mu, \tau \in (0, 2\alpha)$ and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ are three sequences in $[0, 1]$ such that

- (i) $\alpha_n + \beta_n + \gamma_n = 1$,
- (ii) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (iii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$.

Then, $\{x_n\}$ converges strongly to $\bar{x} \in F(S) \cap Y$, where $\bar{x} = P_{F(S) \cap Y} f(\bar{x})$ and $(\bar{x}, \bar{y}, \bar{z})$ is a solution of problem (1.10), where

$$\begin{aligned}\bar{y} &= P_E(\bar{z} - \mu A\bar{z}), \\ \bar{z} &= P_E(\bar{x} - \tau A\bar{x}).\end{aligned}\tag{3.50}$$

Setting $A \equiv B \equiv 0$ (the zero operators), we obtain the following corollary for solving the fixed points problem and the classical variational inequality problems.

Corollary 3.5. *Let E be a nonempty, closed, convex subset of a real Hilbert space H . Let the mapping $A : E \rightarrow H$ be α -inverse-strongly monotone. Let S be a nonexpansive mapping of E into itself such that $F(S) \cap VI(A, E) \neq \emptyset$. Let f be a contraction of H into itself and given $x_0 \in H$ arbitrarily and $\{x_n\}$ is generated by*

$$x_{n+1} = \alpha_n f(x_n) + \beta_n x_n + \gamma_n S P_E(x_n - \lambda A x_n), \quad n \geq 1, \tag{3.51}$$

where $\lambda \in (0, 2\alpha)$ and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ are three sequences in $[0, 1]$ such that

- (i) $\alpha_n + \beta_n + \gamma_n = 1$,
- (ii) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (iii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$.

Then, $\{x_n\}$ converges strongly to $\bar{x} \in F(S) \cap VI(A, E)$, where $\bar{x} = P_{F(S) \cap VI(A, E)} f(\bar{x})$.

4. Some Applications

We recall that a mapping $T : E \rightarrow E$ is called strictly pseudocontractive if there exists some k with $0 \leq k < 1$ such that

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + k\|(I - T)x - (I - T)y\|^2, \quad \forall x, y \in E. \tag{4.1}$$

For recent convergence result for strictly pseudocontractive mappings, put $A = I - T$. Then, we have

$$\|(I - A)x - (I - A)y\|^2 \leq \|x - y\|^2 + k\|Ax - Ay\|^2. \tag{4.2}$$

On the other hand,

$$\|(I - A)x - (I - A)y\|^2 \leq \|x - y\|^2 + \|Ax - Ay\|^2 - 2\langle x - y, Ax - Ay \rangle. \tag{4.3}$$

Hence, we have

$$\langle x - y, Ax - Ay \rangle \geq \frac{1-k}{2} \|Ax - Ay\|^2. \quad (4.4)$$

Consequently, if $T : E \rightarrow E$ is a strictly pseudocontractive mapping with constant k , then the mapping $A = I - T$ is $(1-k)/2$ -inverse-strongly monotone.

Setting $A = I - T$, $B = I - V$, and $C = I - W$, we obtain the following corollary.

Theorem 4.1. *Let E be a nonempty, closed, convex subset of a real Hilbert space H . Let T, V, W be strictly pseudocontractive mappings with constant k of C into itself, and let S be a nonexpansive mapping of E into itself such that $F(S) \cap Y \neq \emptyset$. Let f be a contraction of H into itself and given $x_0 \in H$ arbitrarily and $\{x_n\}$ is generated by*

$$\begin{aligned} z_n &= (I - \tau)x_n + \tau Wx_n, \\ y_n &= (I - \mu)z_n + \mu Vz_n, \\ x_{n+1} &= \alpha_n f(x_n) + \beta_n x_n + \gamma_n S((1 - \lambda)y_n + \lambda Ty_n), \quad n \geq 1, \end{aligned} \quad (4.5)$$

where $\lambda \in (0, 2\alpha)$, $\mu \in (0, 2\beta)$, and $\tau \in (0, 2\gamma)$ and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ are three sequences in $[0, 1]$ such that

- (i) $\alpha_n + \beta_n + \gamma_n = 1$,
- (ii) $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\sum_{n=1}^{\infty} \alpha_n = \infty$,
- (iii) $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$.

Then $\{x_n\}$ converges strongly to $\bar{x} \in F(S) \cap Y$, where $\bar{x} = P_{F(S) \cap Y} f(\bar{x})$ and $(\bar{x}, \bar{y}, \bar{z})$ is a solution of problem (1.9), where

$$\begin{aligned} \bar{y} &= P_E(\bar{z} - \mu B\bar{z}), \\ \bar{z} &= P_E(\bar{x} - \tau C\bar{x}). \end{aligned} \quad (4.6)$$

Proof. Since $A = I - T$, $B = I - V$, and $C = I - W$, we have

$$\begin{aligned} P_E(x_n - \tau Cx_n) &= (I - \tau)x_n + \tau Wx_n, \\ P_E(y_n - \lambda Ay_n) &= (I - \lambda)y_n + \lambda Ty_n, \\ P_E(z_n - \mu Bz_n) &= (I - \mu)z_n + \mu Vz_n. \end{aligned} \quad (4.7)$$

Thus, the conclusion follows immediately from Theorem 3.3. $\square$

If $f(x) = x_0$, for all $x \in E$, and $T = V = W$ in Theorem 4.1, we obtain the following corollary.

Corollary 4.2. *Let E be a nonempty, closed, convex subset of a real Hilbert space H . Let T be strictly pseudocontractive mappings with constant k of C into itself, and let S be a nonexpansive mapping of E into itself such that $F(S) \cap Y \neq \emptyset$. Given $x_0 \in H$ arbitrarily and $\{x_n\}$ is generated by*

$$\begin{aligned} z_n &= (I - \tau)x_n + \tau Tx_n, \\ y_n &= (I - \mu)z_n + \mu Tz_n, \\ x_{n+1} &= \alpha_n x_0 + \beta_n x_n + \gamma_n S((1 - \lambda)y_n + \lambda Ty_n), \quad n \geq 1, \end{aligned} \tag{4.8}$$

where $\lambda \in (0, 2\alpha)$, $\mu \in (0, 2\beta)$, and $\tau \in (0, 2\gamma)$ and $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}$ are three sequences in $[0, 1]$ such that

$$\begin{aligned} \alpha_n + \beta_n + \gamma_n &= 1, \\ \lim_{n \rightarrow \infty} \alpha_n &= 0 \text{ and } \sum_{n=1}^{\infty} \alpha_n = \infty, \\ 0 < \liminf_{n \rightarrow \infty} \beta_n &\leq \limsup_{n \rightarrow \infty} \beta_n < 1. \end{aligned} \tag{4.9}$$

Then, $\{x_n\}$ converges strongly to $\bar{x} \in F(S) \cap Y$, where $\bar{x} = P_{F(S) \cap Y} \bar{x}$ and $(\bar{x}, \bar{y}, \bar{z})$ is a solution of problem (1.10), where

$$\begin{aligned} \bar{y} &= P_E(\bar{z} - \mu A\bar{z}) \\ \bar{z} &= P_E(\bar{x} - \tau A\bar{x}). \end{aligned} \tag{4.10}$$

Acknowledgments

W. Kumam would like to give thanks to the Centre of Excellence in Mathematics, the Commission on Higher Education, Thailand, for their financial support for Ph.D. program at KMUTT. Moreover, the authors also would like to thank the Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission for financial support (under CSEC Project no. 54000267) and P. Kumam was supported by the Commission on Higher Education and the Thailand Research Fund under Grant no. MRG5380044.

References

- [1] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, pp. 197–228, 1967.
- [2] F. Liu and M. Z. Nashed, "Regularization of nonlinear ill-posed variational inequalities and convergence rates," *Set-Valued Analysis*, vol. 6, no. 4, pp. 313–344, 1998.
- [3] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [4] J. C. Yao and O. Chadli, "Pseudomonotone complementarity problems and variational inequalities," in *Handbook of Generalized Convexity and Generalized Monotonicity*, J. P. Crouzeix, N. Haddjissas, and S. Schaible, Eds., vol. 76, pp. 501–558, Springer, Amsterdam, the Netherlands, 2005.

- [5] Y. Yao, Y. C. Liou, and J. C. Yao, "An extragradient method for fixed point problems and variational inequality problems," *Journal of Inequalities and Applications*, vol. 2007, Article ID 38752, 12 pages, 2007.
- [6] L. C. Zeng and J. C. Yao, "Strong convergence theorem by an extragradient method for fixed point problems and variational inequality problems," *Taiwanese Journal of Mathematics*, vol. 10, no. 5, pp. 1293–1303, 2006.
- [7] W. Takahashi and M. Toyoda, "Weak convergence theorems for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 118, no. 2, pp. 417–428, 2003.
- [8] G. M. Korpelevič, "An extragradient method for finding saddle points and for other problems," *Matecon*, vol. 12, no. 4, pp. 747–756, 1976.
- [9] P. Kumam, "A relaxed extragradient approximation method of two inverse-strongly monotone mappings for a general system of variational inequalities, fixed point and equilibrium problems," *Bulletin of the Iranian Mathematical Society*, vol. 36, no. 1, pp. 227–252, 2010.
- [10] W. Kumam and P. Kumam, "Hybrid iterative scheme by a relaxed extragradient method for solutions of equilibrium problems and a general system of variational inequalities with application to optimization," *Nonlinear Analysis: Hybrid Systems*, vol. 3, no. 4, pp. 640–656, 2009.
- [11] N. Nadezhkina and W. Takahashi, "Weak convergence theorem by an extragradient method for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 128, no. 1, pp. 191–201, 2006.
- [12] L. C. Zeng, N. C. Wong, and J. C. Yao, "Strong convergence theorems for strictly pseudocontractive mappings of Browder-Petryshyn type," *Taiwanese Journal of Mathematics*, vol. 10, no. 4, pp. 837–849, 2006.
- [13] Y. Yao and J. C. Yao, "On modified iterative method for nonexpansive mappings and monotone mappings," *Applied Mathematics and Computation*, vol. 186, no. 2, pp. 1551–1558, 2007.
- [14] L. C. Ceng, C. Y. Wang, and J. C. Yao, "Strong convergence theorems by a relaxed extragradient method for a general system of variational inequalities," *Mathematical Methods of Operations Research*, vol. 67, no. 3, pp. 375–390, 2008.
- [15] R. U. Verma, "On a new system of nonlinear variational inequalities and associated iterative algorithms," *Mathematical Sciences Research Hot-Line*, vol. 3, no. 8, pp. 65–68, 1999.
- [16] R. U. Verma, "Iterative algorithms and a new system of nonlinear variational inequalities," *Advances in Nonlinear Variational Inequalities*, vol. 3, no. 8, pp. 117–124.
- [17] G. Kassay, J. Kolumbán, and Z. Páles, "Factorization of minty and stampania variational inequality systems," *European Journal of Operational Research*, vol. 143, no. 2, pp. 377–389, 2002.
- [18] G. Kassay and J. Kolumbán, "System of multi-valued variational inequalities," *Publicationes Mathematicae Debrecen*, vol. 56, no. 1-2, pp. 185–195, 2000.
- [19] S. Zhang, J. Lee, and C. Chan, "Algorithms of common solutions to quasi variational inclusion and fixed point problems," *Applied Mathematics and Mechanics*, vol. 29, no. 5, pp. 571–581, 2008.
- [20] M. O. Osilike and D. I. Igbokwe, "Weak and strong convergence theorems for fixed points of pseudocontractions and solutions of monotone type operator equations," *Computers & Mathematics with Applications*, vol. 40, no. 4-5, pp. 559–567, 2000.
- [21] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter non-expansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [22] H. K. Xu, "Viscosity approximation methods for nonexpansive mappings," *Journal of Mathematical Analysis and Applications*, vol. 298, no. 1, pp. 279–291, 2004.
- [23] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge University Press, Cambridge, UK, 1990.