Research Article

Finite Difference and Iteration Methods for Fractional Hyperbolic Partial Differential Equations with the Neumann Condition

Allaberen Ashyralyev^{1,2} and Fadime Dal³

¹ *Department of Mathematics, Fatih University, Buyukcekmece 34500, Istanbul, Turkey*

² *Department of Mathematics, ITTU, Ashgabad, Turkmenistan*

³ *Department of Mathematics, Ege University, 35100 Izmir, Turkey*

Correspondence should be addressed to Fadime Dal, fadimedal@hotmail.com

Received 19 December 2011; Accepted 18 April 2012

Academic Editor: Chuanxi Qian

Copyright © 2012 A. Ashyralyev and F. Dal. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The numerical and analytic solutions of the mixed problem for multidimensional fractional hyperbolic partial differential equations with the Neumann condition are presented. The stable difference scheme for the numerical solution of the mixed problem for the multidimensional fractional hyperbolic equation with the Neumann condition is presented. Stability estimates for the solution of this difference scheme and for the first- and second-order difference derivatives are obtained. A procedure of modified Gauss elimination method is used for solving this difference scheme in the case of one-dimensional fractional hyperbolic partial differential equations. He's variational iteration method is applied. The comparison of these methods is presented.

1. Introduction

It is known that various problems in fluid mechanics (dynamics, elasticity) and other areas of physics lead to fractional partial differential equations. Methods of solutions of problems for fractional differential equations have been studied extensively by many researchers (see, e.g., [1–15] and the references given therein).

The role played by stability inequalities (well posedness) in the study of boundary-value problems for hyperbolic partial differential equations is well known (see, e.g., [16–29]).

In the present paper, finite difference and He's iteration methods for the approximate solutions of the mixed boundary-value problem for the multidimensional fractional hyperbolic equation

$$\begin{aligned} \frac{\partial^2 u(t, x)}{\partial t^2} - \sum_{r=1}^m (a_r(x) u_{x_r})_{x_r} + D_t^{1/2} u(t, x) + \sigma u(t, x) &= f(t, x), \\ x = (x_1, \dots, x_m) \in \Omega, \quad 0 < t < 1, \\ u(0, x) = 0, \quad u_t(0, x) = 0, \quad x \in \overline{\Omega}; \quad \frac{\partial u(t, x)}{\partial \bar{n}} &= 0, \quad x \in S, \end{aligned} \quad (1.1)$$

are studied. Here Ω is the unit open cube in the m -dimensional Euclidean space: $\mathbb{R}^m : \{\Omega = x = (x_1, \dots, x_m) : 0 < x_j < 1, 1 \leq j \leq m\}$ with boundary S , $\overline{\Omega} = \Omega \cup S$; $a_r(x)$ ($x \in \Omega$) and $f(t, x)$ ($t \in (0, 1), x \in \Omega$) are given smooth functions and $a_r(x) \geq a > 0$.

1.1. Definition

The Caputo fractional derivative of order $\alpha > 0$ of a continuous function $u(t, x)$ is defined by

$$D_{a^+}^\alpha u(t, x) = \frac{1}{\Gamma(1-\alpha)} \int_a^t \frac{u'(s, x)}{(t-s)^\alpha} ds, \quad (1.2)$$

where $\Gamma(\cdot)$ is the gamma function.

2. The Finite Difference Method

In this section, we consider the first order of accuracy in t and the second-orders of accuracy in space variables' stable difference scheme for the approximate solution of problem (1.1). The stability estimates for the solution of this difference scheme and its first- and second-order difference derivatives are established. A procedure of modified Gauss elimination method is used for solving this difference scheme in the case of one-dimensional fractional hyperbolic partial differential equations.

2.1. The Difference Scheme: Stability Estimates

The discretization of problem (1.1) is carried out in two steps. In the first step, let us define the grid space

$$\begin{aligned} \tilde{\Omega}_h &= \{x = x_r = (h_1 r_1, \dots, h_m r_m), r = (r_1, \dots, r_m), 0 \leq r_j \leq N_j, h_j N_j = 1, j = 1, \dots, m\}, \\ \Omega_h &= \tilde{\Omega}_h \cap \Omega, \quad S_h = \tilde{\Omega}_h \cap S. \end{aligned} \quad (2.1)$$

We introduce the Banach space $L_{2h} = L_2(\tilde{\Omega}_h)$ of the grid functions $\varphi^h(x) = \{\varphi(h_1 r_1, \dots, h_m r_m)\}$ defined on $\tilde{\Omega}_h$, equipped with the norm

$$\|\varphi^h\|_{L_2(\tilde{\Omega}_h)} = \left(\sum_{x \in \tilde{\Omega}_h} |\varphi^h(x)|^2 h_1 \cdots h_m \right)^{1/2}. \quad (2.2)$$

To the differential operator A^x generated by problem (1.1), we assign the difference operator A_h^x by the formula

$$A_h^x u^h = - \sum_{r=1}^m \left(a_r(x) u_{\bar{x}_r}^h \right)_{x_{r,j}} + \sigma u_x^h \quad (2.3)$$

acting in the space of grid functions $u^h(x)$, satisfying the conditions $D_h u^h(x) = 0$ for all $x \in S_h$. It is known that A_h^x is a self-adjoint positive definite operator in $L_2(\tilde{\Omega}_h)$. With the help of A_h^x we arrive at the initial boundary value problem

$$\frac{d^2 v^h(t, x)}{dt^2} + D_t^{1/2} v^h(t, x) + A_h^x v^h(t, x) = f^h(t, x), \quad 0 \leq t \leq 1, x \in \Omega_h, \quad (2.4)$$

$$v^h(0, x) = 0, \quad \frac{dv^h(0, x)}{dt} = 0, \quad x \in \tilde{\Omega}_h,$$

for an infinite system of ordinary fractional differential equations.

In the second step, we replace problem (2.4) by the first order of accuracy difference scheme

$$\frac{u_{k+1}^h(x) - 2u_k^h(x) + u_{k-1}^h(x)}{\tau^2} + \frac{1}{\sqrt{\pi}} \sum_{m=1}^k \frac{\Gamma(k-m+1/2)}{(k-m)!} \frac{u_m^h - u_{m-1}^h}{\tau^{1/2}} + A_h^x u_{k+1}^h = f_k^h(x), \quad x \in \tilde{\Omega}_h,$$

$$f_k^h(x) = f(t_k, x), \quad t_k = k\tau, \quad 1 \leq k \leq N-1, \quad N\tau = 1, \quad x \in \tilde{\Omega}_h,$$

$$\frac{u_1^h(x) - u_0^h(x)}{\tau} = 0, \quad u_0^h(x) = 0, \quad x \in \tilde{\Omega}_h. \quad (2.5)$$

Here $\Gamma(k-m+1/2) = \int_0^\infty t^{k-m-1/2} e^{-t} dt$.

Theorem 2.1. Let τ and $|h| = \sqrt{h_1^2 + \dots + h_m^2}$ be sufficiently small numbers. Then, the solutions of difference scheme (2.5) satisfy the following stability estimates:

$$\begin{aligned} \max_{1 \leq k \leq N} \|u_k^h\|_{L_{2h}} + \max_{1 \leq k \leq N} \left\| \frac{u_k^h - u_{k-1}^h}{\tau} \right\|_{L_{2h}} &\leq C_1 \max_{1 \leq k \leq N-1} \|f_k^h\|_{L_{2h}}, \\ \max_{1 \leq k \leq N-1} \left\| \tau^{-2} (u_{k+1}^h - 2u_k^h + u_{k-1}^h) \right\|_{L_{2h}} + \max_{1 \leq k \leq N} \left\| (u_k^h)_{\bar{x}_r \bar{x}_r} \right\|_{L_{2h}} & \\ \leq C_2 \left[\|f_1^h\|_{L_{2h}} + \max_{2 \leq k \leq N-1} \left\| \tau^{-1} (f_k^h - f_{k-1}^h) \right\|_{L_{2h}} \right]. & \end{aligned} \quad (2.6)$$

Here C_1 and C_2 do not depend on τ , h , and $f_k^h, 1 \leq k \leq N-1$.

The proof of Theorem 2.1 is based on the self-adjointness and positive definiteness of operator A_h^x in L_{2h} and on the following theorem on the coercivity inequality for the solution of the elliptic difference problem in L_{2h} .

Theorem 2.2. For the solutions of the elliptic difference problem

$$\begin{aligned} A_h^x u^h(x) &= \omega^h(x), \quad x \in \Omega_h, \\ D_h u^h(x) &= 0, \quad x \in S_h, \end{aligned} \quad (2.7)$$

the following coercivity inequality holds [30]:

$$\sum_{r=1}^m \|u_{x_r \bar{x}_r}^h\|_{L_{2h}} \leq C \|\omega^h\|_{L_{2h}}. \quad (2.8)$$

Finally, applying this difference scheme, the numerical methods are proposed in the following section for solving the one-dimensional fractional hyperbolic partial differential equation. The method is illustrated by numerical examples.

2.2. Numerical Results

For the numerical result, the mixed problem

$$\begin{aligned} D_t^2 u(t, x) + D_t^{1/2} u(t, x) - u_{xx}(t, x) + u(t, x) &= f(t, x), \\ f(t, x) &= \left(2 + t^2 + \frac{8t^{3/2}}{3\sqrt{\pi}} + (\pi t)^2 \right) \cos(\pi x), \quad 0 < t, x < 1, \\ u(0, x) &= 0, \quad u_t(0, x) = 0, \quad 0 \leq x \leq 1, \\ u_x(t, 0) &= u_x(t, 1) = 0, \quad 0 \leq t \leq 1 \end{aligned} \quad (2.9)$$

for solving the one-dimensional fractional hyperbolic partial differential equation is considered. Applying difference scheme (2.5), we obtained

$$\begin{aligned}
 & \frac{u_n^{k+1} - 2u_n^k + u_n^{k-1}}{\tau^2} + \frac{1}{\sqrt{\pi}} \sum_{m=1}^k \frac{\Gamma(k-m+1/2)}{(k-m)!} \left(\frac{u_n^m - u_n^{m-1}}{\tau^{1/2}} \right) - \left(\frac{u_{n+1}^{k+1} - 2u_n^{k+1} + u_{n-1}^{k+1}}{h^2} \right) + u_n^k = \varphi_n^k, \\
 & \varphi_n^k = f(t_k, x_n), \quad 1 \leq k \leq N-1, 1 \leq n \leq M-1, \\
 & u_n^0 = 0, \quad \tau^{-1}(u_n^1 - u_n^0) = 0, \quad 0 \leq n \leq M, \\
 & u_1^k - u_0^k = u_M^k - u_{M-1}^k = 0, \quad 0 \leq k \leq N.
 \end{aligned} \tag{2.10}$$

We get the system of equations in the matrix form:

$$\begin{aligned}
 AU_{n+1} + BU_n + CU_{n-1} &= D\varphi_n, \quad 1 \leq n \leq M-1, \\
 U_1 &= U_0, \quad U_M = U_{M-1},
 \end{aligned} \tag{2.11}$$

where

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & a & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & a & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & \cdots & a & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 \end{bmatrix}_{(N+1) \times (N+1)},$$

$$B = \begin{bmatrix} b_{1,1} & 0 & 0 & 0 & \cdots & 0 & 0 \\ b_{2,1} & b_{2,2} & 0 & 0 & \cdots & 0 & 0 \\ b_{3,1} & b_{3,2} & b_{3,3} & 0 & \cdots & 0 & 0 \\ b_{4,1} & b_{4,2} & b_{4,3} & b_{4,4} & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ b_{N,1} & b_{N,2} & b_{N,3} & b_{N,4} & \cdots & b_{N,N} & 0 \\ b_{N+1,1} & b_{N+1,2} & b_{N+1,3} & b_{N+1,4} & \cdots & b_{N+1,N} & b_{N+1,N+1} \end{bmatrix}_{(N+1) \times (N+1)},$$

$$C = A,$$

$$D = \begin{bmatrix} 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 1 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & \cdots & 1 & 0 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 \end{bmatrix}_{(N+1) \times (N+1)},$$

$$U_s = \begin{bmatrix} U_s^0 \\ U_s^1 \\ U_s^2 \\ U_s^3 \\ \cdots \\ U_s^{N-1} \\ U_s^N \end{bmatrix}_{(N+1) \times (1)}, \quad s = n-1, n, n+1.$$
(2.12)

Here

$$a = -\frac{1}{h^2}, \quad b_{1,1} = 1, \quad b_{2,1} = -1, \quad b_{2,2} = 1, \quad b_{3,1} = \frac{1}{\tau^2} - \frac{1}{\tau^{1/2}},$$

$$b_{3,2} = -\frac{2}{\tau^2} + \frac{1}{\tau^{1/2}}, \quad b_{3,3} = 1 + \frac{1}{\tau^2} + \frac{2}{h^2},$$

$$b_{k+2,1} = -\frac{1}{\sqrt{\pi}} \frac{\Gamma(k-1+1/2)}{\Gamma(k)\tau^{1/2}}, \quad 2 \leq k \leq N-1,$$

$$b_{k+2,k+1} = -\frac{2}{\tau^2} + \frac{1}{\tau^{1/2}}, \quad 1 \leq k \leq N-1,$$

$$b_{k+2,k} = \frac{1}{\tau^2} + \frac{1}{\sqrt{\pi}} \left(\frac{\Gamma(1+0.5)}{\Gamma(2)} - \frac{\Gamma(0.5)}{\Gamma(1)} \right) \frac{1}{\tau^{1/2}}, \quad 2 \leq k \leq N-1,$$

$$b_{k+2,k+2} = 1 + \frac{1}{\tau^2} + \frac{2}{h^2}, \quad 1 \leq k \leq N-1,$$

$$b_{k+2,i+1} = \frac{1}{\sqrt{\pi}} \left(\frac{\Gamma(k-i+1/2)}{\Gamma(k-(i-1))} - \frac{\Gamma(k-(i+1)+1/2)}{\Gamma(k-(i-1)-1)} \right) \frac{1}{\tau^{1/2}}, \quad 3 \leq k \leq N-1, 1 \leq i \leq k-2,$$

$$\varphi_n^k = \left(2 + (k\tau)^2 + \frac{8(k\tau)^{3/2}}{3\sqrt{\pi}} + (\pi k\tau)^2 \right) \cos \pi(nh),$$

$$\varphi_n = \begin{bmatrix} \varphi_n^0 \\ \varphi_n^1 \\ \varphi_n^2 \\ \cdots \\ \varphi_n^N \end{bmatrix}_{(N+1) \times 1}.$$
(2.13)

So, we have the second-order difference equation with respect to n matrix coefficients. To solve this difference equation, we have applied a procedure of modified Gauss elimination method for difference equation with respect to k matrix coefficients. Hence, we seek a solution of the matrix equation in the following form:

$$U_j = \alpha_{j+1}U_{j+1} + \beta_{j+1}, \quad (2.14)$$

$n = M-1, \dots, 2, 1$, $\alpha_j (j = 1, \dots, M)$ are $(N+1) \times (N+1)$ square matrices, and $\beta_j (j = 1, \dots, M)$ are $(N+1) \times 1$ column matrices defined by

$$\begin{aligned} \alpha_{n+1} &= (B + C\alpha_n)^{-1}(-A), \\ \beta_{n+1} &= (B + C\alpha_n)^{-1}(D\varphi_n - C\beta_n), \quad n = 2, 3, \dots, M, \end{aligned} \quad (2.15)$$

where

$$\begin{aligned} \alpha_1 &= \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}_{(N+1) \times (N+1)}, \\ \beta_1 &= \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cdots \\ 0 \end{bmatrix}_{(N+1) \times 1}. \end{aligned} \quad (2.16)$$

Now, we will give the results of the numerical analysis. First, we give an estimate for the constants C_1 and C_2 figuring in the stability estimates of Theorem 2.1. We have

$$\begin{aligned} C_1 &= \max_{f,u}(C_{t1}), \quad C_2 = \max_{f,u}(C_{t2}), \\ C_{t1} &= \left[\max_{1 \leq k \leq N} \|u_k^h\|_{L_{2h}} + \max_{1 \leq k \leq N} \|\tau^{-1}(u_k^h - u_{k-1}^h)\|_{L_{2h}} \right] \times \left(\max_{1 \leq k \leq N-1} \|f_k^h\|_{L_{2h}} \right)^{-1}, \\ &= \left[\max_{1 \leq k \leq N-1} \|\tau^{-2}(u_{k+1}^h - 2u_k^h + u_{k-1}^h)\|_{L_{2h}} + \max_{1 \leq k \leq N} \sum_{r=1}^n \|(u_k^h)_{\bar{x}_r, \bar{x}_r}\|_{L_{2h}} \right] \\ &\quad \times \left(\max_{2 \leq k \leq N-1} \|\tau^{-1}(f_k^h - f_{k-1}^h)\|_{L_{2h}} + \|f_1^h\|_{L_{2h}} \right)^{-1}. \end{aligned} \quad (2.17)$$

The constants C_{t1} and C_{t2} in the case of numerical solution of initial-boundary value problem (2.9) are computed. The constants C_{t1} and C_{t2} are given in Table 1 for $N = 20, 40, 80$, and $M = 80$, respectively.

Table 1: Stability estimates for (2.9).

	$M = 80$ $N = 20$	$M = 80$ $N = 40$	$M = 80$ $N = 80$
The values of C_{t1}	0.2096	0.2073	0.2061
The values of C_{t2}	0.2075	0.1223	0.0670

Table 2: Comparison of the errors for the difference scheme.

Method	$M = 80$ $N = 20$	$M = 80$ $N = 40$	$M = 80$ $N = 60$
Comparison of errors (E_0) for approximate solutions	0.0071	0.0037	0.0008
Comparison of errors (E_1) for approximate solutions	0.1030	0.0521	0.0491
Comparison of errors (E_2) for approximate solutions	0.1224	0.0806	0.0882

Second, for the accurate comparison of the difference scheme considered, the errors computed by

$$\begin{aligned}
 E_0 &= \max_{1 \leq k \leq N-1} \left(\sum_{n=1}^{M-1} |u(t_k, x_n) - u_n^k|^2 h \right)^{1/2}, \\
 E_1 &= \max_{1 \leq k \leq N-1} \left(\sum_{n=1}^{M-1} \left| u_t(t_k, x_n) - \frac{(u_n^{k+1} - u_n^{k-1}))}{2\tau} \right|^2 h \right)^{1/2}, \\
 E_2 &= \max_{1 \leq k \leq N-1} \left(\sum_{n=1}^{M-1} \left| u_{tt}(t_k, x_n) - \frac{(u_n^{k+1} - 2u_n^k + u_n^{k-1}))}{\tau^2} \right|^2 h \right)^{1/2}
 \end{aligned} \tag{2.18}$$

of the numerical solutions are recorded for higher values of $N = M$, where $u(t_k, x_n)$ represents the exact solution and u_n^k represents the numerical solution at (t_k, x_n) . The errors E_0, E_1 and E_2 results are shown in Table 2 for $N = 20, 40, 60$ and $M = 60$, respectively.

The figure of the difference scheme solution of (2.9) is given by the Figure 2. The exact solution of (2.9) is given by as follows:

$$u(t, x) = t^2 \cos(\pi x). \tag{2.19}$$

The figure of the exact solution of (2.9) is shown by the Figure 1.

Figure 1: The surface shows the exact solution $u(t, x)$ for (2.9).

Figure 2: Difference scheme solution for (2.9).

3. He's Variational Iteration Method

In the present paper, the mixed boundary value problem for the multidimensional fractional hyperbolic equation (1.1) is considered. The correction functional for (1.1) can be approximately expressed as follows:

$$u_{n+1}(t, x) = u_n(t, x) + \int_0^t \lambda \left[\frac{\partial^2 u(s, x)}{\partial s^2} - \sum_{r=1}^m (a_r(x) \tilde{u}_{x_r})_{x_r} + D_s^{1/2} \tilde{u}(s, x) + \sigma \tilde{u}(s, x) - f(s, x) \right] ds, \quad (3.1)$$

where λ is a general Lagrangian multiplier (see, e.g., [31]) and $\tilde{u}$ is considered as a restricted variation as a restricted variation (see, e.g., [32]); that is, $\delta \tilde{u} = 0$, $u_0(t, x)$ is its initial

approximation. Using the above correction functional stationary and noticing that $\delta\tilde{u} = 0$, we obtain

$$\begin{aligned}\delta u_{n+1}(t, x) &= \delta u_n(t, x) + \int_0^t \delta \lambda \left[\frac{\partial u_n^2(t, x)}{\partial s^2} \right] ds, \\ \delta u_{n+1}(t, x) &= \delta u_n(t, x) - \frac{\partial \lambda}{\partial s} \delta u_n(s, x) \Big|_{s=t} + \lambda \frac{\partial}{\partial s} (\delta u_n(s, x)) \Big|_{s=t} + \int_0^t \frac{\partial^2 \lambda(t, s)}{\partial s^2} \delta u_n(s, x) ds = 0.\end{aligned}\tag{3.2}$$

From the above relation for any δu_n , we get the Euler-Lagrange equation:

$$\frac{\partial \lambda^2(t, s)}{\partial s^2} = 0, \tag{3.3}$$

with the following natural boundary conditions:

$$\begin{aligned}1 - \frac{\partial \lambda(t, s)}{\partial s} \Big|_{s=t} &= 0, \\ \lambda(t, s) \Big|_{s=t} &= 0.\end{aligned}\tag{3.4}$$

Therefore, the Lagrange multiplier can be identified as follows:

$$\lambda(t, s) = s - t. \tag{3.5}$$

Substituting the identified Lagrange multiplier into (3.1), the following variational iteration formula can be obtained:

$$u_{n+1}(t, x) = u_n(t, x) + \int_0^t (s-t) \left[\frac{\partial^2 u_n(s, x)}{\partial s^2} - \sum_{r=1}^m (a_r(x) u_{n_{x_r}})_{x_r} + D_s^{1/2} u_n(s, x) + u_n(s, x) - f(s, x) \right] ds. \tag{3.6}$$

In this case, let an initial approximation $u_0(t, x) = u(0, x) + t u_t(0, x)$. Then approximate solution takes the form $u(t, x) = \lim_{n \rightarrow \infty} u_n(t, x)$.

3.1. Variational Iteration Solution 1

For the numerical result, the mixed problem

$$\begin{aligned}
 D_t^2 u(t, x) + D_t^{1/2} u(t, x) - u_{xx}(t, x) + u(t, x) &= f(t, x), \\
 f(t, x) &= \left(2 + t^2 + \frac{8t^{3/2}}{3\sqrt{\pi}} + (\pi t)^2 \right) \cos(\pi x), \quad 0 < t, x < 1, \\
 u(0, x) &= 0, \quad u_t(0, x) = 0, \quad 0 \leq x \leq 1, \\
 u_x(t, 0) &= u_x(t, 1) = 0, \quad 0 \leq t \leq 1
 \end{aligned} \tag{3.7}$$

for solving the one-dimensional fractional hyperbolic partial differential equation is considered.

According to formula (3.6), the iteration formula for (3.7) is given by

$$\begin{aligned}
 u_{n+1}(t, x) &= u_n(t, x) \\
 &+ \int_0^t (s-t) \left[\frac{\partial u_n^2(s, x)}{\partial s^2} + D_s^{1/2} u_n(s, x) - \frac{\partial u_n^2(s, x)}{\partial x^2} + u_n(s, x) - f(s, x) \right] ds.
 \end{aligned} \tag{3.8}$$

Now we start with an initial approximation

$$u_0(t, x) = u(0, x) + t u_t(0, x). \tag{3.9}$$

Using the above iteration formula (3.8), we can obtain the other components as

$$\begin{aligned}
 u_0(t, x) &= 0, \\
 u_1(t, x) &= \frac{1}{420\sqrt{\pi}} \left(128t^{7/2} + 35t^4\sqrt{\pi} + 35t^4\pi^{5/2} + 420t^2\sqrt{\pi} \right) \cos(\pi x), \\
 u_2(t, x) &= \frac{1}{420\sqrt{\pi}} \cos(\pi x) \left(128t^{7/2} + 35t^4\sqrt{\pi} + 35t^4\pi^{5/2} + 420t^2\sqrt{\pi} \right) \\
 &\quad + \cos(\pi x) \left[-0.9058003666t^4 - 0.1510268880t^{11/2} \right. \\
 &\quad \left. - 0.1719434921t^{7/2} - 0.3281897218t^6 - 0.01666666667t^5 \right] \\
 &\quad \vdots
 \end{aligned} \tag{3.10}$$

The figure of (3.10) is given by the Figure 3.

Figure 3: Variational iteration method for (3.10).

Figure 4: Variational iteration method for (3.14).

3.2. Variational Iteration Solution 2

For the numerical result, the mixed problem

$$\begin{aligned}
 D_t^2 u(t, x, y) + D_t^{1/2} u(t, x, y) - u_{xx}(t, x, y) - u_{yy}(t, x, y) + u(t, x, y) &= f(t, x, y), \\
 f(t, x, y) &= \left(2 + t^2 + \frac{8t^{3/2}}{3\sqrt{\pi}} + (\pi t)^2 \right) \cos(\pi x) \cos(\pi y), \quad 0 < t, x < 1, y < 1, \\
 u(0, x, y) &= 0, \quad u_t(0, x, y) = 0, \quad 0 \leq x \leq 1, 0 \leq y \leq 1, \\
 u_x(t, 0, y) &= u_x(t, 1, y) = 0, \quad 0 \leq t \leq 1, 0 \leq y \leq 1, \\
 u_y(t, x, 0) &= u_y(t, x, 1) = 0, \quad 0 \leq t \leq 1, 0 \leq x \leq 1
 \end{aligned} \tag{3.11}$$

for solving the two-dimensional fractional hyperbolic partial differential equation is considered.

Figure 5: The surface shows the exact solution $u(t, x, y)$ for (3.11).

According to formula (3.6), the iteration formula for (3.11) is given by

$$\begin{aligned}
 u_{n+1}(t, x, y) = & u_n(t, x, y) \\
 & + \int_0^t (s-t) \left[\frac{\partial^2 u_n(s, x, y)}{\partial s^2} + D_s^{1/2} u_n(s, x, y) - \frac{\partial u_n^2(s, x, y)}{\partial x^2} \right. \\
 & \left. - \frac{\partial u_n^2(s, x, y)}{\partial y^2} + u_n(s, x, y) - f(s, x, y) \right] ds;
 \end{aligned} \tag{3.12}$$

we start with an initial approximation

$$u_0(t, x, y) = u(0, x, y) + tu_t(0, x, y). \tag{3.13}$$

Using the above iteration formula (3.12), we can obtain the other components as

$$\begin{aligned}
 u_0(t, x, y) &= 0, \\
 u_1(t, x, y) &= \frac{1}{420\sqrt{\pi}} \left(128t^{7/2} + 35t^4\sqrt{\pi} + 35t^4\pi^{5/2} + 420t^2\sqrt{\pi} \right) \cos(\pi x) \cos(\pi y),
 \end{aligned}$$

$$\begin{aligned}
u_2(t, x, y) = & \cos(\pi x) \cos(\pi y) \left[\frac{32}{105} \left(\frac{1}{t}\right)^{5/2} t^{7/2} + \left(\frac{1}{12}\right) \left(\frac{1}{t}\right)^{5/2} t^4 \sqrt{\pi} \right. \\
& + \frac{1}{12} \left(\frac{1}{t}\right)^{5/2} t^4 \pi^{5/2} + \left(\frac{1}{t}\right)^{5/2} t^2 \sqrt{\pi} \left. \right] + \cos(\pi x) \cos(\pi y) \\
& \times \left[-0.2195931203 t^{11/2} \sqrt{\pi} \left(\frac{1}{t}\right)^{5/2} - 0.1719434921 \sqrt{\pi} t^{7/2} \right. \\
& \times \left(\frac{1}{t}\right)^{5/2} - 0.6261860981 t^6 \sqrt{\pi} \left(\frac{1}{t}\right)^{5/2} \\
& \left. - 1.728267400 \sqrt{\pi} t^4 \left(\frac{1}{t}\right)^{5/2} - 0.01666666667 \sqrt{\pi} t^{5/2} \right] \\
& \vdots
\end{aligned} \tag{3.14}$$

The exact solution of (3.11) is given by as follows:

$$u(t, x, y) = t^2 \cos(\pi x) \cos(\pi y). \tag{3.15}$$

The figure of the exact solution of (3.11) is shown by the Figure 5.

The figure of (3.14) is given by the Figure 4, and so on; in the same manner the rest of the components of the iteration formula (3.12) can be obtained using the Maple package.

References

- [1] I. Podlubny, *Fractional Differential Equations*, vol. 198 of *Mathematics in Science and Engineering*, Academic Press, San Diego, Calif, USA, 1999.
- [2] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives*, Gordon and Breach Science Publishers, London, UK, 1993.
- [3] J.-L. Lavoie, T. J. Osler, and R. Tremblay, "Fractional derivatives and special functions," *SIAM Review*, vol. 18, no. 2, pp. 240–268, 1976.
- [4] V. E. Tarasov, "Fractional derivative as fractional power of derivative," *International Journal of Mathematics*, vol. 18, no. 3, pp. 281–299, 2007.
- [5] A. E. M. El-Mesiry, A. M. A. El-Sayed, and H. A. A. El-Saka, "Numerical methods for multi-term fractional (arbitrary) orders differential equations," *Applied Mathematics and Computation*, vol. 160, no. 3, pp. 683–699, 2005.
- [6] A. M. A. El-Sayed and F. M. Gaafar, "Fractional-order differential equations with memory and fractional-order relaxation-oscillation model," *Pure Mathematics and Applications*, vol. 12, no. 3, pp. 296–310, 2001.
- [7] A. M. A. El-Sayed, A. E. M. El-Mesiry, and H. A. A. El-Saka, "Numerical solution for multi-term fractional (arbitrary) orders differential equations," *Computational & Applied Mathematics*, vol. 23, no. 1, pp. 33–54, 2004.
- [8] R. Gorenflo and F. Mainardi, "Fractional calculus: integral and differential equations of fractional order," in *Fractals and Fractional Calculus in Continuum Mechanics*, vol. 378 of *CISM Courses and Lectures*, pp. 223–276, Springer, Vienna, Austria, 1997.
- [9] D. Matignon, "Stability results for fractional differential equations with applications to control processing," in *Proceedings of the Computational Engineering in System Application 2*, Lille, France, 1996.
- [10] M. De la Sen, "Positivity and stability of the solutions of Caputo fractional linear time-invariant systems of any order with internal point delays," *Abstract and Applied Analysis*, vol. 2011, Article ID 161246, 25 pages, 2011.

- [11] A. Ashyralyev, "A note on fractional derivatives and fractional powers of operators," *Journal of Mathematical Analysis and Applications*, vol. 357, no. 1, pp. 232–236, 2009.
- [12] A. Ashyralyev, F. Dal, and Z. Pinar, "A note on the fractional hyperbolic differential and difference equations," *Applied Mathematics and Computation*, vol. 217, no. 9, pp. 4654–4664, 2011.
- [13] A. Ashyralyev, F. Dal, and Z. Pinar, "On the numerical solution of fractional hyperbolic partial differential equations," *Mathematical Problems in Engineering*, vol. 2009, Article ID 730465, 11 pages, 2009.
- [14] F. Dal, "Application of variational iteration method to fractional hyperbolic partial differential equations," *Mathematical Problems in Engineering*, vol. 2009, Article ID 824385, 10 pages, 2009.
- [15] I. Podlubny and A. M. A. El-Sayed, *On Two Definitions of Fractional Calculus*, Slovak Academy of Science-Institute of Experimental Phys, 1996.
- [16] S. G. Kreĭn, *Linear Differential Equations in a Banach Space*, Nauka, Moscow, Russia, 1967.
- [17] P. E. Sobolevskii and L. M. Chebotaryeva, "Approximate solution by method of lines of the Cauchy problem for an abstract hyperbolic equations," *Izvestiya Vysshikh Uchebnykh Zavedenii, Matematika*, vol. 5, pp. 103–116, 1977 (Russian).
- [18] A. Ashyralyev, M. Martinez, J. Paster, and S. Piskarev, "Weak maximal regularity for abstract hyperbolic problems in function spaces," in *Proceedings of the 6th International ISAAC Congress*, p. 90, Ankara, Turkey, 2007.
- [19] A. Ashyralyev and N. Aggez, "A note on the difference schemes of the nonlocal boundary value problems for hyperbolic equations," *Numerical Functional Analysis and Optimization*, vol. 25, no. 5-6, pp. 439–462, 2004.
- [20] A. Ashyralyev and I. Muradov, "On difference schemes a second order of accuracy for hyperbolic equations," in *Modelling Processes of Exploitation of Gas Places and Applied Problems of Theoretical Gasohydrodynamics*, pp. 127–138, Ashgabat, Ilim, 1998.
- [21] A. Ashyralyev and P. E. Sobolevskii, *New Difference Schemes for Partial Differential Equations*, vol. 148 of *Operator Theory: Advances and Applications*, Birkhäuser, Boston, Mass, USA, 2004.
- [22] A. Ashyralyev and Y. Ozdemir, "On nonlocal boundary value problems for hyperbolic-parabolic equations," *Taiwanese Journal of Mathematics*, vol. 11, no. 4, pp. 1075–1089, 2007.
- [23] A. Ashyralyev and O. Yildirim, "On multipoint nonlocal boundary value problems for hyperbolic differential and difference equations," *Taiwanese Journal of Mathematics*, vol. 14, no. 1, pp. 165–194, 2010.
- [24] A. A. Samarskii, I. P. Gavrilyuk, and V. L. Makarov, "Stability and regularization of three-level difference schemes with unbounded operator coefficients in Banach spaces," *SIAM Journal on Numerical Analysis*, vol. 39, no. 2, pp. 708–723, 2001.
- [25] A. Ashyralyev and P. E. Sobolevskii, "Two new approaches for construction of the high order of accuracy difference schemes for hyperbolic differential equations," *Discrete Dynamics in Nature and Society*, no. 2, pp. 183–213, 2005.
- [26] A. Ashyralyev and M. E. Koksai, "On the second order of accuracy difference scheme for hyperbolic equations in a Hilbert space," *Numerical Functional Analysis and Optimization*, vol. 26, no. 7-8, pp. 739–772, 2005.
- [27] A. Ashyralyev and M. E. Koksai, "On the second order of accuracy difference scheme for hyperbolic equations in a Hilbert space," *Numerical Functional Analysis and Optimization*, vol. 26, no. 7-8, pp. 739–772, 2005.
- [28] M. Ashyraliyev, "A note on the stability of the integral-differential equation of the hyperbolic type in a Hilbert space," *Numerical Functional Analysis and Optimization*, vol. 29, no. 7-8, pp. 750–769, 2008.
- [29] A. Ashyralyev and P. E. Sobolevskii, "A note on the difference schemes for hyperbolic equations," *Abstract and Applied Analysis*, vol. 6, no. 2, pp. 63–70, 2001.
- [30] P. E. Sobolevskii, *Difference Methods for the Approximate Solution of Differential Equations*, Izdatelstvo Voronezhskogo Gosud Universiteta, Voronezh, Russia, 1975.
- [31] M. Inokti, H. Sekine, and T. Mura, "General use of the Lagrange multiplier in nonlinear mathematical physic," in *Variational Method in the Mechanics of Solids*, S. Nemer-Nasser, Ed., Pergamon Press, Oxford, UK, 1978.
- [32] J. H. He, *Generalized Variational Principles in Fluids*, Science & Culture Publishing House of China, Hong Kong, 2003.