Research Article

On the Periods of 2-Step General Fibonacci Sequences in the Generalized Quaternion Groups

Bahram Ahmadi and Hossein Doostie

*Department of Mathematics, Science and Research Branch, Islamic Azad University,
P.O. Box 14515-1775, Tehran 14778-93855, Iran*

Correspondence should be addressed to Bahram Ahmadi, b.ahmadi.math@gmail.com

Received 12 October 2012; Accepted 3 December 2012

Academic Editor: Carlo Piccardi

Copyright © 2012 B. Ahmadi and H. Doostie. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study 2-step general Fibonacci sequences in the generalized quaternion groups Q_{4n} . In cases where the sequences are proved to be simply periodic, we obtain the periods of 2-step general Fibonacci sequences.

1. Introduction

The study of the Fibonacci sequences in groups began with the earlier work of Wall [1] in 1960, where the ordinary Fibonacci sequences in cyclic groups were investigated. In the mid-eighties, Wilcox [2] extended the problem to the abelian groups. In 1990, Campbell et al. [3] expanded the theory to some classes of finite groups. In 1992, Knox proved that the periods of k -nacci (k -step Fibonacci) sequences in the dihedral groups are equal to $2k + 2$, in [4]. In the progress of this study, the article [5] of Aydin and Smith proves that the lengths of the ordinary 2-step Fibonacci sequences are equal to the lengths of the ordinary 2-step Fibonacci recurrences in finite nilpotent groups of nilpotency class 4 and a prime exponent, in 1994.

Since 1994, the theory has been generalized and many authors had nice contributions in computations of recurrence sequences in groups and we may give here a brief of these attempts. In [6, 7] the definition of the Fibonacci sequence has been generalized to the ordinary 3-step Fibonacci sequences in finite nilpotent groups. Then in [8] it is proved that the period of 2-step general Fibonacci sequence is equal to the length of the fundamental period of the 2-step general recurrence constructed by two generating elements of a group of nilpotency class 2 and exponent p . In [9] Karaduman and Yavuz showed that the periods of the 2-step Fibonacci recurrences in finite nilpotent groups of nilpotency class 5 and a prime exponent are $p \cdot k(p)$, for $2 < p \leq 2927$, where p is a prime and $k(p)$ is the period of the ordinary

2-step Fibonacci sequence. The main role of [10, 11] in generalizing the theory was to study the 2-step general Fibonacci sequences in finite nilpotent groups of nilpotency class 4 and exponent p and to the 2-step Fibonacci sequences in finite nilpotent groups of nilpotency class n and exponent p , respectively.

One may consult [12, 13] to see the results of the Fibonacci sequences in the modular groups concerning the periodicity of 2-step Fibonacci sequences constructed by two generating elements.

Going on a detailed literature in this area of research, we have to mention certain essential computation on the Fibonacci lengths of new structures like the semidirect products, the direct products, and the automorphism groups of finite groups which have been studied in [14–19]. Finally, we refer to [20] where Karaduman and Aydin studied the periodicity property of 2-step general Fibonacci sequences in dihedral groups and the goal of this paper is to calculate the periods of 2-step general Fibonacci sequences in the generalized quaternion groups.

Let $G = \langle a_0, a_1, a_2, \dots, a_{j-1} \rangle$ be a finite group. A k -nacci sequence in group G is a sequence $\{x_n\}_{n=0}^{\infty}$ of group elements for which each element is defined by $x_0 = a_0, x_1 = a_1, \dots, x_{j-1} = a_{j-1}$,

$$x_n = \begin{cases} x_0 x_1 \cdots x_{n-1}, & \text{for } j \leq n < k, \\ x_{n-k} x_{n-k+1} \cdots x_{n-1}, & \text{for } n \geq k. \end{cases} \quad (1.1)$$

This sequence of the group G is denoted by $F_k(G; a_0, a_1, a_2, \dots, a_{j-1})$. We also call a 2-nacci sequence of group elements a *Fibonacci sequence* of a finite group. A finite group G is *k-nacci sequenceable* if there exists a k -nacci sequence of G such that every element of the group appears in the sequence. A sequence of group elements is *periodic* if after a certain point, it consists only of repetitions of a fixed subsequence. The number of elements in the repeating subsequence is called the *period* of the sequence. For example, the sequence $a, b, c, d, d, e, b, c, d, e, \dots$ is periodic after the initial element a and has period 4. We denote the period of a k -nacci sequence $F_k(G; a_0, a_1, a_2, \dots, a_{j-1})$ by $P_k(G; a_0, a_1, a_2, \dots, a_{j-1})$. A sequence of group elements is called *simply periodic* with period k if the first k elements in the sequence form a repeating subsequence. For example, the sequence $a, b, c, d, e, f, a, b, c, d, e, f, \dots$ is simply periodic with period 6. The following theorem is well known.

Theorem 1.1 (see [4]). *Every k -nacci sequence in a finite group is simply periodic.*

2. Main Theorems

The generalized quaternion group Q_{4n} , ($n \geq 2$) is a group with a presentation of the form

$$\langle a, b \mid a^{2n} = 1, a^n = b^2, b^{-1}ab = a^{-1} \rangle. \quad (2.1)$$

It is easy to see that Q_{4n} is of order $4n$, a has order $2n$, b has order 4, and the relation $a^k b = b a^{-k}$ holds for all $k \in \mathbb{Z}$.

First we consider the following lemma which will be used frequently without further reference.

Lemma 2.1. For an integer r , where $1 \leq r < 2n$, and for $s \in \{1, 3\}$, the following relations hold in Q_{4n} :

- (i) $(a^r b^s)^t = b^{st}$, where t is any even integer;
- (ii) $(a^r b^s)^t = a^r b^{st}$, where t is any odd integer.

Proof. We may use an induction method on t for both parts, simultaneously. □

Theorem 2.2. $P_k(Q_{4n}; a, b) = 2k + 2$.

Proof. Obviously, the order of ab is 4. If $k = 2$, then

$$\begin{aligned} x_0 &= a, & x_1 &= b, & x_2 &= ab, & x_3 &= a^{n-1}, \\ x_4 &= a^{n+2}b, & x_5 &= ab, & x_6 &= a, & x_7 &= b, \dots, \end{aligned} \quad (2.2)$$

and hence the sequence has period 6. If $k = 3$, then

$$\begin{aligned} x_0 &= a, & x_1 &= b, & x_2 &= ab, & x_3 &= (ab)^2 = b^2, & x_4 &= a^{-1}, \\ x_5 &= a^2b^3, & x_6 &= ab, & x_7 &= 1, & x_8 &= a, & x_9 &= b, \dots, \end{aligned} \quad (2.3)$$

and so the period is 8. Now let $k \geq 4$. Then, the first $k + 1$ elements of $F_k(Q_{4n}; a, b)$ are

$$x_0 = a, \quad x_1 = b, \quad x_2 = ab, \quad x_3 = (ab)^2, \quad x_4 = (ab)^4, \dots, x_k = (ab)^{2^{k-2}}. \quad (2.4)$$

Since ab is of order 4, this sequence reduces to

$$x_0 = a, \quad x_1 = b, \quad x_2 = ab, \quad x_3 = b^2, \quad x_j = 1, \quad (4 \leq j \leq k). \quad (2.5)$$

Thus,

$$\begin{aligned} x_{k+1} &= \prod_{i=1}^k x_i = b(ab)b^2 = a^{-1}, \\ x_{k+2} &= \prod_{i=2}^{k+1} x_i = (ab)b^2a^{-1} = a^2b^3, \\ x_{k+3} &= \prod_{i=3}^{k+2} x_i = b^2a^{-1}(a^2b^3) = ab, \\ x_{k+4} &= \prod_{i=4}^{k+3} x_i = a^{-1}(a^2b^3)(ab) = 1. \end{aligned} \quad (2.6)$$

It follows that $x_{k+j} = 1$ for j ($4 \leq j \leq k+1$). We also have

$$\begin{aligned} x_{2k+2} &= \prod_{i=k+2}^{2k+1} x_i = (a^2 b^3)(ab) = a, \\ x_{2k+3} &= \prod_{i=k+3}^{2k+2} x_i = (ab)a = b. \end{aligned} \quad (2.7)$$

It shows that $P_k(Q_{4n}; a, b) = 2k + 2$. □

Let $G = \langle a, b \rangle$ be a finite nonabelian 2-generated group. A 2-step general Fibonacci sequence in the group G is defined by $x_0 = a$, $x_1 = b$, $x_i = x_{i-2}^m x_{i-1}^l$, for $i \geq 2$, and the integers m and l . Now we study this sequence for group Q_{4n} .

Theorem 2.3. *Let m and l be integers. If $m \equiv 0 \pmod{2n}$ or $m \equiv 0 \pmod{4}$, then 2-step general Fibonacci sequence in Q_{4n} is not simply periodic.*

Proof. First we consider the case $m \equiv 0 \pmod{2n}$. Then the sequence is

$$x_0 = a, \quad x_1 = b, \quad x_2 = a^m b^l = b^l, \quad x_3 = b^m b^{l^2} = b^{m+l^2}, \dots \quad (2.8)$$

Obviously, the cycle does not begin again with a and b , and hence the sequence is not simply periodic.

Now let $m \equiv 0 \pmod{4}$. Four cases occur.

Case 1 ($l \equiv 0 \pmod{4}$). Then

$$\begin{aligned} x_0 &= a, & x_1 &= b, & x_2 &= a^m, & x_3 &= a^{ml}, \\ x_4 &= a^{m^2+ml^2}, & x_5 &= a^{2m^2l+ml^3}, & x_6 &= a^{m^3+3m^2l^2+ml^4}, \dots \end{aligned} \quad (2.9)$$

Since the cycle does not begin again with a and b , the sequence is not simply periodic.

Case 2 ($l \equiv 1 \pmod{4}$). Then

$$x_0 = a, \quad x_1 = b, \quad x_2 = a^m b, \quad x_3 = a^m b, \quad x_4 = a^m b, \dots \quad (2.10)$$

Clearly, the sequence is not simply periodic.

Case 3 ($l \equiv 2 \pmod{4}$). Note that b^2 is a central element of Q_{4n} . Thus

$$x_0 = a, \quad x_1 = b, \quad x_2 = a^m b^2, \quad x_3 = a^{ml}, \quad x_4 = a^{m^2+ml^2}, \quad x_5 = a^{2m^2l+ml^3}, \dots \quad (2.11)$$

Similar to Case 1, the sequence is not simply periodic.

Case 4 ($l \equiv 3 \pmod{4}$). So

$$x_0 = a, \quad x_1 = b, \quad x_2 = a^m b^3, \quad x_3 = a^m b, \quad x_4 = a^m b^3, \quad x_5 = a^m b, \dots \quad (2.12)$$

Thus the sequence is not simply periodic. $\square$

Theorem 2.4. *Let n be an even integer.*

- (i) *If $m \equiv 1 \pmod{2n}$ and $l \equiv 0 \pmod{2n}$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period 2.*
- (ii) *If $m \equiv -1 \pmod{2n}$ and $l \equiv -1 \pmod{2n}$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period 3.*
- (iii) *If $m \equiv 1 \pmod{2n}$ and $l \equiv 1 \pmod{2n}$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period 6.*

Proof. (i) Since n is even, $m \equiv 1 \pmod{4}$ and $l \equiv 0 \pmod{4}$. Thus

$$x_0 = a, \quad x_1 = b, \quad x_2 = a, \quad x_3 = b, \dots, \quad (2.13)$$

and the period is 2.

(ii) Since n is even, $m \equiv 3 \pmod{4}$ and $l \equiv 3 \pmod{4}$. Note that m and l are odd integers. So the sequence reduces to

$$x_0 = a, \quad x_1 = b, \quad x_2 = a^{-1}b^3, \quad x_3 = a, \quad x_4 = b, \dots, \quad (2.14)$$

and the period is 3.

(iii) Because n is even, $m \equiv 1 \pmod{4}$ and $l \equiv 1 \pmod{4}$. Furthermore, m and l are odd integers. Then the sequence reduces to

$$\begin{aligned} x_0 &= a, & x_1 &= b, & x_2 &= ab, & x_3 &= a^{n-1}, \\ x_4 &= a^2b^3, & x_5 &= ab, & x_6 &= a, & x_7 &= b, \dots, \end{aligned} \quad (2.15)$$

and the period is 6. $\square$

Theorem 2.5. *Let n be an odd integer.*

- (i) *If $m \equiv 1 \pmod{2n}$, $m \equiv 1 \pmod{4}$, $l \equiv 0 \pmod{4}$ and $(l, n) = 1$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period $2n$.*
- (ii) *If $m \equiv 1 \pmod{2n}$, $m \equiv 3 \pmod{4}$, $l \equiv 0 \pmod{4}$ and $(l, n) = 1$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period $4n$.*

Proof. (i) By induction on $k \geq 0$, we can show that $x_{2k} = a$ and $x_{2k+1} = a^{-kl}b$. In particular, the period must be even. Now we have

$$x_{2k+1} = b \iff 2n \mid kl \iff n \mid k \quad (\text{because } (l, n) = 1), \quad (2.16)$$

and so the period is $2n$.

(ii) By induction on $k \geq 0$, it may be shown that

$$x_{2k} = a, \quad x_{2k+1} = \begin{cases} a^{-kl}b & \text{if } k \text{ is even,} \\ a^{-kl}b^3 & \text{if } k \text{ is odd.} \end{cases} \quad (2.17)$$

Therefore, the period must be even. Now we have

$$x_{2k+1} = b \iff 2n \mid kl \iff 2n \mid k \quad (\text{because } (l, n) = 1 \text{ and } n \text{ is odd}), \quad (2.18)$$

and thus the period is $4n$. $\square$

Theorem 2.6. Let n and l be odd integers, $m \equiv -1 \pmod{2n}$ and $(l+1, n) = 1$.

- (i) If $m \equiv 1 \pmod{4}$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period $6n$.
- (ii) If $m \equiv 3 \pmod{4}$ and $l \equiv 1 \pmod{4}$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period $6n$.
- (iii) If $m \equiv 3 \pmod{4}$ and $l \equiv 3 \pmod{4}$, then 2-step general Fibonacci sequence in Q_{4n} is simply periodic with period $3n$.

Proof. (i) Two cases occur.

Case 1 ($l \equiv 1 \pmod{4}$). By induction on $k \geq 0$, we can prove that the following relations hold:

$$\begin{aligned} x_{6k} &= a, \\ x_{6k+1} &= a^{-2k(l+1)}b, \\ x_{6k+2} &= a^{-(2k(l+1)+1)}b, \\ x_{6k+3} &= a^{n+1}, \\ x_{6k+4} &= a^{-(n+(2k+1)(l+1))}b, \\ x_{6k+5} &= a^{-((2k+1)(l+1)+1)}b. \end{aligned} \quad (2.19)$$

Consequently, the period must be a multiple of 6. Now we have

$$x_{6k+1} = b \iff 2n \mid 2k(l+1) \iff n \mid k \quad (\text{because } (l+1, n) = 1), \quad (2.20)$$

and hence the period is $6n$.

Case 2 ($l \equiv 3 \pmod{4}$). The proof is similar to Case 1 except that

$$\begin{aligned} x_{6k+2} &= a^{-(n+2k(l+1)+1)}b, \\ x_{6k+4} &= a^{-(2k+1)(l+1)}b. \end{aligned} \quad (2.21)$$

(ii) It is easily shown that the following relations hold for every $k \geq 0$:

$$\begin{aligned} x_{3k} &= a, & x_{6k} &= a, & x_{6k+1} &= a^{-2k(l+1)}b, & x_{6k+2} &= a^{-(2k(l+1)+1)}b, \\ x_{6k+3} &= a, & x_{6k+4} &= a^{-(n+(2k+1)(l+1))}b, & x_{6k+5} &= a^{-(n+1+(2k+1)(l+1))}b. \end{aligned} \quad (2.22)$$

Particularly, the period must be a multiple of 3. Since $n + (2k + 1)(l + 1)$ is odd, then it is not a multiple of $2n$. Thus $x_{6k+4} \neq b$. Further,

$$x_{6k+1} = b \iff 2n \mid 2k(l+1) \iff n \mid k \quad (\text{because } (l+1, n) = 1), \quad (2.23)$$

and hence the period is $6n$.

(iii) By induction on $k \geq 0$, we can prove the following relations:

$$\begin{aligned} x_{3k} &= a, \\ x_{3k+1} &= a^{-k(l+1)}b, \\ x_{3k+2} &= a^{-(n+1+k(l+1))}b. \end{aligned} \quad (2.24)$$

Therefore, the period must be a multiple of 3. Now we have

$$x_{3k+1} = b \iff 2n \mid k(l+1) \iff n \mid k \quad (\text{because } (l+1, n) = 1), \quad (2.25)$$

and so the period is $3n$. □

Acknowledgments

The authors would like to express their appreciations to the referees for the valuable and constructive comments regarding the presentation of this paper.

References

- [1] D. D. Wall, "Fibonacci series modulo m ," *The American Mathematical Monthly*, vol. 67, pp. 525–532, 1960.
- [2] H. J. Wilcox, "Fibonacci sequences of period n in groups," *The Fibonacci Quarterly*, vol. 24, no. 4, pp. 356–361, 1986.
- [3] C. M. Campbell, H. Doostie, and E. F. Robertson, "Fibonacci length of generating pairs in groups," in *Applications of Fibonacci Numbers*, vol. 3, pp. 27–35, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1990.
- [4] S. W. Knox, "Fibonacci sequences in finite groups," *The Fibonacci Quarterly*, vol. 30, no. 2, pp. 116–120, 1992.
- [5] H. Aydin and G. C. Smith, "Finite p -quotients of some cyclically presented groups," *Journal of the London Mathematical Society*, vol. 49, no. 1, pp. 83–92, 1994.
- [6] R. Dikici and G. C. Smith, "Fibonacci sequences in finite nilpotent groups," *Turkish Journal of Mathematics*, vol. 21, no. 2, pp. 133–142, 1997.
- [7] R. Dikici and G. C. Smith, "Recurrences in finite groups," *Turkish Journal of Mathematics*, vol. 19, no. 3, pp. 321–329, 1995.

- [8] H. Aydin and R. Dikici, "General Fibonacci sequences in finite groups," *The Fibonacci Quarterly*, vol. 36, no. 3, pp. 216–221, 1998.
- [9] E. Karaduman and U. Yavuz, "On the period of Fibonacci sequences in nilpotent groups," *Applied Mathematics and Computation*, vol. 142, no. 2-3, pp. 321–332, 2003.
- [10] E. Karaduman and H. Aydin, "General 2-step Fibonacci sequences in nilpotent groups of exponent p and nilpotency class 4," *Applied Mathematics and Computation*, vol. 141, no. 2-3, pp. 491–497, 2003.
- [11] E. Karaduman and H. Aydin, "On Fibonacci sequences in nilpotent groups," *Mathematica Balkanica*, vol. 17, no. 3-4, pp. 207–214, 2003.
- [12] R. Dikici and A. Işik, "Fibonacci sequences in modular groups," *Institute of Mathematics & Computer Sciences*, vol. 10, no. 3, pp. 187–189, 1997.
- [13] R. Dikici and E. Özkan, "General recurrences in modular groups," *Journal of the Faculty of Science*, vol. 21, no. 2, pp. 93–101, 1998.
- [14] H. Doostie and R. Golamie, "Computing on the Fibonacci lengths of finite groups," *International Journal of Applied Mathematics*, vol. 4, no. 2, pp. 149–156, 2000.
- [15] H. Doostie and M. Maghasedi, "Fibonacci length of direct products of groups," *Vietnam Journal of Mathematics*, vol. 33, no. 2, pp. 189–197, 2005.
- [16] H. Doostie and C. M. Campbell, "Fibonacci length of automorphism groups involving Tribonacci numbers," *Vietnam Journal of Mathematics*, vol. 28, no. 1, pp. 57–65, 2000.
- [17] C. M. Campbell and P. P. Campbell, "The Fibonacci length of certain centro-polyhedral groups," *Journal of Applied Mathematics & Computing*, vol. 19, no. 1-2, pp. 231–240, 2005.
- [18] C. M. Campbell, P. P. Campbell, H. Doostie, and E. F. Robertson, "Fibonacci lengths for certain metacyclic groups," *Algebra Colloquium*, vol. 11, no. 2, pp. 215–222, 2004.
- [19] H. Doostie and M. Hashemi, "Fibonacci lengths involving the Wall number $k(n)$," *Journal of Applied Mathematics & Computing*, vol. 20, no. 1-2, pp. 171–180, 2006.
- [20] E. Karaduman and H. Aydin, "On the periods of 2-step general Fibonacci sequences in dihedral groups," *Matematichki Vesnik*, vol. 58, no. 1-2, pp. 47–56, 2006.