Research Article

Some Identities on Bernoulli and Euler Numbers

D. S. Kim,¹ T. Kim,² J. Choi,³ and Y. H. Kim³

¹ Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

² Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

³ Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to T. Kim, tkkim@kw.ac.kr

Received 15 November 2011; Accepted 23 December 2011

Academic Editor: Delfim F. M. Torres

Copyright © 2012 D. S. Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently, Kim introduced the fermionic p -adic integral on $\mathbb{Z}_p$. By using the equations of the fermionic and bosonic p -adic integral on $\mathbb{Z}_p$, we give some interesting identities on Bernoulli and Euler numbers.

1. Introduction/Preliminaries

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will denote the ring of p -adic integers, the field of p -adic rational numbers, and the completion of algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. The p -adic absolute value $|\cdot|_p$ is normally defined by $|p|_p = 1/p$.

Let $\text{UD}(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$ and $C(\mathbb{Z}_p)$ the space of continuous function on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the fermionic p -adic integral on $\mathbb{Z}_p$ is defined by Kim as follows:

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x, \quad (\text{see [1]}). \quad (1.1)$$

The following fermionic p -adic integral equation on $\mathbb{Z}_p$ is well known (see [1–3]):

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0), \quad (1.2)$$

where $f_1(x) = f(x+1)$.

From (1.1) and (1.2), we can derive the generating function of Euler polynomials as follows:

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (1.3)$$

where $E_n(x)$ is the n th ordinary Euler polynomial (see [1–4]). In the special case, $x = 0$, $E_n(0) = E_n$ is called the n th ordinary Euler number.

By (1.3), we get Witt's formula for the n th Euler polynomial as follows:

$$\int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(y) = E_n(x), \quad \text{for } n \in \mathbb{Z}_+. \quad (1.4)$$

Thus, by (1.4), we have

$$E_n(x) = (E+x)^n = \sum_{l=0}^n \binom{n}{l} x^{n-l} E_l, \quad (1.5)$$

with the usual convention about replacing E^n by E_n (see [5, 6]). From (1.3), we note that

$$(E+1)^n + E_n = 2\delta_{0,n}, \quad (1.6)$$

where $\delta_{k,n}$ is the Kronecker symbol (see [3]). By (1.2) and (1.4), we get

$$\int_{\mathbb{Z}_p} (x+y+1)^n d\mu_{-1}(y) + \int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(y) = 2x^n. \quad (1.7)$$

Thus, by (1.4) and (1.7), we have

$$E_n(x+1) + E_n(x) = 2x^n, \quad \text{for } n \in \mathbb{Z}_+. \quad (1.8)$$

Equation (1.8) is equivalent to

$$x^n = E_n(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_l(x). \quad (1.9)$$

From (1.6), we can derive the following equation:

$$E_n(2) = 2 - E_n(1) = 2 + E_n - 2\delta_{0,n}, \quad \text{for } n \in \mathbb{Z}_+. \quad (1.10)$$

For $f \in \text{UD}(\mathbb{Z}_p)$, the bosonic p -adic integral on $\mathbb{Z}_p$ is defined by

$$I_1(f) = \int_{\mathbb{Z}_p} f(x) d\mu_1(x) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x), \quad (\text{see [4]}). \quad (1.11)$$

From (1.11), we can easily derive the following I_1 -integral equation:

$$I_1(f_1) = I(f) + f'(0), \quad (\text{see } [4, 7, 8]), \quad (1.12)$$

where $f_1(x) = f(x+1)$ and $f'(0) = df(x)/dx|_{x=0}$.

It is well known that the Bernoulli polynomial can be represented by the bosonic p -adic integral on $\mathbb{Z}_p$ as follows:

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_1(y) = \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}, \quad (1.13)$$

where $B_n(x)$ is called the n th Bernoulli polynomial (see [4, 7–13]). In the special case, $x = 0$, $B_n(0) = B_n$ is called the n th Bernoulli number. By the definition of Bernoulli numbers and polynomials, we get

$$B_n(x) = \int_{\mathbb{Z}_p} (x+y)^n d\mu_1(y) = \sum_{l=0}^n \binom{n}{l} x^{n-l} B_l. \quad (1.14)$$

Thus, by (1.13) and (1.14), we see that

$$B_0 = 1, \quad (B+1)^n - B_n = \delta_{1,n}, \quad (1.15)$$

with the usual convention about replacing B^n by B_n (see [1–22]).

By (1.11), we easily get

$$\int_{\mathbb{Z}_p} (1-x+y)^n d\mu_1(y) = (-1)^n \int_{\mathbb{Z}_p} (x+y)^n d\mu_1(y). \quad (1.16)$$

From (1.13), (1.14), and (1.16), we have

$$B_n(1-x) = (-1)^n B_n(x) \quad \text{for } n \in \mathbb{Z}_+. \quad (1.17)$$

By (1.15), we get

$$B_n(2) = n + B_n(1) = n + B_n + \delta_{1,n}. \quad (1.18)$$

Thus, by (1.17) and (1.18), we have

$$(-1)^n B_n(-1) = B_n(2) = n + B_n + \delta_{1,n}, \quad (\text{see } [4]). \quad (1.19)$$

From (1.12) and (1.13), we get

$$\int_{\mathbb{Z}_p} (x+1+y)^{n+1} d\mu_1(y) - \int_{\mathbb{Z}_p} (x+y)^{n+1} d\mu_1(y) = (n+1)x^n. \quad (1.20)$$

Thus, by (1.13) and (1.20), we have

$$B_{n+1}(x+1) - B_{n+1}(x) = (n+1)x^n \quad \text{for } n \in \mathbb{Z}_+. \quad (1.21)$$

Equation (1.21) is equivalent to the following equation:

$$x^n = \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} B_l(x) \quad \text{for } n \in \mathbb{Z}_+. \quad (1.22)$$

In this paper we derive some interesting and new identities for the Bernoulli and Euler numbers from the p -adic integral equations on $\mathbb{Z}_p$.

2. Some Identities on Bernoulli and Euler Numbers

From (1.1), we note that

$$\int_{\mathbb{Z}_p} (1-x+y)^n d\mu_{-1}(y) = (-1)^n \int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(y). \quad (2.1)$$

By (1.14) and (2.1), we get

$$E_n(1-x) = (-1)^n E_n(x), \quad \text{where } n \in \mathbb{Z}_+. \quad (2.2)$$

In the special case, $x = -1$, we have

$$E_n(2) = (-1)^n E_n(-1) = 2 + E_n - 2\delta_{0,n}. \quad (2.3)$$

Let us consider the following fermionic p -adic integral on $\mathbb{Z}_p$ as follows:

$$\begin{aligned} \int_{\mathbb{Z}_p} x^n d\mu_{-1}(x) &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \int_{\mathbb{Z}_p} B_l(x) d\mu_{-1}(x) \\ &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} \int_{\mathbb{Z}_p} x^k d\mu_{-1}(x) \\ &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} E_k. \end{aligned} \quad (2.4)$$

Therefore, by (1.4) and (2.4), we obtain the following theorem.

Theorem 2.1. *For $n \in \mathbb{Z}_+$, one has*

$$E_n = \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} E_k. \quad (2.5)$$

It is known that $B_n(x) = (-1)^n B_n(1-x)$. If we take the fermionic p -adic integral on both sides of (1.22), then we have

$$\begin{aligned}
 \int_{\mathbb{Z}_p} x^n d\mu_{-1}(x) &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \int_{\mathbb{Z}_p} B_l(x) d\mu_{-1}(x) \\
 &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \int_{\mathbb{Z}_p} B_l(1-x) d\mu_{-1}(x) \\
 &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} B_{l-k} \int_{\mathbb{Z}_p} (1-x)^k d\mu_{-1}(x) \\
 &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} B_{l-k} (-1)^k E_k(-1).
 \end{aligned} \tag{2.6}$$

From (2.2) and (2.6), we note that

$$\begin{aligned}
 \int_{\mathbb{Z}_p} x^n d\mu_{-1}(x) &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} B_{l-k} E_k(2) \\
 &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} B_{l-k} (2 + E_k - 2\delta_{0,k}) \\
 &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \left(2B_l(1) + \sum_{k=0}^l \binom{l}{k} B_{l-k} E_k - 2B_l \right) \\
 &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \left(\sum_{k=0}^l \binom{l}{k} B_{l-k} E_k + 2\delta_{1,l} \right).
 \end{aligned} \tag{2.7}$$

Therefore, by (1.4) and (2.7), we obtain the following theorem.

Theorem 2.2. For $n \in \mathbb{Z}_+$, one has

$$E_n = \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \left(\sum_{k=0}^l \binom{l}{k} B_{l-k} E_k + 2\delta_{1,l} \right). \tag{2.8}$$

Corollary 2.3. For $n \in \mathbb{N}$, one has

$$2 + E_n = \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} (-1)^l \left(\sum_{k=0}^l \binom{l}{k} B_{l-k} E_k \right). \tag{2.9}$$

Let us take the bosonic p -adic integral on both sides of (1.9) as follows:

$$\begin{aligned}
\int_{\mathbb{Z}_p} x^n d\mu_1(x) &= \int_{\mathbb{Z}_p} \left(E_n(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_l(x) \right) d\mu_1(x) \\
&= \sum_{l=0}^n \binom{n}{l} E_{n-l} \int_{\mathbb{Z}_p} x^l d\mu_1(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} \int_{\mathbb{Z}_p} x^k d\mu_1(x) \quad (2.10) \\
&= \sum_{l=0}^n \binom{n}{l} E_{n-l} B_l + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} B_k.
\end{aligned}$$

Thus, by (1.14) and (2.10), we obtain the following theorem.

Theorem 2.4. *For $n \in \mathbb{Z}_+$, one has*

$$B_n = \sum_{l=0}^n \binom{n}{l} E_{n-l} B_l + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} B_k. \quad (2.11)$$

On the other hand, by (2.2) and (2.10), we get

$$\begin{aligned}
\int_{\mathbb{Z}_p} x^n d\mu_1(x) &= (-1)^n \int_{\mathbb{Z}_p} E_n(1-x) d\mu_1(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \int_{\mathbb{Z}_p} E_l(1-x) d\mu_1(x) \\
&= (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} \int_{\mathbb{Z}_p} (1-x)^l d\mu_1(x) \\
&\quad + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} \int_{\mathbb{Z}_p} (1-x)^k d\mu_1(x) \\
&= (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} (-1)^l B_l (-1) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} (-1)^k B_k (-1) \\
&= (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} B_l (2) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} B_k (2) \\
&= (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} (l + B_l + \delta_{1,l}) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} (k + B_k + \delta_{1,k})
\end{aligned}$$

$$\begin{aligned}
&= (-1)^n n E_{n-1}(1) + (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} B_l + (-1)^n n E_{n-1} + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l l E_{l-1}(1) \\
&\quad + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} B_k + \frac{1}{2} \sum_{l=1}^{n-1} \binom{n}{l} (-1)^l l E_{l-1} \\
&= (-1)^n n (2 + E_{n-1} - 2\delta_{0,n-1}) + (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} B_l + (-1)^n n E_{n-1} \\
&\quad + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l l (2 + E_{l-1} - \delta_{0,l-1}) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} B_k \\
&\quad + \frac{1}{2} \sum_{l=1}^{n-1} \binom{n}{l} (-1)^l l E_{l-1},
\end{aligned} \tag{2.12}$$

where $n \in \mathbb{N}$ with $n \geq 2$. Therefore, by (2.12), we obtain the following theorem.

Theorem 2.5. For $n \in \mathbb{N}$ with $n \geq 2$, one has

$$\begin{aligned}
B_{2n-1} &= -\frac{2n-1}{2} - (2n-1)E_{2n-2}(-1) - \sum_{l=0}^{2n-1} \binom{2n-1}{l} E_{2n-1-l} B_l \\
&\quad + \frac{1}{2} \sum_{l=0}^{2n-2} \binom{2n-1}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} B_k.
\end{aligned} \tag{2.13}$$

By (1.9) and (1.22), we get

$$\begin{aligned}
&\iint_{\mathbb{Z}_p} x^m y^n d\mu_{-1}(x) d\mu_1(y) \\
&= \iint_{\mathbb{Z}_p} \left(\frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} B_k(x) \right) \left(E_n(y) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_l(y) \right) d\mu_{-1}(x) d\mu_1(y) \\
&= \frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} \iint_{\mathbb{Z}_p} B_k(x) E_n(y) d\mu_{-1}(x) d\mu_1(y) \\
&\quad + \frac{1}{2(m+1)} \sum_{k=0}^m \sum_{l=0}^{n-1} \binom{m+1}{k} \binom{n}{l} \iint_{\mathbb{Z}_p} B_k(x) E_l(y) d\mu_{-1}(x) d\mu_1(y) \\
&= \frac{1}{m+1} \sum_{k=0}^m \sum_{l=0}^k \sum_{p=0}^n \binom{m+1}{k} \binom{k}{l} \binom{n}{p} B_{k-l} E_{n-p} B_p E_l \\
&\quad + \frac{1}{2(m+1)} \sum_{k=0}^m \sum_{l=0}^{n-1} \sum_{s=0}^k \sum_{p=0}^l \binom{m+1}{k} \binom{n}{l} \binom{k}{s} \binom{l}{p} B_{k-s} E_{l-p} E_s B_p.
\end{aligned} \tag{2.14}$$

Therefore, by (1.4), (1.14), and (2.14), we obtain the following theorem.

Theorem 2.6. For $m \in \mathbb{Z}_+$ and $n \in \mathbb{N}$, one has

$$\begin{aligned} E_m B_n &= \frac{1}{m+1} \sum_{k=0}^m \sum_{l=0}^k \sum_{p=0}^n \binom{m+1}{k} \binom{k}{l} \binom{n}{p} B_{k-l} E_{n-p} B_p E_l \\ &\quad + \frac{1}{2(m+1)} \sum_{k=0}^m \sum_{l=0}^{n-1} \sum_{s=0}^k \sum_{p=0}^l \binom{m+1}{k} \binom{n}{l} \binom{k}{s} \binom{l}{p} B_{k-s} E_{l-p} E_s B_p. \end{aligned} \quad (2.15)$$

It is easy to show that

$$\begin{aligned} \int_{\mathbb{Z}_p} x^{m+n} d\mu_{-1}(x) &= \int_{\mathbb{Z}_p} \left(\frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} B_k(x) \right) \left(E_n(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_l(x) \right) d\mu_{-1}(x) \\ &= \frac{1}{m+1} \sum_{k=0}^m \sum_{i=0}^k \sum_{j=0}^n \binom{m+1}{k} \binom{k}{i} \binom{n}{j} B_{k-i} E_{n-j} \int_{\mathbb{Z}_p} x^{i+j} d\mu_{-1}(x) \\ &\quad + \frac{1}{2(m+1)} \sum_{k=0}^m \sum_{l=0}^{n-1} \sum_{i=0}^k \sum_{j=0}^l \binom{m+1}{k} \binom{n}{l} \binom{k}{i} \binom{l}{j} B_{k-i} E_{l-j} \int_{\mathbb{Z}_p} x^{i+j} d\mu_{-1}(x) \\ &= \frac{1}{m+1} \sum_{k=0}^m \sum_{i=0}^k \sum_{j=0}^n \binom{m+1}{k} \binom{k}{i} \binom{n}{j} B_{k-i} E_{n-j} E_{i+j} \\ &\quad + \frac{1}{2(m+1)} \sum_{k=0}^m \sum_{l=0}^{n-1} \sum_{i=0}^k \sum_{j=0}^l \binom{m+1}{k} \binom{n}{l} \binom{k}{i} \binom{l}{j} B_{k-i} E_{l-j} E_{i+j}. \end{aligned} \quad (2.16)$$

Therefore, by (2.16), we obtain the following corollary.

Corollary 2.7. For $m \in \mathbb{Z}_+$ and $n \in \mathbb{N}$, one has

$$\begin{aligned} E_{m+n} &= \frac{1}{m+1} \sum_{k=0}^m \sum_{i=0}^k \sum_{j=0}^n \binom{m+1}{k} \binom{k}{i} \binom{n}{j} B_{k-i} E_{n-j} E_{i+j} \\ &\quad + \frac{1}{2(m+1)} \sum_{k=0}^m \sum_{l=0}^{n-1} \sum_{i=0}^k \sum_{j=0}^l \binom{m+1}{k} \binom{n}{l} \binom{k}{i} \binom{l}{j} B_{k-i} E_{l-j} E_{i+j}. \end{aligned} \quad (2.17)$$

For $f \in C(\mathbb{Z}_p)$, p -adic analogue of Bernstein operator of order n for f is given by

$$\mathbb{B}_n(f | x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x), \quad (2.18)$$

where $B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}$ for $n, k \in \mathbb{Z}_+$ is called the Bernstein polynomial of degree n (see [8]). From the definition of $B_{k,n}(x)$, we note that $B_{n-k,n}(1-x) = B_{k,n}(x)$.

Let us take the fermionic p -adic integral on $\mathbb{Z}_p$ for the product of x^m and $B_{k,n}(x)$ as follows:

$$\begin{aligned}
 \int_{\mathbb{Z}_p} x^m B_{k,n}(x) d\mu_{-1}(x) &= \frac{1}{m+1} \sum_{l=0}^m \binom{m+1}{l} \int_{\mathbb{Z}_p} B_l(x) B_{k,n}(x) d\mu_{-1}(x) \\
 &= \frac{\binom{n}{k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l \binom{m+1}{l} \binom{l}{j} B_{l-j} \int_{\mathbb{Z}_p} x^{j+k} (1-x)^{n-k} d\mu_{-1}(x) \\
 &= \frac{\binom{n}{k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^{n-k} (-1)^i B_{l-j} \binom{m+1}{l} \binom{l}{j} \binom{n-k}{i} \int_{\mathbb{Z}_p} x^{i+j+k} d\mu_{-1}(x) \\
 &= \frac{\binom{n}{k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^{n-k} (-1)^i \binom{m+1}{l} \binom{l}{j} \binom{n-k}{i} B_{l-j} E_{i+j+k}.
 \end{aligned} \tag{2.19}$$

From (2.18), we note that

$$\begin{aligned}
 \int_{\mathbb{Z}_p} x^m B_{k,n}(x) d\mu_{-1}(x) &= \binom{n}{k} \int_{\mathbb{Z}_p} x^{m+k} (1-x)^{n-k} d\mu_{-1}(x) \\
 &= \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j \int_{\mathbb{Z}_p} x^{m+k+j} d\mu_{-1}(x) \\
 &= \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j E_{m+k+j}.
 \end{aligned} \tag{2.20}$$

Therefore, by (2.19) and (2.20), we obtain the following theorem.

Theorem 2.8. For $m, n, k \in \mathbb{Z}_+$, one has

$$\sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j E_{m+k+j} = \frac{1}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^{n-k} (-1)^i \binom{m+1}{l} \binom{l}{j} \binom{n-k}{i} B_{l-j} E_{i+j+k}. \tag{2.21}$$

In particular,

$$(m+1) E_{m+n} = \sum_{l=0}^m \sum_{j=0}^l \binom{m+1}{l} \binom{l}{j} B_{l-j} E_{j+n}. \tag{2.22}$$

By (1.17) and the symmetric property of $B_{k,n}(x)$, we get

$$\begin{aligned}
 \int_{\mathbb{Z}_p} x^m B_{k,n}(x) d\mu_{-1}(x) &= \int_{\mathbb{Z}_p} x^m B_{n-k,n}(1-x) d\mu_{-1}(x) \\
 &= \frac{1}{m+1} \sum_{l=0}^m (-1)^l \binom{m+1}{l} \int_{\mathbb{Z}_p} B_l(1-x) B_{n-k,n}(1-x) d\mu_{-1}(x) \\
 &= \frac{\binom{n}{k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^k (-1)^{i+l} \binom{m+1}{l} \binom{l}{j} \binom{k}{i} B_{l-j} \int_{\mathbb{Z}_p} (1-x)^{i+j+n-k} d\mu_{-1}(x).
 \end{aligned} \tag{2.23}$$

From (1.4) and (2.2), we note that

$$\int_{\mathbb{Z}_p} (1-x)^n d\mu_{-1}(x) = (-1)^n E_n(-1) = E_n(2) = 2 + E_n - 2\delta_{0,n}. \tag{2.24}$$

By (2.23) and (2.24), we see that

$$\int_{\mathbb{Z}_p} x^m B_{k,n}(x) d\mu_{-1}(x) = \frac{\binom{n}{k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^k (-1)^{i+l} \binom{m+1}{l} \binom{l}{j} \binom{k}{i} B_{l-j} (2 + E_{i+j+n-k} - 2\delta_{0,i+j+n-k}). \tag{2.25}$$

From (2.20) and (2.25), we have

$$\begin{aligned}
 &\sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j E_{m+k+j} \\
 &= \frac{2\delta_{0,k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l (-1)^l \binom{m+1}{l} \binom{l}{j} B_{l-j} - \frac{2}{m+1} \sum_{l=0}^m (-1)^l \binom{m+1}{l} B_l \delta_{k,n} \\
 &\quad + \frac{1}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^k (-1)^{i+l} \binom{m+1}{l} \binom{l}{j} \binom{k}{i} B_{l-j} E_{i+j+n-k} \\
 &= \frac{2\delta_{0,k}}{m+1} \sum_{l=0}^m \sum_{j=0}^l (-1)^l \binom{m+1}{l} \binom{l}{j} B_{l-j} - \frac{2}{m+1} (B_{m+1}(2) + (-1)^m B_{m+1}) \delta_{k,n} \\
 &\quad + \frac{1}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^k (-1)^{i+l} \binom{m+1}{l} \binom{l}{j} \binom{k}{i} B_{l-j} E_{i+j+n-k}.
 \end{aligned} \tag{2.26}$$

Therefore, by (1.19) and (2.26), we obtain the following theorem.

Theorem 2.9. For $m, n, k \in \mathbb{N}$ with $n \geq k$, one has

$$\sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j E_{m+k+j} = \frac{1}{m+1} \sum_{l=0}^m \sum_{j=0}^l \sum_{i=0}^k (-1)^{i+l} \binom{m+1}{l} \binom{l}{j} \binom{k}{i} B_{l-j} E_{i+j+n-k} \\ - \frac{2}{m+1} (B_{m+1} + m + 1 + (-1)^m B_{m+1}). \quad (2.27)$$

In particular,

$$(2m+2)(E_{2m+n+1} + 2) = \sum_{l=0}^{2m+1} \sum_{j=0}^l \sum_{i=0}^n (-1)^{i+l} \binom{2m+2}{l} \binom{l}{j} \binom{n}{i} B_{l-j} E_{i+j}. \quad (2.28)$$

Acknowledgment

The first author was supported by National Research Foundation of Korea Grant funded by the Korean Government 2011-0002486.

References

- [1] T. Kim, "Euler numbers and polynomials associated with zeta functions," *Abstract and Applied Analysis*, vol. 2008, Article ID 581582, 11 pages, 2008.
- [2] T. Kim, "Note on the Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 17, no. 2, pp. 131–136, 2008.
- [3] T. Kim, "Symmetry of power sum polynomials and multivariate fermionic p -adic invariant integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 1, pp. 93–96, 2009.
- [4] T. Kim, "Symmetry p -adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials," *Journal of Difference Equations and Applications*, vol. 14, no. 12, pp. 1267–1277, 2008.
- [5] A. Bayad, "Modular properties of elliptic Bernoulli and Euler functions," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 389–401, 2010.
- [6] D. Ding and J. Yang, "Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 1, pp. 7–21, 2010.
- [7] T. Kim, " q -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [8] T. Kim, "A note on q -Bernstein polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 1, pp. 73–82, 2011.
- [9] A. Kudo, "A congruence of generalized Bernoulli number for the character of the first kind," *Advanced Studies in Contemporary Mathematics*, vol. 2, pp. 1–8, 2000.
- [10] Q.-M. Luo and F. Qi, "Relationships between generalized Bernoulli numbers and polynomials and generalized Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 7, no. 1, pp. 11–18, 2003.
- [11] Q.-M. Luo, "Some recursion formulae and relations for Bernoulli numbers and Euler numbers of higher order," *Advanced Studies in Contemporary Mathematics*, vol. 10, no. 1, pp. 63–70, 2005.
- [12] H. Ozden, I. N. Cangul, and Y. Simsek, "Remarks on q -Bernoulli numbers associated with Daehee numbers," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 1, pp. 41–48, 2009.
- [13] Y.-H. Kim and K.-W. Hwang, "Symmetry of power sum and twisted Bernoulli polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 2, pp. 127–133, 2009.
- [14] G. Kim, B. Kim, and J. Choi, "The DC algorithm for computing sums of powers of consecutive integers and Bernoulli numbers," *Advanced Studies in Contemporary Mathematics*, vol. 17, no. 2, pp. 137–145, 2008.
- [15] L. C. Jang, "A note on Kummer congruence for the Bernoulli numbers of higher order," *Proceedings of the Jangjeon Mathematical Society*, vol. 5, no. 2, pp. 141–146, 2002.

- [16] L. C. Jang and H. K. Pak, "Non-Archimedean integration associated with q -Bernoulli numbers," *Proceedings of the Jangjeon Mathematical Society*, vol. 5, no. 2, pp. 125–129, 2002.
- [17] S.-H. Rim, J.-H. Jin, E.-J. Moon, and S.-J. Lee, "Some identities on the q -Genocchi polynomials of higher-order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *International Journal of Mathematics and Mathematical Sciences*, vol. 2010, Article ID 860280, 14 pages, 2010.
- [18] C. S. Ryoo, "On the generalized Barnes type multiple q -Euler polynomials twisted by ramified roots of unity," *Proceedings of the Jangjeon Mathematical Society*, vol. 13, no. 2, pp. 255–263, 2010.
- [19] C. S. Ryoo, "Some relations between twisted q -Euler numbers and Bernstein polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 2, pp. 217–223, 2011.
- [20] Y. Simsek, "Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 251–278, 2008.
- [21] I. Buyukyazici, "On generalized q -Bernstein polynomials," *The Global Journal of Pure and Applied Mathematics*, vol. 6, pp. 1331–1348, 2010.
- [22] L.-C. Jang, W.-J. Kim, and Y. Simsek, "A study on the p -adic integral representation on $\mathbb{Z}_p$ associated with Bernstein and Bernoulli polynomials," *Advances in Difference Equations*, vol. 2010, Article ID 163217, 6 pages, 2010.