Research Article

On Reciprocal Series of Generalized Fibonacci Numbers with Subscripts in Arithmetic Progression

Neşe Ömür

Department of Mathematics, Kocaeli University, 41380 Izmit, Turkey

Correspondence should be addressed to Neşe Ömür, neseomur@kocaeli.edu.tr

Received 24 October 2011; Accepted 16 November 2011

Academic Editor: Elena Braverman

Copyright © 2012 Neşe Ömür. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate formulas for closely related series of the forms: $\sum_{n=0}^{\infty} 1/(U_{an+b} + c)$, $\sum_{n=0}^{\infty} (-1)^n U_{an+b}/(U_{an+b} + c)^2$, $\sum_{n=0}^{\infty} U_{2(an+b)}/(U_{an+b}^2 + c)^2$ for certain values of a , b , and c .

1. Introduction

Let p be a nonzero integer such that $\Delta = p^2 + 4 \neq 0$. The generalized Fibonacci and Lucas sequences are defined by the following recurrences:

$$\begin{aligned}U_{n+1} &= pU_n + U_{n-1}, \\V_{n+1} &= pV_n + V_{n-1},\end{aligned}\tag{1.1}$$

where $U_0 = 0$, $U_1 = 1$ and $V_0 = 2$, $V_1 = p$, respectively. When $p = 1$, $U_n = F_n$ (n th Fibonacci number) and $V_n = L_n$ (n th Lucas number).

If α and β are the roots of equation $x^2 - px - 1 = 0$, the Binet formulas of the sequences $\{U_n\}$ and $\{V_n\}$ have the forms:

$$U_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad V_n = \alpha^n + \beta^n,\tag{1.2}$$

respectively.

In [1], Backstrom developed formulas for closely related series of the form:

$$\sum_{n=0}^{\infty} \frac{1}{F_{an+b} + c},\tag{1.3}$$

for certain values of a , b , and c . For example, he obtained the following series:

$$\sum_{n=0}^{\infty} \frac{1}{F_{2n+1} + F_K} = \frac{K\sqrt{5}}{2L_K},$$

$$\sum_{n=0}^{\infty} \frac{1}{F_{(2n+1)K+2t} + F_K} = \begin{cases} \frac{(\sqrt{5} - 5F_t/L_t)}{2L_K}, & t \text{ even,} \\ \frac{(\sqrt{5} - L_t/F_t)}{2L_K}, & t \text{ odd,} \end{cases} \quad (1.4)$$

where K represents an odd integer and t is an integer in the range $-(K-1)/2$ to $(K-1)/2$ inclusive. Also, he gave the similar results for Lucas numbers.

In [2], Popov found in explicit form series of the form:

$$\sum_{n=0}^{\infty} \frac{1}{F_{an+b} \pm c}, \quad \sum_{n=0}^{\infty} \frac{1}{F_{an+b} F_{cn+d}}, \quad \sum_{n=0}^{\infty} \frac{1}{F_{an+b}^2 \pm F_{cn+d}^2}, \quad (1.5)$$

for certain values of a , b , c , and d .

In [3], Popov generalized some formulas of Backstrom [1] related to sums of reciprocal series of Fibonacci and Lucas numbers. For example,

$$\Delta \sum_{n=0}^{\infty} \frac{(-q)^{s+nr}}{V_{(2n+1)r+2s} - (-q)^{s+nr} V_r} = \begin{cases} \frac{\beta^s}{U_r U_s}, & \left| \frac{\beta}{\alpha} \right| < 1, \\ \frac{\alpha^s}{U_r U_s}, & \left| \frac{\alpha}{\beta} \right| < 1, \end{cases} \quad (1.6)$$

where s and r are integers.

In [4], Gauthier found the closed form expressions for the following sums:

$$\sum_{k=1}^m \frac{(-1)^{kn} f_{(2k+1)n}}{f_{(k+1)n}^2 f_{kn}^2}, \quad m, n \geq 1,$$

$$\sum_{k=0}^m \frac{(-1)^{kn} f_{(2k+1)n}}{l_{(k+1)n}^2 l_{kn}^2}, \quad m, n \geq 0, \quad (1.7)$$

where for $x \neq 0$ an indeterminate, the generalized Fibonacci and Lucas polynomials $\{f_n\}_n$ and $\{l_n\}_n$ are given by the following recurrences:

$$f_{n+2} = x f_{n+1} + f_n, \quad f_0 = 0, \quad f_1 = 1, \quad n \geq 0, \quad (1.8)$$

$$l_{n+2} = x l_{n+1} + l_n, \quad l_0 = 2, \quad l_1 = x, \quad n \geq 0,$$

respectively.

In this paper, we investigate formulas for closely related series of the forms:

$$\sum_{n=0}^{\infty} \frac{1}{U_{an+b} + c}, \quad \sum_{n=0}^{\infty} \frac{(-1)^n U_{an+b}}{(U_{an+b} + c)^2}, \quad \sum_{n=0}^{\infty} \frac{U_{2(an+b)}}{(U_{an+b}^2 + c)^2}, \quad (1.9)$$

for certain values of a , b and c .

2. On Some Series of Reciprocals of Generalized Fibonacci Numbers

In this section, firstly, we will give the following lemmas for further use.

Lemma 2.1. *Let n be an arbitrary nonzero integer. For integer $m \geq 1$,*

$$\sum_{k=1}^m \frac{(-1)^{kn} U_{(2k+1)n}}{U_{(k+1)n}^2 U_{kn}^2} = \frac{1}{4U_n} \left(\frac{V_n^2}{U_n^2} - \frac{V_{(m+1)n}^2}{U_{(m+1)n}^2} \right), \quad (2.1)$$

and for integer $m \geq 0$,

$$\sum_{k=0}^m \frac{(-1)^{kn} U_{(2k+1)n}}{V_{(k+1)n}^2 V_{kn}^2} = \frac{U_{(m+1)n}^2}{4U_n V_{(m+1)n}^2}. \quad (2.2)$$

Proof. We give the proof of Lemma 2.1 as the proofs of the sums in [4], using the following equalities:

$$\begin{aligned} \frac{U_{(2k+1)n}}{U_{(k+1)n} U_{kn}} &= \frac{1}{2} \left(\frac{V_{kn}}{U_{kn}} + \frac{V_{(k+1)n}}{U_{(k+1)n}} \right), \\ \frac{U_{(2k+1)n}}{V_{(k+1)n} V_{kn}} &= \frac{1}{2} \left(\frac{U_{(k+1)n}}{V_{(k+1)n}} + \frac{U_{kn}}{V_{kn}} \right), \\ \frac{(-1)^{kn} U_n}{U_{(k+1)n} U_{kn}} &= \frac{1}{2} \left(\frac{V_{kn}}{U_{kn}} - \frac{V_{(k+1)n}}{U_{(k+1)n}} \right), \\ \frac{(-1)^{kn} U_n}{V_{(k+1)n} V_{kn}} &= \frac{1}{2} \left(\frac{U_{(k+1)n}}{V_{(k+1)n}} - \frac{U_{kn}}{V_{kn}} \right). \end{aligned} \quad (2.3)$$

Lemma 2.2. *For arbitrary integers n and t ,* □

$$V_{2n} - (-1)^{n-t} V_{2t} = \Delta U_{n-t} U_{n+t}, \quad (2.4)$$

$$V_{2n} + (-1)^{n-t} V_{2t} = V_{n-t} V_{n+t},$$

$$U_n^2 - (-1)^{n-t} U_t^2 = U_{n-t} U_{n+t}, \quad (2.5)$$

$$V_n^2 - (-1)^{n-t} V_t^2 = \Delta U_{n-t} U_{n+t}.$$

Proof. From Binet formulas of sequences $\{U_n\}$ and $\{V_n\}$, the desired results are obtained. □

Theorem 2.3. *For an odd integer t ,*

$$\begin{aligned} \sum_{n=1}^m \frac{1}{U_{(2n+1)t} + U_t} &= \frac{1}{2V_t} \left(\frac{2 - V_{2t}}{U_{2t}} - \frac{2 - V_{2(m+1)t}}{U_{2(m+1)t}} \right), \\ \sum_{n=1}^m \frac{1}{U_{(2n+1)t} - U_t} &= \frac{1}{2V_t} \left(\frac{2 + V_{2t}}{U_{2t}} - \frac{2 + V_{2(m+1)t}}{U_{2(m+1)t}} \right). \end{aligned} \quad (2.6)$$

Proof. By replacing n with $(2n+1)t$ in (2.5), we have

$$U_{(2n+1)t}^2 - U_t^2 = U_{2nt}U_{2(n+1)t}, \quad (2.7)$$

or

$$\frac{1}{U_{(2n+1)t} + U_t} = \frac{U_{(2n+1)t} - U_t}{U_{2nt}U_{2(n+1)t}}. \quad (2.8)$$

Taking $r = (2n+1)t$ and $s = t$ in the equality $V_s U_r = U_{r+s} + (-1)^s U_{r-s}$ [5], the equality (2.8) is rewritten as follows:

$$\frac{1}{U_{(2n+1)t} + U_t} = \frac{1}{V_t} \left(\frac{1}{U_{2nt}} + \frac{(-1)^t}{U_{2(n+1)t}} \right) - \frac{U_t}{U_{2nt}U_{2(n+1)t}}. \quad (2.9)$$

We have the sum

$$\sum_{n=1}^m \frac{1}{U_{(2n+1)t} + U_t} = \frac{1}{V_t} \sum_{n=1}^m \left(\frac{1}{U_{2nt}} + \frac{(-1)^t}{U_{2(n+1)t}} \right) - U_t \sum_{n=1}^m \frac{1}{U_{2nt}U_{2(n+1)t}}. \quad (2.10)$$

For an odd integer t , we have

$$\sum_{n=1}^m \left(\frac{1}{U_{2nt}} - \frac{1}{U_{2(n+1)t}} \right) = \frac{1}{U_{2t}} - \frac{1}{U_{2(m+1)t}}, \quad (2.11)$$

and taking $s = 2nt$ and $r = 2t$ in identity [5]:

$$U_{s+r}V_s - U_sV_{s+r} = 2(-1)^s U_r, \quad (2.12)$$

we get

$$\sum_{n=1}^m \frac{1}{U_{2nt}U_{2(n+1)t}} = \frac{1}{2U_{2t}} \sum_{n=1}^m \left(\frac{V_{2nt}}{U_{2nt}} - \frac{V_{2(n+1)t}}{U_{2(n+1)t}} \right) = \frac{1}{2U_{2t}} \left(\frac{V_{2t}}{U_{2t}} - \frac{V_{2(m+1)t}}{U_{2(m+1)t}} \right). \quad (2.13)$$

Substituting (2.11) and (2.13) in (2.10), we have the desired result. $\square$

For example, if we take $t = 1$ and $p = 1$ in (2.6), we have

$$\sum_{n=1}^m \frac{1}{F_{2n+1} + 1} = \frac{F_{2m+1} - 1}{F_{2(m+1)}}. \quad (2.14)$$

Note that

$$F_{2(m+1)} \sum_{n=1}^m \frac{1}{F_{2n+1} + 1} = \sum_{n=1}^m F_{2n}. \quad (2.15)$$

Corollary 2.4. For an odd integer t ,

$$\sum_{n=1}^m \frac{1}{U_{(2n+1)t} + U_t} = \begin{cases} \frac{1}{2V_t} \left(\frac{V_{(m+1)t}}{U_{(m+1)t}} - \frac{V_t}{U_t} \right), & m \text{ is even,} \\ \frac{1}{2V_t} \left(\frac{\Delta U_{(m+1)t}}{V_{(m+1)t}} - \frac{V_t}{U_t} \right), & m \text{ is odd,} \end{cases}$$

$$\sum_{n=1}^m \frac{1}{U_{(2n+1)t} - U_t} = \begin{cases} \frac{\Delta}{2V_t} \left(\frac{U_t}{V_t} - \frac{U_{(m+1)t}}{V_{(m+1)t}} \right), & m \text{ is even,} \\ \frac{1}{2V_t} \left(\frac{\Delta U_t}{V_t} - \frac{V_{(m+1)t}}{U_{(m+1)t}} \right), & m \text{ is odd.} \end{cases} \quad (2.16)$$

Proof. Using the equalities $V_{2n} = V_n^2 - 2(-1)^n = \Delta U_n^2 + 2(-1)^n$ and $U_{2n} = U_n V_n$ in Theorem 2.3, the results are obtained. $\square$

Corollary 2.5. Let t be an odd integer. For $|\beta/\alpha| < 1, t > 0$ and $|\alpha/\beta| < 1, t < 0$,

$$\sum_{n=1}^{\infty} \frac{1}{U_{(2n+1)t} + U_t} = \frac{1}{2V_t} \left(\sqrt{\Delta} - \frac{V_t}{U_t} \right),$$

$$\sum_{n=1}^{\infty} \frac{1}{U_{(2n+1)t} - U_t} = \frac{\Delta}{2V_t} \left(\frac{U_t}{V_t} - \frac{1}{\sqrt{\Delta}} \right), \quad (2.17)$$

and for $|\beta/\alpha| < 1, t < 0$ and $|\alpha/\beta| < 1, t > 0$,

$$\sum_{n=1}^{\infty} \frac{1}{U_{(2n+1)t} + U_t} = \frac{-1}{2V_t} \left(\sqrt{\Delta} + \frac{V_t}{U_t} \right),$$

$$\sum_{n=1}^{\infty} \frac{1}{U_{(2n+1)t} - U_t} = \frac{\Delta}{2V_t} \left(\frac{U_t}{V_t} + \frac{1}{\sqrt{\Delta}} \right). \quad (2.18)$$

Proof. Since

$$\lim_{n \rightarrow \infty} \left(\frac{V_{an+b}}{U_{an+b}} \right) = \begin{cases} \sqrt{\Delta} & \left| \frac{\beta}{\alpha} \right| < 1, a > 0, & \left| \frac{\alpha}{\beta} \right| < 1, a < 0, \\ -\sqrt{\Delta} & \left| \frac{\beta}{\alpha} \right| < 1, a < 0, & \left| \frac{\alpha}{\beta} \right| < 1, a > 0, \end{cases} \quad (2.19)$$

the results are easily seen by equalities (2.16). $\square$

Theorem 2.6. For an integer $m \geq 1$ and an arbitrary nonzero integer t ,

$$\sum_{n=1}^m \frac{(-1)^{nt} U_{(2n+1)t}}{(V_{(2n+1)t} - (-1)^{nt} V_t)^2} = \frac{1}{4\Delta^2 U_t} \left(\frac{V_t^2}{U_t^2} - \frac{V_{(m+1)t}^2}{U_{(m+1)t}^2} \right). \quad (2.20)$$

Proof. By replacing n with $(2n+1)t/2$ and t with $t/2$ in (2.4), we have

$$V_{(2n+1)t} - (-1)^{nt} V_t = \Delta U_{nt} U_{(n+1)t}, \quad (2.21)$$

or

$$\frac{1}{V_{(2n+1)t} - (-1)^{nt} V_t} = \frac{1}{\Delta U_{nt} U_{(n+1)t}}. \quad (2.22)$$

Multiplying equality (2.22) by $(-1)^{nt} U_{(2n+1)t} / U_{nt} U_{(n+1)t}$, we get

$$\frac{(-1)^{nt} U_{(2n+1)t}}{U_{nt} U_{(n+1)t} (V_{(2n+1)t} - (-1)^{nt} V_t)} = \frac{(-1)^{nt} U_{(2n+1)t}}{\Delta U_{nt}^2 U_{(n+1)t}^2}. \quad (2.23)$$

We have the sum:

$$\sum_{n=1}^m \frac{(-1)^{nt} U_{(2n+1)t}}{U_{nt} U_{(n+1)t} (V_{(2n+1)t} - (-1)^{nt} V_t)} = \frac{1}{\Delta} \sum_{n=1}^m \frac{(-1)^{nt} U_{(2n+1)t}}{U_{tn}^2 U_{t(n+1)}^2}. \quad (2.24)$$

Using the equalities (2.1) and (2.21), the proof is obtained. $\square$

Corollary 2.7. For an arbitrary nonzero integer t ,

$$\sum_{n=1}^{\infty} \frac{(-1)^{nt} U_{(2n+1)t}}{(V_{(2n+1)t} - (-1)^{nt} V_t)^2} = \frac{1}{4\Delta^2 U_t} \left(\frac{V_t^2}{U_t^2} - \Delta \right). \quad (2.25)$$

Proof. Taking $m \rightarrow \infty$ in Theorem 2.6 and using (2.19), the result is easily obtained. $\square$

Theorem 2.8. For an integer $m \geq 0$ and an arbitrary nonzero integer t ,

$$\sum_{n=0}^m \frac{(-1)^{nt} U_{(2n+1)t}}{(V_{(2n+1)t} + (-1)^{nt} V_t)^2} = \frac{U_{(m+1)t}^2}{4U_t V_{(m+1)t}^2}. \quad (2.26)$$

Proof. The proof of the theorem is similar to the proof of Theorem 2.6. $\square$

Corollary 2.9. For an arbitrary nonzero integer t ,

$$\sum_{n=0}^{\infty} \frac{(-1)^{nt} U_{(2n+1)t}}{(V_{(2n+1)t} + (-1)^{nt} V_t)^2} = \frac{1}{4\Delta U_t}. \quad (2.27)$$

Proof. Taking $m \rightarrow \infty$ in Theorem 2.8 and using (2.19), the result is easily obtained. $\square$

For example, if we take $t = 3$ and $p = 1$ in (2.27), we have

$$\sum_{n=0}^{\infty} \frac{(-1)^n F_{3(2n+1)}}{(L_{3(2n+1)} + (-1)^n 4)^2} = \frac{1}{40}. \quad (2.28)$$

Theorem 2.10. For an integer $m \geq 1$ and an arbitrary nonzero integer t ,

$$\begin{aligned} \sum_{n=1}^m \frac{U_{2(2n+1)t}}{(U_{(2n+1)t}^2 - U_t^2)^2} &= \frac{1}{4U_{2t}} \left(\frac{V_{2t}^2}{U_{2t}^2} - \frac{V_{2(m+1)t}^2}{U_{2(m+1)t}^2} \right), \\ \sum_{n=1}^m \frac{U_{2(2n+1)t}}{(V_{(2n+1)t}^2 - V_t^2)^2} &= \frac{1}{4\Delta^2 U_{2t}} \left(\frac{V_{2t}^2}{U_{2t}^2} - \frac{V_{2(m+1)t}^2}{U_{2(m+1)t}^2} \right). \end{aligned} \quad (2.29)$$

Proof. The proof of theorem is similar to the proof of Theorem 2.6. □

Corollary 2.11. For an arbitrary nonzero integer t ,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{U_{2(2n+1)t}}{(U_{(2n+1)t}^2 - U_t^2)^2} &= \frac{1}{4U_{2t}} \left(\frac{V_{2t}^2}{U_{2t}^2} - \Delta \right), \\ \sum_{n=1}^{\infty} \frac{U_{2(2n+1)t}}{(V_{(2n+1)t}^2 - V_t^2)^2} &= \frac{1}{4\Delta^2 U_{2t}} \left(\frac{V_{2t}^2}{U_{2t}^2} - \Delta \right). \end{aligned} \quad (2.30)$$

Proof. Taking $m \rightarrow \infty$ in Theorem 2.10 and using (2.19), the result is easily obtained. □

For example, if we take $t = 2$ in the equality (2.30), we have

$$\sum_{n=1}^{\infty} \frac{U_{4(2n+1)}}{(V_{2(2n+1)}^2 - V_2^2)^2} = \frac{1}{\Delta^2 U_4^3}. \quad (2.31)$$

References

- [1] R. P. Backstrom, "On reciprocal series related to Fibonacci numbers with subscripts in arithmetic progression," *The Fibonacci Quarterly*, vol. 19, pp. 14–21, 1981.
- [2] B. S. Popov, "On certain series of reciprocals of Fibonacci numbers," *The Fibonacci Quarterly*, vol. 22, no. 3, pp. 261–265, 1984.
- [3] B. S. Popov, "Summation of reciprocal series of numerical functions of second order," *The Fibonacci Quarterly*, vol. 24, no. 1, pp. 17–21, 1986.
- [4] N. Gauthier, "Solution to problem H-680," *The Fibonacci Quarterly*, vol. 49, no. 1, pp. 90–92, 2011.
- [5] S. Vajda, *Fibonacci & Lucas Numbers, and the Golden Section. Theory and Applications*, John Wiley & Sons, New York, NY, USA, 1989.