Research Article

Bifurcation Analysis in a Delayed Diffusive Leslie-Gower Model

Shuling Yan, Xinze Lian, Weiming Wang, and Youbin Wang

College of Mathematics and Information Science, Wenzhou University, Wenzhou 325035, China

Correspondence should be addressed to Weiming Wang; weimingwang2003@163.com

Received 29 September 2012; Accepted 18 November 2012

Academic Editor: Yanbin Sang

Copyright © 2013 Shuling Yan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate a modified delayed Leslie-Gower model under homogeneous Neumann boundary conditions. We give the stability analysis of the equilibria of the model and show the existence of Hopf bifurcation at the positive equilibrium under some conditions. Furthermore, we investigate the stability and direction of bifurcating periodic orbits by using normal form theorem and the center manifold theorem.

1. Introduction

The dynamic relationship between predators and their preys has long been and will continue to be one of dominant themes in both ecology and mathematical ecology due to its universal existence and importance. A major trend in theoretical work on prey-predator dynamics has been to derive more realistic models, trying to keep to maximum the unavoidable increase in complexity of their mathematics [1]. In this optic, recent years, the important Leslie-Gower predator-prey model [2, 3] has been extensively studied in [4–7]. A modified version of Leslie-Gower predator-prey model with Holling-type II functional response takes the form

$$\begin{aligned}\frac{dH}{dt} &= H(a_1 - bH) - \frac{c_1 HP}{k_1 + H}, \\ \frac{dP}{dt} &= P\left(a_2 - \frac{c_2 P}{k_2 + H}\right),\end{aligned}\quad (1)$$

where H and P represent prey and predator population densities at time t , respectively. a_1 , a_2 , b , c_1 , c_2 , k_1 , and k_2 are positive constants. a_1 is the growth rate of prey H . a_2 describes the growth rate of predator P . b measures the strength of competition among individuals of species H . c_1 is the maximum value of the per capita reduction of H due to P , and c_2 is the maximum value of the per capita reduction of P due to H , which is not available in abundance. k_1 measures the extent to which environment provides protection to prey

H . k_2 measures the extent to which environment provides protection to the predator P .

On the other hand, time delay plays an important role in many biological dynamical systems, being particularly relevant in ecology [1]. For some predator-prey systems, the rate of the prey population depends on the predation of predator in the earlier times [8–14]. The results indicated that delay differential equations exhibit much more complicated dynamics than ordinary differential equations since a time delay could cause a stable equilibrium to become unstable and induce bifurcations.

In this paper, we will focus on the complex dynamics of the delay effect in the extended reaction-diffusion model. The reproduction of the individuals is modeled by diffusion with diffusion coefficients $D_1 > 0$ and $D_2 > 0$ for the prey and predator, respectively. This basic model is described by a system of two partial differential equations:

$$\begin{aligned}\frac{\partial H}{\partial t} &= H(a_1 - bH) - \frac{c_1 HP(t - \tau)}{k_1 + H} \\ &\quad + D_1 \Delta H, \quad x \in \Omega, \quad t > 0, \\ \frac{\partial P}{\partial t} &= P\left(a_2 - \frac{c_2 P}{k_2 + H}\right) + D_2 \Delta P, \quad x \in \Omega, \quad t > 0, \\ \frac{\partial H}{\partial \mathbf{n}} &= \frac{\partial P}{\partial \mathbf{n}} = 0, \quad x \in \partial\Omega, \quad t > 0,\end{aligned}\quad (2)$$

where $H = H(t, x)$, $P = P(t, x)$. $\Delta = \partial^2 / \partial x^2$, Ω is a bounded open domain in $\mathbb{R}$ with boundary $\partial\Omega$, $\mathbf{n}$ is the outward unit

FIGURE 1: The bifurcation diagram displays the distribute of the positive roots; the number indicates the number of positive equilibria.

normal vector on $\partial\Omega$, and homogeneous Neumann boundary conditions reflect the situation where the population cannot move across the boundary of the domain. And we incorporate a single discrete delay $\tau > 0$ in the negative feedback of the predator's density.

The rest of the paper is organized as follows. In Section 2, we give the stability property of the equilibria of model (1). In Section 3, we mainly analyze the distribution of the roots of the characteristic equation and show the occurrence of Hopf bifurcation at the positive equilibrium of model (2) under some conditions. In Section 4, we investigate the stability and direction of bifurcating periodic orbits by using normal form of theorem and the center manifold theorem, corresponding to theorems we also give some numerical simulations.

2. Equilibria Stability

In this section, we consider the existence and stability of the equilibria of model (1).

It is easy to verify that model (1) always has three boundary equilibria:

- (i) $E_1 = (0, 0)$ (extinction of prey and predator), which is a nodal source point;
- (ii) $E_2 = (a_1/b, 0)$ (extinction of the predator), which is a saddle point;
- (iii) $E_3 = (0, a_2k_2/c_2)$ (extinction of the prey), which is a stable node when $a_2k_2/c_2 > a_1k_1/c_1$.

For the positive equilibria, we have

$$a_1 - bH - \frac{c_1P}{k_1 + H} = 0, \quad a_2 - \frac{c_2P}{k_2 + H} = 0, \quad (3)$$

which yields

$$bc_2H^2 - (a_1c_2 - a_2c_1 - bc_2k_1)H + a_2c_1k_2 - a_1c_2k_1 = 0. \quad (4)$$

For simplicity, we define

$$\begin{aligned} A &= bc_2, & B &= a_1c_2 - a_2c_1 - bc_2k_1, \\ C &= a_2c_1k_2 - a_1c_2k_1, \end{aligned} \quad (5)$$

then (4) can be written as

$$AH^2 - BH + C = 0, \quad (6)$$

which has two roots given by

$$h_+ = \frac{B + \sqrt{B^2 - 4AC}}{2A}, \quad h_- = \frac{B - \sqrt{B^2 - 4AC}}{2A}. \quad (7)$$

Case 1 ($C < 0$, i.e., $k_2 < a_1c_2k_1/a_2c_1$). Model (1) has a unique positive equilibrium $E^* = (h^*, p^*) = (h_+, (a_2/c_2)(h_+ + k_2))$.

Case 2 ($C = 0$, i.e., $k_2 = a_1c_2k_1/a_2c_1$).

- (i) If $B > 0$, that is, $k_1 < (a_1c_2 - a_2c_1)/bc_2$, model (1) has a unique positive equilibrium $E = (h, p) = (h_+, (a_2/c_2)(h_+ + k_2))$;
- (ii) If $B \leq 0$, that is, $k_1 \geq (a_1c_2 - a_2c_1)/bc_2$, (4) has no positive root; hence model (1) has no positive equilibrium.

Case 3 ($C > 0$, i.e., $k_2 > a_1c_2k_1/a_2c_1$).

- (i) Suppose that $B > 0$, that is, $k_1 < (a_1c_2 - a_2c_1)/bc_2$, then
 - (i1) if $B^2 - 4AC > 0$, that is, $k_2 < (a_2c_1 - a_1c_2 - bc_2k_1)^2 / (4a_2bc_1c_2 + k_1)$, model (1) has two positive equilibria $E_+ = (h_+, p_+) = (h_+, (a_2/c_2)(h_+ + k_2))$ and $E_- = (h_-, p_-) = (h_-, (a_2/c_2)(h_- + k_2))$;
 - (i2) if $B^2 - 4AC = 0$, that is, $k_2 = (a_2c_1 - a_1c_2 - bc_2k_1)^2 / (4a_2bc_1c_2 + k_1)$, (4) has a unique positive root of multiplicity 2 given by $h_e = B/2A = h_+ = h_-$, then model (1) has a unique positive equilibrium $E_e = (h_e, p_e) = (h_e, (a_2/c_2)(h_e + k_2))$;
 - (i3) if $B^2 - 4AC < 0$, that is, $k_2 > (a_2c_1 - a_1c_2 - bc_2k_1)^2 / (4a_2bc_1c_2 + k_1)$, (4) has no positive root; hence model (1) has no positive equilibrium;
- (ii) if $B < 0$, that is, $k_1 > (a_1c_2 - a_2c_1)/bc_2$, (4) has no positive root; hence model (1) has no positive equilibrium.

We show the bifurcation diagram to display the distribute of the positive roots; in Figure 1, the whole region has been divided into six parts; the number indicates the number of positive equilibria.

Next, we analyze the stability of these positive equilibria. Let $E = (h, p)$ be arbitrary positive equilibrium, and the Jacobian matrix for $E = (h, p)$ is given by

$$J(E) = \begin{pmatrix} -bh + \frac{c_1hp}{(h+k_1)^2} & \frac{-c_1h}{h+k_1} \\ \frac{a_2^2}{c_2} & -a_2 \end{pmatrix}. \quad (8)$$

FIGURE 2: Phase portraits of model (1) with the parameters $a_1 = 2$, $a_2 = 0.5$, $b = 1$, $c_1 = c_2 = 2$, $k_1 = 0.4$, and $k_2 = 2.1$. In this case, $E_1 = (0, 0)$ is a nodal source point; $E_2 = (2, 0)$ is a saddle point; $E_3 = (0, 0.525)$ is a nodal sink point, which is locally asymptotically stable; $E_+ = (0.7791, 0.7197)$ is a spiral sink, which is locally asymptotically stable; and $E_- = (0.3208, 0.6052)$ is a saddle point. There exists a separatrix curve determined by the stable manifold of E_- . The dashed curve is the H -nullcline $H(a_1 - bH) - c_1HP/(k_1 + H)$, and the dotted curve is the P -nullcline $P(a_2 - c_2P/(k_2 + H))$.

The corresponding characteristic equation is

$$\lambda^2 - \text{tr}(J(E))\lambda + \det(J(E)) = 0, \quad (9)$$

where

$$\begin{aligned} \text{tr}(J(E)) &= \frac{c_1 h p}{(h + k_1)^2} - a_2 - b h \\ &= \frac{h(a_1 - a_2 - 2bh) - (a_2 + bh)k_1}{h + k_1}, \end{aligned} \quad (10)$$

$$\det(J(E)) = \frac{a_2 h(a_2 c_1 - a_1 c_2 + b c_2 k_1 + 2b c_2 h)}{c_2(h + k_1)}.$$

And the sign of $\det(J(E))$ is determined by

$$F_1(h) = a_2 c_1 - a_1 c_2 + b c_2 k_1 + 2b c_2 h = 2Ah - B. \quad (11)$$

Thus,

$$\begin{aligned} F_1(h^*) &= 2Ah^* - B = \sqrt{B^2 - 4AC} > 0, \\ F_1(\bar{h}) &= 2A\bar{h} - B = B > 0, \\ F_1(h_+) &= 2Ah_+ - B = \sqrt{B^2 - 4AC} > 0; \\ F_1(h_-) &= 2Ah_- - B = -\sqrt{B^2 - 4AC} < 0, \\ F_1(h_e) &= 2Ah_e - B = 0. \end{aligned} \quad (12)$$

Hence, $\det(J(E^*)) > 0$, $\det(J(\bar{E})) > 0$, $\det(J(E_+)) > 0$, $\det(J(E_-)) < 0$, and $\det(J(E_e)) = 0$. Obviously, the positive equilibrium E_- is a saddle point.

In the following, we study the stability of other positive equilibria. The sign of $\text{tr}(J(E))$ is determined by

$$G_1(h) = h(a_1 - a_2 - 2bh) - (a_2 + bh)k_1. \quad (13)$$

Then we can get

$$\begin{aligned} G_1(h^*) &= \frac{h^* a_2 (c_1 - c_2)}{c_2} - \frac{h^* \sqrt{B^2 - 4AC}}{c_2} - a_2 k_1, \\ G_1(\bar{h}) &= \frac{\bar{h} a_2 (c_1 - c_2)}{c_2} - \frac{\bar{h} B}{c_2} - a_2 k_1, \\ G_1(h_+) &= \frac{h_+ a_2 (c_1 - c_2)}{c_2} - \frac{h_+ \sqrt{B^2 - 4AC}}{c_2} - a_2 k_1, \end{aligned} \quad (14)$$

$$G_1(h_e) = \frac{h_e a_2 (c_1 - c_2)}{c_2} - a_2 k_1.$$

Hence, if $c_1 \leq c_2$, then $\text{tr}(J(E^*)) < 0$, $\text{tr}(J(\bar{E})) < 0$, and $\text{tr}(J(E_+)) < 0$ are true. Summarizing the above, we can obtain the following theorem.

Theorem 1. For model (1),

- (i) if $k_2 < a_1 c_2 k_1 / a_2 c_1$ holds, the unique positive equilibrium E^* is locally asymptotically stable for $c_1 \leq c_2$;
- (ii) if $k_2 = a_1 c_2 k_1 / a_2 c_1$ and $k_1 < (a_1 c_2 - a_2 c_1) / b c_2$ hold, the unique positive equilibrium $\bar{E}$ is locally asymptotically stable for $c_1 \leq c_2$;
- (iii) if $a_1 c_2 k_1 / a_2 c_1 < k_2 < (a_2 c_1 - a_1 c_2 - b c_2 k_1)^2 / 4 a_2 b c_1 c_2 + k_1$ and $k_1 < (a_1 c_2 - a_2 c_1) / b c_2$ hold, model (1) has two positive equilibria, the positive equilibrium E_+ is locally asymptotically stable for $c_1 \leq c_2$, and E_- is a saddle point.

Figure 2 shows the dynamics of model (1). In this case, E_1 is a nodal source point; E_2 is a saddle point; E_3 is a nodal sink point, which is locally asymptotically stable; E_+ is locally asymptotically stable; E_- is a saddle point. There exists a separatrix curve determined by the stable manifold of E_- , which divides the behavior of trajectories; that is, the stable manifold of saddle E_- splits the feasible region into two parts such that orbits initiating inside tend to the positive equilibrium E_+ , while orbits initiating outside tend to E_3 except for the stable manifolds of E_- . This means that, in this situation, the trajectories of the model can have different behavior strongly depending on the initial conditions.

Theorem 2. For model (1), if the unique positive equilibrium $E_e = (h_e, (a_2/c_2)(h_e + k_2))$ exists, and $k_1 = (a_1 c_2 - a_2 c_1)(c_1 - c_2) / b c_2 (c_1 + c_2)$, then it is a cusp of codimension 2.

Proof. The Jacobian matrix at $E_e = (h_e, p_e)$ is

$$J_{E_e} = \begin{pmatrix} -\frac{a_2 c_1 (a_2 c_1 - a_1 c_2 + b c_2 k_1)}{c_2 (b c_2 k_1 - a_2 c_1 + a_1 c_2)} & \frac{c_1 (a_2 c_1 - a_1 c_2 + b c_2 k_1)}{b c_2 k_1 - a_2 c_1 + a_1 c_2} \\ \frac{a_2^2}{c_2} & -a_2 \end{pmatrix}, \quad (15)$$

we have know that $\det(J(E_e)) = 0$. Moreover, $\text{tr}(J(E_e)) = 0$, if and only if

$$k_1 = \frac{(a_1 c_2 - a_2 c_1)(c_1 - c_2)}{b c_2 (c_1 + c_2)}. \quad (16)$$

Then

$$J_{E_e} = \begin{pmatrix} a_2 & -c_2 \\ \frac{a_2^2}{c_2} & -a_2 \end{pmatrix} = a_2 \begin{pmatrix} 1 & -\frac{c_2}{a_2} \\ \frac{a_2}{c_2} & -1 \end{pmatrix}, \quad (17)$$

and the associate Jordan matrix is

$$\hat{J} = \begin{pmatrix} 0 & \frac{c_1 (a_2 c_1 - a_1 c_2 + b c_2 k_1)}{b c_2 k_1 - a_2 c_1 + a_1 c_2} \\ 0 & 0 \end{pmatrix}. \quad (18)$$

Hence, following [15, 16], we know that the unique positive equilibrium E_e is a cusp of codimension 2. $\square$

3. Stability and Hopf Bifurcation Analysis in Delayed Reaction-Diffusion Model (2)

According to the previous section, for model (1), we know that E_1 , E_2 , and E_3 are unstable and E_- is a saddle point, and note that a solution of the model (1) is also a solution of the model (2), so they are also unstable for model (2). In the following, we will focus on the dynamics of the positive equilibria of model (2). As an example, we only give the proof of the unique positive equilibrium E^* of model (2).

Introducing small perturbations $\tilde{H} = H - h^*$, and $\tilde{P} = P - p^*$ and dropping the hats for simplicity of notation, then we have

$$\begin{aligned} \frac{\partial H}{\partial t} &= (H + h^*) (a_1 - b(H + h^*)) \\ &\quad - \frac{c_1 (H + h^*) (P(t - \tau) + p^*)}{k_1 + H + h^*} \\ &\quad + D_1 \Delta H, \quad x \in \Omega, \quad t > 0, \\ \frac{\partial P}{\partial t} &= (P + p^*) \left(a_2 - \frac{c_2 (P + p^*)}{k_2 + H + h^*} \right) \\ &\quad + D_2 \Delta P, \quad x \in \Omega, \quad t > 0, \\ \frac{\partial H}{\partial \mathbf{n}} &= \frac{\partial P}{\partial \mathbf{n}} = 0, \quad x \in \partial\Omega, \quad t > 0. \end{aligned} \quad (19)$$

Denote

$$X = \left\{ H, P \in W^{2,2}(\Omega) : \frac{\partial H}{\partial \mathbf{n}} = \frac{\partial P}{\partial \mathbf{n}} = 0, \quad x \in \partial\Omega \right\}. \quad (20)$$

In the abstract space $C([- \tau, 0], X)$, model (19) can be regarded as the following abstract functional differential equation.

$$\frac{\partial U}{\partial t} = D \Delta U + L(U_t) + F(U_t), \quad (21)$$

where $U = (H, P)^T$, $U_t = U(t + \theta)$, $\theta \in [- \tau, 0]$, $D = \begin{pmatrix} D_1 & 0 \\ 0 & D_2 \end{pmatrix}$, $\text{dom}(\Delta) \subset X$, and $L : C([- \tau, 0], X) \mapsto X$, $F : C([- \tau, 0], X) \mapsto X$ are given by

$$\begin{aligned} L(\phi) &= \begin{pmatrix} \left(-bh^* + \frac{c_1 h^* p^*}{(h^* + k_1)^2} \right) \phi_1(0) + \frac{-c_1 h^*}{h^* + k_1} \phi_2(-\tau) \\ \frac{a_2^2}{c_2} \phi_1(0) - a_2 \phi_2(0) \end{pmatrix}, \\ F(\phi) &= \begin{pmatrix} \left(\frac{c_1 k_1 p^*}{(k_1 + h^*)^3} - b \right) \phi_1^2(0) - \frac{c_1 k_1}{(k_1 + h^*)^2} \phi_1(0) \phi_2(-\tau) \\ -\frac{c_2 p^{*2}}{(k_2 + h^*)^3} \phi_1^2(0) + \frac{2c_2 p^*}{(k_2 + h^*)^2} \phi_1(0) \phi_2(0) - \frac{c_2}{k_2 + h^*} \phi_2^2(0) \end{pmatrix} \end{aligned} \quad (22)$$

here $\phi = (\phi_1, \phi_2)^T = U_t \in C([- \tau, 0], X)$. Then the linearization of model (19) near (h^*, p^*) is

$$\frac{\partial U}{\partial t} = D \Delta U + L(U_t). \quad (23)$$

Following [17], we obtain that the characteristic equation for liner model (23) is

$$\lambda y - D \Delta y - L(e^{\lambda \cdot} y) = 0, \quad y \in \text{dom}(\Delta) \subset X, \quad y \neq 0. \quad (24)$$

It is well known that the eigenvalue problem

$$\begin{aligned} -\Delta \psi &= \mu \psi, \quad x \in \Omega, \\ \frac{\partial \psi}{\partial \mathbf{n}} &= 0, \quad x \in \partial\Omega, \end{aligned} \quad (25)$$

has eigenvalues $\mu_n = \{\mu_n^i = -D_i n^2, i = 1, 2, \text{ and } n = 0, 1, 2, \dots\}$, with the corresponding eigenfunctions $\psi_n(x) = \cos nx$ ($n \in \mathbb{N} = \{0, 1, 2, \dots\}$).

Substituting $y = \sum_{n=0}^{\infty} \cos nx (y_{1n}, y_{2n})^T$ into characteristic equation (24), we obtain

$$\begin{aligned} &\begin{pmatrix} -bh^* + \frac{c_1 h^* p^*}{(h^* + k_1)^2} - D_1 n^2 & \frac{-c_1 h^*}{h^* + k_1} e^{-\lambda \tau} \\ \frac{a_2^2}{c_2} & -a_2 - D_2 n^2 \end{pmatrix} \begin{pmatrix} y_{1n} \\ y_{2n} \end{pmatrix} \\ &= \lambda \begin{pmatrix} y_{1n} \\ y_{2n} \end{pmatrix}. \end{aligned} \quad (26)$$

Therefore the characteristic equation (24) is equivalent to

$$\lambda^2 + A_n \lambda + B_n e^{-\lambda \tau} + C_n = 0, \quad n \in \mathbb{N}, \quad (27)$$

where

$$\begin{aligned} A_n &= (D_1 + D_2)n^2 + a_2 + bh^* - \frac{c_1 h^* p^*}{(h^* + k_1)^2}, \\ B_n &= \frac{a_2^2 c_1 h^*}{c_2 (h^* + k_1)^2}, \\ C_n &= D_1 D_2 n^4 + \left(D_1 + D_2 + a_2 + bh^* - \frac{c_1 h^* p^*}{(h^* + k_1)^2} \right) n^2 \\ &\quad + a_2 bh^* - \frac{a_2 c_1 h^* p^*}{(h^* + k_1)^2}. \end{aligned} \quad (28)$$

The stability of the positive equilibrium E^* can be determined by the distribution of the roots of (27); that is, the equilibrium E^* is locally asymptotically stable if all the roots of (27) have negative real parts. From the result of [18], the sum of the multiplicities of the roots of (27) in the open right half plane changes only if a root appears on or crosses the imaginary axis. It can be verified that $\lambda = 0$ is not a root of (27) for $n \in \mathbb{N}$.

Theorem 3. *If $k_2 < \min\{a_1 c_2 k_1 / a_2 c_1, bc_2(h^* + k_1)^2 / a_2 c_1 - 2h^* - k_1\}$ holds, then the unique positive equilibrium E^* of model (2) is locally asymptotically stable.*

Proof. Let $\pm i\omega$ ($\omega > 0$) be a pair of roots of (27); substituting $i\omega$ into (27), then we have

$$-\omega^2 + iA_n \omega + B_n e^{-i\omega \tau} + C_n = 0. \quad (29)$$

Separating the real part from image part, we have

$$\begin{aligned} -\omega^2 + B_n \cos \omega \tau + C_n &= 0, \\ A_n \omega - B_n \sin \omega \tau &= 0, \end{aligned} \quad (30)$$

then

$$\omega^4 + (A_n^2 - 2C_n)\omega^2 + C_n^2 - B_n^2 = 0, \quad (31)$$

where

$$\begin{aligned} A_n^2 - 2C_n &= \left(bh^* - \frac{c_1 h^* p^*}{(h^* + k_1)^2} + D_1 n^2 \right)^2 \\ &\quad + (a_2 + D_2 n^2)^2 > 0, \\ C_n^2 - B_n^2 &= D_1 D_2 n^4 \\ &\quad + \left(D_1 + D_2 + a_2 + bh^* - \frac{c_1 h^* p^*}{(h^* + k_1)^2} \right) n^2 \\ &\quad - \frac{a_2 c_1 h^* p^*}{(h^* + k_1)^2} - \frac{a_2^4 c_1^2 (h^*)^2}{c_2^2 (h^* + k_1)^2} + a_2 bh^*. \end{aligned} \quad (32)$$

Obviously, if $k_2 < bc_2(h^* + k_1)^2 / a_2 c_1 - 2h^* - k_1$, $C_n^2 - B_n^2 > 0$ is true. Thus (31) has no positive roots for all $n \in \mathbb{N}$. Hence, all the roots of (27) have negative real part. This completes the proof. $\square$

If there exists an integer $n_0 \in \mathbb{N}$ such that for $0 \leq n \leq n_0$, $C_n^2 - B_n^2 < 0$, then (31) has a unique positive real root

$$\omega_0^n = \sqrt{\frac{-(A_n^2 - 2C_n) + \sqrt{(A_n^2 - 2C_n)^2 - 4(C_n^2 - B_n^2)}}{2}}, \quad (33)$$

and (27) has a pair of pure imaginary roots $\pm i\omega_0^n$, and

$$\tau_n^j = \tau_n^0 + \frac{2\pi j}{\omega_n}, \quad j = 0, 1, 2, \dots, \quad 0 \leq n \leq n_0, \quad n_0 \in \mathbb{N}, \quad (34)$$

where $\tau_n^0 = \arccos((\omega_0^{n^2} - C_n)/B_n)/\omega_0^n$.

Let $\lambda(\tau) = \gamma(\tau) + i\omega(\tau)$ be the root of (24), where $\gamma(\tau_n^j) = 0$ and $\omega(\tau_n^j) = \omega_0^n$ when τ is close to τ_n^j . Then we have the following transversality condition.

Lemma 4. *For $0 \leq n \leq n_0$ ($n_0 \in \mathbb{N}$), if*

$$(H1) \quad D_1 D_2 n^4 + (D_1 + D_2 + a_2 + bh^* - c_1 h^* p^* / (h^* + k_1)^2) n^2 < a_2 c_1 h^* p^* / (h^* + k_1)^2 + a_2^4 c_1^2 (h^*)^2 / c_2^2 (h^* + k_1)^2 - a_2 bh^* \text{ holds, then } (d\gamma/d\tau)|_{\tau=\tau_n^j} > 0 \text{ for } j = 0, 1, 2, \dots$$

From this transversality condition, we know that when τ passes through these critical values τ_n^j , the sum of the multiplicities of the roots of (27) in the open right half plane will increase at least two.

Summarizing the above results, we can obtain the following theorem.

Theorem 5. *For $0 \leq n \leq n_0$ ($n_0 \in \mathbb{N}$) if (H1) holds, the following statements are true:*

- (i) *if $\tau \in [0, \tau_n^0)$, then the equilibrium point E^* is locally asymptotically stable;*
- (ii) *if $\tau > \tau_n^0$, then the equilibrium E^* is unstable;*
- (iii) *$\tau = \tau_n^j$ ($j = 0, 1, 2, \dots$) are Hopf bifurcation values of model (2).*

4. Direction and Stability of Spatial Hopf Bifurcation

In the previous section, we have obtained the conditions under which model (2) undergoes a Hopf bifurcation at the equilibrium point E^* when τ crosses through the critical value τ_n^j ($0 \leq n \leq n_0$, $n_0 \in \mathbb{N}$, $j = 0, 1, 2, \dots$). In this section, we will study the direction of the Hopf bifurcation and the stability of the bifurcating periodic solutions by employing the center manifold theorem and normal form method [17, 19] for partial differential equations with delay.

Then we compute the direction and stability of the Hopf bifurcation when $\tilde{\tau} = \tau_n^j$ for fixed $j \in \{0, 1, 2, \dots\}$.

Define

$$\begin{aligned}\mu &= -\frac{\operatorname{Re}(c_1(0))}{\operatorname{Re}(\lambda'(\tilde{\tau}))}, \\ \beta_2 &= 2 \operatorname{Re}(c_1(0)), \\ T_2 &= -\frac{\operatorname{Im}(c_1(0)) + \mu_2 \operatorname{Im}(\lambda'(\tilde{\tau}))}{\omega_0^n \tilde{\tau}},\end{aligned}\quad (35)$$

where $c_1(0)$ is defined in the appendix. Then we can get the following theorem.

Theorem 6. For model (19), if (H1) holds, we have the following:

- (i) μ determines the direction of the Hopf bifurcation: if $\mu > 0$ ($\mu < 0$), then the bifurcating periodic solutions exist for $\tau > \tilde{\tau}$ ($\tau < \tilde{\tau}$);
- (ii) β_2 determines the stability of bifurcating periodic solutions: the bifurcating periodic solutions are orbitally asymptotically stable (unstable) if $\beta_2 < 0$ ($\beta_2 > 0$);
- (iii) T_2 determines the period of the bifurcating periodic solutions: the period increases (decreases) if $T_2 > 0$ ($T_2 < 0$).

The proof is deferred to the appendix.

5. Conclusions and Remarks

In this paper, we have considered a modified version of Leslie-Gower model with Holling-type II functional and delayed diffusive predator-prey model under homogeneous Neumann boundary conditions. The value of this study lies in two folds. First, it presents local asymptotic stability of the equilibria of model with and without delay and the existence of Hopf bifurcation, which indicates that the dynamics induced by time delay are rich and complex. Second, it give the analysis of direction and stability of spatial Hopf bifurcation, from which one can find that small sufficiently delays cannot change the stability of the positive equilibrium and large delays cannot only destabilize the positive equilibrium but also induce oscillatory behaviors near the positive equilibrium.

In the following, we give some numerical examples to illustrate the dynamical behaviors of model (2). In Figure 3, $\tau = 2 < \tau_0^0 = 3.435144529$, the unique positive equilibrium $E^* = (2, 1.5)$ remains the stability; the population of the predator and the prey will tend to a steady state. However, in Figure 4, $\tau = 4 > \tau_0^0$, the positive equilibrium E^* losses its stability and Hopf bifurcation occurs, which means that a family of stable periodic solutions bifurcate from E^* and the system goes into oscillations; it means that the predator coexists with the prey with oscillatory behaviors.

Our results show that time-delay can make a stable equilibrium to become unstable and induce Hopf bifurcation and the system goes into oscillations; that's to say, the dynamical

behaviors of the delay reaction-diffusion equations are much more complex and rich than reaction-diffusion equations.

Appendix

A. The Proof of Theorem 6

Setting $\tau = \tilde{\tau} + \alpha$, then $\alpha = 0$ is the Hopf bifurcation of model (19). Let $\tilde{H}(t, x) = H(\tau t, x)$, $\tilde{P}(t, x) = P(\tau t, x)$, and drop the tilde for the sake of simplicity, then (19) can be transformed into

$$\begin{aligned}\frac{\partial H}{\partial t} &= \tau(H + h^*)(a_1 - b(H + h^*)) \\ &\quad - \frac{c_1(H + h^*)(P(t-1) + p^*)}{k_1 + H + h^*} \\ &\quad + D_1 \Delta H, \quad x \in \Omega, \quad t > 0, \\ \frac{\partial P}{\partial t} &= \tau(P + p^*) \left(a_2 - \frac{c_2(P + p^*)}{k_2 + H + h^*} \right) \\ &\quad + D_2 \Delta P, \quad x \in \Omega, \quad t > 0, \\ \frac{\partial H}{\partial \mathbf{n}} &= \frac{\partial P}{\partial \mathbf{n}} = 0, \quad x \in \partial\Omega, \quad t > 0.\end{aligned}\quad (A.1)$$

And the abstract functional differential equation can also be written in the form

$$\frac{dU}{dt} = \tilde{\tau} D \Delta U + \tilde{\tau} L(U_t) + G(U_t, \alpha), \quad (A.2)$$

where

$$\begin{aligned}L(\phi) &= \begin{pmatrix} \left(-bh^* + \frac{c_1 h^* p^*}{(h^* + k_1)^2} \right) \phi_1(0) - \frac{c_1 h^*}{h^* + k_1} \phi_2(-1) \\ \frac{a_2^2}{c_2} \phi_1(0) - a_2 \phi_2(0) \end{pmatrix}, \\ G(\phi, \alpha) &= \alpha D \Delta \phi + \alpha L(\phi) + (\tilde{\tau} + \alpha) F(\phi), \\ F(\phi) &= \begin{pmatrix} \left(\frac{c_1 k_1 p^*}{(k_1 + h^*)^3} - b \right) \phi_1^2(0) - \frac{c_1 k_1}{(k_1 + h^*)^2} \phi_1(0) \phi_2(-1) \\ \frac{c_2 p^{*2}}{(k_2 + h^*)^3} \phi_1^2(0) + \frac{2c_2 p^*}{(k_2 + h^*)^2} \phi_1(0) \phi_2(0) - \frac{c_2}{k_2 + h^*} \phi_2^2(0) \end{pmatrix},\end{aligned}\quad (A.3)$$

for $\phi = (\phi_1, \phi_2) = U_t \in C([-1, 0], X)$.

From Section 2, we know that $\pm i\omega_0^n \tilde{\tau}$ are a pair of simple purely imaginary eigenvalues of the liner system

$$\frac{dU}{dt} = \tilde{\tau} L(U_t) + \tilde{\tau} D \Delta U, \quad (A.4)$$

where $U(t) = (H(t), P(t))^T \in \mathbb{R}^2$, and $U_t(\theta)$ is defined by $U_t(\theta) = U(t + \theta)$.

FIGURE 3: The solutions of model (2) tends to aperiodically oscillatory orbit. The parameters are taken as $D_1 = 0.01$, $D_2 = 0.1$, $a_1 = 2$, $a_2 = 1$, $b = 0.5$, $c_1 = c_2 = 2$, $k_1 = k_2 = 1$, and the initial values $H(x, t) = 2 + 0.1t \sin x$, $P(x, t) = 1.5 + 0.1t \cos x$, $t \in [-2, 0]$, $x \in [0, \pi]$. In this case, $\tau = 4 > \tau_0^0$.

FIGURE 4: The solutions of model (2) tends to aperiodically oscillatory orbit. The parameters are taken as $D_1 = 0.01$, $D_2 = 0.1$, $a_1 = 2$, $a_2 = 1$, $b = 0.5$, $c_1 = c_2 = 2$, $k_1 = k_2 = 1$, and the initial values $H(x, t) = 2 + 0.1t \sin x$, $P(x, t) = 1.5 + 0.1t \cos x$, $t \in [-4, 0]$, $x \in [0, \pi]$. In this case, $\tau = 4 > \tau_0^0$.

By using the Riesz representation theorem [20], we have a function $\eta(\theta, \alpha)$ of bounded variation for $\theta \in [-1, 0]$ such that

$$L(\phi) + D\Delta\phi(0) = \int_{-1}^0 d\eta(\theta, \alpha) \phi(\theta), \quad (\text{A.5})$$

for $\phi(\theta) \in C([-1, 0], R^2)$.

Because in this paper, we discuss the existence of the Hopf bifurcation when $\tau = \tau_0^0$, that is $n = 0$; here we choose

$$\eta(\theta, \alpha) = \begin{pmatrix} -bh^* + \frac{c_1 h^* p^*}{(h^* + k_1)^2} & 0 \\ \frac{a_2^2}{c_2} & -a_2 \end{pmatrix} \delta(\theta) + \begin{pmatrix} 0 & -c_1 h^* \\ 0 & 0 \end{pmatrix} \delta(\theta + 1), \quad (\text{A.6})$$

where δ is the Dirac delta function. For $\phi(\theta) \in C^1([-1, 0], R^2)$, define $A(\alpha)$ as

$$A(\alpha)\phi = \begin{cases} \frac{d\phi(\theta)}{d\theta}, & \theta \in [-1, 0), \\ \int_{-1}^0 d\eta(\theta, \alpha) \phi(\theta), & \theta = 0, \end{cases} \quad (\text{A.7})$$

and for $\psi = (\psi_1, \psi_2) \in C^1([0, 1], (R^2)^*)$, define

$$A^*(\psi(s)) = \begin{cases} -\frac{d\psi(s)}{ds}, & s \in [-1, 0), \\ \int_{-1}^0 \psi(-\xi) d\eta(\theta, 0), & s = 0. \end{cases} \quad (\text{A.8})$$

Then $A(0)$ and A^* are adjoint operators under the bilinear form

$$\begin{aligned} & (\psi(s), \phi(\theta)) \\ &= \bar{\psi}(0) \phi(0) - \int_{-1}^0 \int_0^\theta \bar{\psi}(\xi - \theta) d\eta(\theta, \alpha) \phi(\xi) d\xi. \end{aligned} \quad (\text{A.9})$$

We know that $\pm i\omega_0^n$ are eigenvalues of $A(0)$. Since $A(0)$ and $A^*(0)$ are two adjoint operators, then $\pm i\omega_0^n$ are also eigenvalue of A^* ; we shall first try to obtain eigenvector of $A(0)$ and A^* corresponding to the eigenvalues $i\omega_0^n$ and $-i\omega_0^n$, respectively. Let $q(\theta) = (1, \rho)^T e^{i\omega_0^n \theta}$ be the eigenvector of $A(0)$ corresponding to the eigenvalue $i\omega_0^n$. Then by definition of eigenvector we have $A(0)q(\theta) = q(\theta)i\omega_0^n$. Therefore, from (A.5), (A.6), and definition of $A(0)$ we get

$$\begin{pmatrix} -bh^* + \frac{c_1 h^* p^*}{(h^* + k_1)^2} - i\omega_0^n & \frac{-c_1 h^*}{h^* + k_1} e^{-i\omega_0^n \bar{\tau}} \\ \frac{a_2^2}{c_2} & -a_2 - i\omega_0^n \end{pmatrix} \begin{pmatrix} 1 \\ \rho \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (\text{A.10})$$

we choose $\rho = a_2^2 / c_2 (a_2 + i\omega_0^n)$, and then we get $q(\theta)$. On the other hand, $q^*(s) = M(1, \gamma) e^{i\omega_0^n s}$ is the eigenvector of A^* corresponding to the eigenvalue $-i\omega_0^n$. From the definition of A^* , we have

$$\begin{pmatrix} -bh^* + \frac{c_1 h^* p^*}{(h^* + k_1)^2} + i\omega_0^n & \frac{a_2^2}{c_2} \\ \frac{-c_1 h^*}{h^* + k_1} e^{-i\omega_0^n \bar{\tau}} & -a_2 + i\omega_0^n \end{pmatrix} \begin{pmatrix} M \\ M\gamma \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (\text{A.11})$$

where

$$\gamma = \left(bh^* - \frac{c_1 h^* p^*}{(h^* + k_1)^2} - i\omega_0^n \right) \frac{c_2}{a_2^2}. \quad (\text{A.12})$$

We also assume that $(q^*(s), q(\theta)) = 1$. To obtain the value of M , from (A.9) we have

$$\begin{aligned} & (q^*(s), q(\theta)) \\ &= \bar{M} \left\{ (1, \bar{\gamma}) (1, \rho)^T \right. \\ & \quad \left. - \int_{-1}^0 \int_{\xi=0}^\theta (1, \bar{\gamma}) e^{-i(\xi-\theta)\omega_0^n} d\eta(\theta) (1, \rho)^T e^{i\xi\omega_0^n} d\xi \right\}, \end{aligned} \quad (\text{A.13})$$

then we choose

$$\bar{M} = \left(1 + \rho \bar{\gamma} + \frac{\rho c_1 h^*}{h^* + k_1} \right)^{-1} e^{i\omega_0^n} \quad (\text{A.14})$$

such that $(q^*(s), q(\theta)) = 0$ and $(q^*(s), q(\theta)) = 1$. In other words, let $\Phi = (q(\theta), \bar{q}(\theta))$, $\Psi = (q^*(s), \bar{q}^*(s))^T$, then $(\Psi, \Phi) = I$, and I is unit matrix.

Then the center subspace of model (A.4) is $P = \text{span}\{q(\theta), \bar{q}(\theta)\}$, and the adjoint subspace is $P^* = \text{span}\{q^*(s), \bar{q}^*(s)\}$. Let $v = (v^1, v^2)^T$, where

$$v^1 = (1, 0)^T, \quad v^2 = (0, 1)^T. \quad (\text{A.15})$$

Let $m \cdot v$ be defined by

$$m \cdot v = m_1 v^1 + m_2 v^2, \quad (\text{A.16})$$

for $m = (m_1, m_2)^T \in [-1, 0] \times X$. Hence the center subspace of linear system (A.4) is given by $P_{CN}\mathcal{E}$, where

$$\begin{aligned} P_{CN}\mathcal{E} &= \Phi(\Psi, \langle \phi, v \rangle) \cdot v, \quad \phi \in \mathcal{E}, \\ P_{CN}\mathcal{E} &= \{(q(\theta)z + \bar{q}(\theta)\bar{z}) \cdot v, z \in \mathbb{C}\}, \end{aligned} \quad (\text{A.17})$$

and $\mathcal{E} = P_{CN}\mathcal{E} \oplus P_S\mathcal{E}$, where $P_S\mathcal{E}$ is the stable subspace.

From [17], we know that the infinitesimal generator A_U of linear model (A.4) satisfies $A_U\psi = \dot{\psi}(\theta)$; moreover $\psi \in \text{dom}(A_U)$ if and only if

$$\begin{aligned} & \dot{\psi}(\theta) \in \mathcal{E}, \quad \psi(0) \in \text{dom}(\Delta), \\ & \dot{\psi}(\theta)(0) = \tau_0 \Delta \psi(0) + \tau_0 L_0(\psi). \end{aligned} \quad (\text{A.18})$$

First we define the coordinate to describe the center manifold at $\alpha = 0$; from center manifold we have

$$\begin{aligned} & w(t, \theta) = w(z(t), \bar{z}(t), \theta) \\ &= w_{20}(\theta) \frac{z^2}{2} + w_{11}(\theta) z\bar{z} + w_{02}(\theta) \frac{\bar{z}^2}{2} + \dots. \end{aligned} \quad (\text{A.19})$$

The flow of model (A.2) in the center manifold can be written as follows:

$$U_t = \Phi(z, \bar{z})^T \cdot v + w(z, \bar{z}), \quad (\text{A.20})$$

where

$$\begin{aligned} \dot{z} &= i\omega_0^n z + q^*(0) \langle F(\Phi(z, \bar{z})^T \cdot v + w(z, \bar{z}), 0), v \rangle \\ &= i\omega_0^n z + g(z, \bar{z}) \end{aligned} \quad (\text{A.21})$$

with

$$\begin{aligned} g(z, \bar{z}) &= q^*(0) \langle F(\Phi(z, \bar{z})^T \cdot v + w(z, \bar{z}), 0), v \rangle \\ &= g_{20} \frac{z^2}{2} + g_{11} z\bar{z} + g_{02} \frac{\bar{z}^2}{2} + g_{21} \frac{z^2 \bar{z}}{2} + \dots. \end{aligned} \quad (\text{A.22})$$

Let

$$\begin{aligned} f^1 &= (H + h^*) (a_1 - b(H + h^*)) \\ &\quad - \frac{c_1 (H + h^*) (P(t-1) + p^*)}{k_1 + H + h^*}, \end{aligned}$$

$$f^2 = (P + p^*) \left(a_2 - \frac{c_2 (P + p^*)}{k_2 + H + h^*} \right). \quad (\text{A.23})$$

From above equations, we have

$$\begin{aligned}
 & \frac{g_{20}}{2} \\
 & = \overline{M} \begin{pmatrix} \rho(f_{110}^1 + \bar{\gamma}f_{110}^2) + \rho e^{-i\omega_0^n}(f_{101}^1 + \bar{\gamma}f_{101}^2) \\ + \rho^2 e^{-i\omega_0^n}(f_{011}^1 + \bar{\gamma}f_{011}^2) \\ + \frac{f_{200}^1 + \bar{\gamma}f_{200}^2}{2} + \frac{\rho^2(f_{020}^1 + \bar{\gamma}f_{020}^2)}{2} \\ + \frac{\rho^2 e^{-2i\omega_0^n}(f_{002}^1 + \bar{\gamma}f_{002}^2)}{2} \end{pmatrix}, \\
 & g_{11}(\theta) \\
 & = \overline{M} \begin{pmatrix} \rho(f_{110}^1 + \bar{\gamma}f_{110}^2) + \bar{\rho}(f_{110}^1 + \bar{\gamma}f_{110}^2) \\ + \bar{\rho}e^{i\omega_0^n}(f_{101}^1 + \bar{\gamma}f_{101}^2) + \rho e^{-i\omega_0^n}(f_{101}^1 + \bar{\gamma}f_{101}^2) \\ + \bar{\rho}\rho e^{-i\omega_0^n}(f_{011}^1 + \bar{\gamma}f_{011}^2) + \bar{\rho}\rho e^{i\omega_0^n}(f_{011}^1 + \bar{\gamma}f_{011}^2) \\ + (f_{200}^1 + \bar{\gamma}f_{200}^2) + \rho\bar{\rho}(f_{020}^1 + \bar{\gamma}f_{020}^2) + \rho\bar{\rho}(f_{011}^1 + \bar{\gamma}f_{011}^2) \end{pmatrix}, \\
 & \frac{g_{02}}{2} \\
 & = \overline{M} \begin{pmatrix} \bar{\rho}(f_{110}^1 + \bar{\gamma}f_{110}^2) + \bar{\rho}e^{-i\omega_0^n}(f_{101}^1 + \bar{\gamma}f_{101}^2) \\ + \bar{\rho}^2 e^{-i\omega_0^n}(f_{011}^1 + \bar{\gamma}f_{011}^2) \\ + \frac{f_{200}^1 + \bar{\gamma}f_{200}^2}{2} + \frac{\bar{\rho}^2(f_{020}^1 + \bar{\gamma}f_{020}^2)}{2} \\ + \frac{\bar{\rho}^2 e^{2i\omega_0^n}(f_{002}^1 + \bar{\gamma}f_{002}^2)}{2} \end{pmatrix}, \\
 & \frac{g_{21}}{2} \\
 & = \overline{M} \begin{pmatrix} (f_{110}^1 + \bar{\gamma}f_{110}^2) \left(\rho w_{11}^1(0) + \frac{\bar{\rho}w_{20}^1(0)}{2} \right. \\ \left. + \frac{w_{20}^2(0)}{2} + w_{11}^2(0) \right) \\ + (f_{101}^1 + \bar{\gamma}f_{101}^2) \left(\rho e^{-i\omega_0^n} w_{11}^1(0) + \frac{\bar{\rho}w_{20}^1(0)e^{i\omega_0^n}}{2} \right. \\ \left. + \frac{w_{20}^2(-1)}{2} + w_{11}^2(-1) \right) \\ + (f_{011}^1 + \bar{\gamma}f_{011}^2) \left(\rho e^{-i\omega_0^n} w_{11}^2(0) + \frac{\bar{\rho}w_{20}^2(0)e^{i\omega_0^n}}{2} \right. \\ \left. + \bar{\rho} \frac{w_{20}^2(-1)}{2} + w_{11}^2(-1)\rho \right) \\ + \frac{1}{2}(f_{200}^1 + \bar{\gamma}f_{200}^2)(2w_{11}^1(0) + w_{20}^1(0)) \\ + \frac{1}{2}(f_{020}^1 + \bar{\gamma}f_{020}^2)(2w_{11}^2(0)\rho + w_{20}^2(0)\bar{\rho}) \\ + \frac{1}{2}(f_{002}^1 + \bar{\gamma}f_{002}^2)(2w_{11}^2(-1)e^{-i\omega_0^n}\rho \\ + w_{20}^2(-1)\bar{\rho}e^{-i\omega_0^n}) \end{pmatrix}. \quad (\text{A.24})
 \end{aligned}$$

From [17], we can know that $w(z, \bar{z})$ satisfies

$$\dot{w} = A_U w + H(z, \bar{z}), \quad (\text{A.25})$$

where

$$\begin{aligned}
 H(z, \bar{z}) &= H_{20}(\theta) \frac{z^2}{2} + H_{11}(\theta) z\bar{z} + H_{02}(\theta) \frac{\bar{z}^2}{2} + \dots \\
 &= X_0 F(U_t, 0) - \Phi(\Psi, \langle X_0 F(U_t, 0), v \rangle) \cdot v. \quad (\text{A.26})
 \end{aligned}$$

Again we write near the origin on C_0

$$\dot{w} = \dot{z}w_z + \dot{\bar{z}}w_{\bar{z}}. \quad (\text{A.27})$$

By comparing (A.21) and (A.25) we get

$$\begin{aligned}
 (A_U - 2i\omega_0^n)w_{20}(\theta) &= -H_{20}(\theta), \\
 A_U w_{11}(\theta) &= -H_{11}(\theta). \quad (\text{A.28})
 \end{aligned}$$

When $-1 \leq \theta < 0$, $H(z, \bar{z}) = -\Phi(\theta)\Psi(0)\langle F(U_t, 0), v \rangle \cdot v$. Therefore, for $-1 \leq \theta < 0$,

$$\begin{aligned}
 H_{20}(\theta) &= -[g_{20}q(\theta) + \bar{g}_{02}\bar{q}(\theta)] \cdot v, \\
 H_{21}(\theta) &= -[g_{21}q(\theta) + \bar{g}_{12}\bar{q}(\theta)] \cdot v, \quad (\text{A.29})
 \end{aligned}$$

and for $\theta = 0$, $H(z, \bar{z})(0) = F(U_t, 0) - \Phi(\Psi, \langle F(U_t, 0), v \rangle) \cdot v$, then we obtain

$$\begin{aligned}
 & H_{20}(0) \\
 &= -[g_{20}q(0) + \bar{g}_{02}\bar{q}(0)] \cdot v \\
 &+ \bar{\tau} \begin{pmatrix} \left(\rho(f_{110}^1) + \rho e^{-i\omega_0^n}(f_{101}^1) + \rho^2 e^{-i\omega_0^n}(f_{011}^1) \right. \\ \left. + \frac{f_{200}^1}{2} + \frac{\rho^2 f_{020}^1}{2} + \frac{1}{2}\rho^2 e^{-2i\omega_0^n} f_{002}^1 \right) \\ \left(\rho\bar{\gamma}f_{110}^2 + \rho e^{-i\omega_0^n}(\bar{\gamma}f_{101}^2) + \rho^2 e^{-i\omega_0^n}(\bar{\gamma}f_{011}^2) \right. \\ \left. + \frac{\bar{\gamma}f_{200}^2}{2} + \frac{\bar{\gamma}\rho^2 f_{020}^2}{2} + \frac{1}{2}\rho^2 e^{-2i\omega_0^n} \bar{\gamma}f_{002}^2 \right) \end{pmatrix}, \\
 & H_{11}(0) \\
 &= -[g_{11}q(0) + \bar{g}_{11}\bar{q}(0)] \cdot v \\
 &+ \bar{\tau} \begin{pmatrix} \left(f_{110}^1(\bar{\rho} + \rho) + f_{101}^1(\bar{\rho}e^{i\omega_0^n} + \rho e^{-i\omega_0^n}) \right. \\ \left. + f_{011}^1(\rho\bar{\rho}e^{i\omega_0^n} + \rho\bar{\rho}e^{-i\omega_0^n}) + f_{200}^1 \right. \\ \left. + f_{020}^1\rho\bar{\rho} + f_{002}^1\rho\bar{\rho} \right) \\ \left(f_{110}^2(\bar{\rho} + \rho) + f_{101}^2(\bar{\rho}e^{i\omega_0^n} + \rho e^{-i\omega_0^n}) \right. \\ \left. + f_{011}^2(\rho\bar{\rho}e^{i\omega_0^n} + \rho\bar{\rho}e^{-i\omega_0^n}) + f_{200}^2 \right. \\ \left. + f_{020}^2\rho\bar{\rho} + f_{002}^2\rho\bar{\rho} \right) \end{pmatrix}. \quad (\text{A.30})
 \end{aligned}$$

By the definition of A_U , we have from (A.28)

$$w_{20}(\theta) = \frac{ig_{20}}{\omega_0^n} q(0) e^{i\theta\omega_0^n} \cdot v - \frac{ig_{02}}{3\omega_0^n} \bar{q}(0) e^{-i\theta\omega_0^n} \cdot v + E e^{2i\theta\omega_0^n}, \quad (\text{A.31})$$

where $E = (E^1, E^2) \in R^2$ is a constant vector.

Similarly, we get

$$w_{11}(\theta) = -\frac{ig_{11}}{\omega_0^n} q(0) e^{i\theta\omega_0^n} \cdot v + \frac{i\bar{g}_{11}}{\omega_0^n} \bar{q}(0) e^{-i\theta\omega_0^n} \cdot v + \tilde{E}, \quad (\text{A.32})$$

where $\tilde{E} = (\tilde{E}^1, \tilde{E}^2) \in R^2$ is a constant vector.

Combining (A.5) and (A.28), we obtain

$$H_{20}(0) = 2i\omega_0^n \tilde{\tau} w_{20}(0) - \tilde{\tau} D \Delta w_{20}(0) - \tilde{\tau} L(w_{20}(\theta)), \quad (\text{A.33})$$

therefore

$$\begin{aligned} & \left(\begin{aligned} & \rho(f_{110}^1) + \rho e^{-i\omega_0^n}(f_{101}^1) + \rho^2 e^{-i\omega_0^n}(f_{011}^1) \\ & + \frac{f_{200}^1}{2} + \frac{\rho^2 f_{020}^1}{2} + \frac{1}{2} \rho^2 e^{-2i\omega_0^n} f_{002}^1 \end{aligned} \right) \\ & \left(\begin{aligned} & \rho \bar{\gamma} f_{110}^2 + \rho e^{-i\omega_0^n}(\bar{\gamma} f_{101}^2) + \rho^2 e^{-i\omega_0^n}(\bar{\gamma} f_{011}^2) \\ & + \frac{\bar{\gamma} f_{200}^2}{2} + \frac{\bar{\gamma} \rho^2 f_{020}^2}{2} + \frac{1}{2} \rho^2 e^{-2i\omega_0^n} \bar{\gamma} f_{002}^2 \end{aligned} \right) \\ & = g_{20} q(\theta) + \bar{g}_{02} \bar{q}(\theta) + 2i\omega_0^n w_{20}(\theta) \\ & - \int_{-1}^0 d\eta(\theta) w_{20}(\theta) \\ & = -g_{20} q(0) + \frac{\bar{g}_{02} \bar{q}(0)}{3} + E 2i\omega_0^n \\ & - \int_{-1}^0 d\eta(\theta) \left[\frac{ig_{20}}{\omega_0^n} q(\theta) - \frac{ig_{02}}{3\omega_0^n} \bar{q}(\theta) - E e^{2i\theta\omega_0^n} \right] \\ & = E \left(2i\omega_0^n I - \int_{-1}^0 d\eta(\theta) e^{2i\theta\omega_0^n} \right) \\ & = 2iE\omega_0^n \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ & - \begin{pmatrix} -bh_+ + \frac{c_1 h_+ p_+}{(h_+ + k_1)^2} & \frac{-c_1 h_+}{h_+ + k_1} e^{2i\omega_0^n} \\ \frac{a_2^2}{c_2} & -a_2 \end{pmatrix} E \\ & = \begin{pmatrix} 2i\omega_0^n + bh_+ - \frac{c_1 h_+ p_+}{(h_+ + k_1)^2} & \frac{c_1 h_+}{h_+ + k_1} e^{2i\omega_0^n} \\ -\frac{a_2^2}{c_2} & 2i\omega_0^n + a_2 \end{pmatrix} E. \end{aligned} \quad (\text{A.34})$$

From the above equation we can find the values of E^1 and E^2 . From (A.28) we have $\int_{-1}^0 d\eta(\theta) w_{11}(\theta) = -H_{11}(\theta)$. Therefore

$$\begin{aligned} & \left(\begin{aligned} & (f_{110}^1(\bar{\rho} + \rho) + f_{101}^1(\bar{\rho} e^{i\omega_0^n} + \rho e^{-i\omega_0^n})) \\ & + f_{011}^1(\rho \bar{\rho} e^{i\omega_0^n} + \rho \bar{\rho} e^{-i\omega_0^n}) + f_{200}^1 + f_{020}^1 \rho \bar{\rho} + f_{002}^1 \rho \bar{\rho} \\ & (f_{110}^2(\bar{\rho} + \rho) + f_{101}^2(\bar{\rho} e^{i\omega_0^n} + \rho e^{-i\omega_0^n})) \\ & + f_{011}^2(\rho \bar{\rho} e^{i\omega_0^n} + \rho \bar{\rho} e^{-i\omega_0^n}) + f_{200}^2 + f_{020}^2 \rho \bar{\rho} + f_{002}^2 \rho \bar{\rho} \end{aligned} \right) \\ & = g_{11} q(0) + \bar{g}_{11} \bar{q}(0) - \int_{-1}^0 d\eta(\theta) w_{11}(\theta) \\ & = g_{11} q(0) + \bar{g}_{11} \bar{q}(0) \\ & - \int_{-1}^0 d\eta(\theta) \left[-\frac{ig_{11}}{\omega_0^n} q(0) e^{i\omega_0^n \theta} + \frac{i\bar{g}_{11}}{\omega_0^n} e^{-i\omega_0^n \theta} + \tilde{E} \right] \\ & = - \int_{-1}^0 d\eta(\theta) \tilde{E} \\ & = \begin{pmatrix} bh_+ - \frac{c_1 h_+ p_+}{(h_+ + k_1)^2} & \frac{c_1 h_+}{h_+ + k_1} \\ -\frac{a_2^2}{c_2} & a_2 \end{pmatrix} \tilde{E}. \end{aligned} \quad (\text{A.35})$$

In a similar manner we can compute the corresponding results in $\tilde{E}^1$ and $\tilde{E}^2$. Then g_{21} can be determined. Based on the above analysis, we can see that each g_{ij} can be determined by the parameters. Thus we can compute the following values which determine the direction and stability of bifurcating periodic orbits:

$$\begin{aligned} c_1(0) &= \frac{i}{2\omega_0^n} \left(g_{11} g_{20} - 2|g_{11}|^2 - \frac{|g_{02}|^2}{3} \right) + \frac{g_{21}}{2}, \\ \mu &= -\frac{\text{Re}(c_1(0))}{\text{Re}(\lambda'(\tilde{\tau}))}, \quad \beta_2 = 2 \text{Re}(c_1(0)), \quad (\text{A.36}) \\ T_2 &= -\frac{\text{Im}(c_1(0)) + \mu_2 \text{Im}(\lambda'(\tilde{\tau}))}{\omega_0^n \tilde{\tau}}. \end{aligned}$$

Acknowledgments

The authors would like to thank the anonymous referee for very helpful suggestions and comments which led to improvements of their original paper. This research was supported by Natural Science Foundation of Zhejiang Province (LY12A01014 and LQ12A01009), the National Basic Research Program of China (2012CB426510), and the Fund Project of Zhejiang Provincial Education Department (Y201223449 and Y201120383).

References

- [1] A. F. Nindjin, M. A. Aziz-Alaoui, and M. Cadivel, "Analysis of a predator-prey model with modified Leslie-Gower and Holling-type II schemes with time delay," *Nonlinear Analysis: Real World Applications*, vol. 7, no. 5, pp. 1104–1118, 2006.
- [2] P. H. Leslie, "Some further notes on the use of matrices in population mathematics," *Biometrika*, vol. 35, pp. 213–245, 1948.
- [3] P. H. Leslie, "A stochastic model for studying the properties of certain biological systems by numerical methods," *Biometrika*, vol. 45, pp. 16–31, 1958.
- [4] P. Aguirre, E. González-Olivares, and E. Sáez, "Two limit cycles in a Leslie-Gower predator-prey model with additive Allee effect," *Nonlinear Analysis: Real World Applications*, vol. 10, no. 3, pp. 1401–1416, 2009.
- [5] P. Aguirre, E. González-Olivares, and E. Sáez, "Three limit cycles in a Leslie-Gower predator-prey model with additive Allee effect," *SIAM Journal on Applied Mathematics*, vol. 69, no. 5, pp. 1244–1262, 2009.
- [6] F. Chen, L. Chen, and X. Xie, "On a Leslie-Gower predator-prey model incorporating a prey refuge," *Nonlinear Analysis: Real World Applications*, vol. 10, no. 5, pp. 2905–2908, 2009.
- [7] X. Guan, W. Wang, and Y. Cai, "Spatiotemporal dynamics of a Leslie-Gower predator-prey model incorporating a prey refuge," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 4, pp. 2385–2395, 2011.
- [8] T. Faria, "Stability and bifurcation for a delayed predator-prey model and the effect of diffusion," *Journal of Mathematical Analysis and Applications*, vol. 254, no. 2, pp. 433–463, 2001.
- [9] R. Yafa, F. El Adnani, and H. T. Alaoui, "Limit cycle and numerical simulations for small and large delays in a predator-prey model with modified Leslie-Gower and Holling-type II schemes," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 5, pp. 2055–2067, 2008.
- [10] B. I. Camara and M. A. Aziz-Alaoui, "Dynamics of a predator-prey model with diffusion," *Dynamics of Continuous, Discrete & Impulsive Systems*, vol. 15, no. 6, pp. 897–906, 2008.
- [11] S. Ruan, "On nonlinear dynamics of predator-prey models with discrete delay," *Mathematical Modelling of Natural Phenomena*, vol. 4, no. 2, pp. 140–188, 2009.
- [12] Y. Song, S. Yuan, and J. Zhang, "Bifurcation analysis in the delayed Leslie-Gower predator-prey system," *Applied Mathematical Modelling. Simulation and Computation for Engineering and Environmental Systems*, vol. 33, no. 11, pp. 4049–4061, 2009.
- [13] G.-P. Hu and W.-T. Li, "Hopf bifurcation analysis for a delayed predator-prey system with diffusion effects," *Nonlinear Analysis: Real World Applications*, vol. 11, no. 2, pp. 819–826, 2010.
- [14] W. Zuo and J. Wei, "Stability and Hopf bifurcation in a diffusive predatory-prey system with delay effect," *Nonlinear Analysis: Real World Applications*, vol. 12, no. 4, pp. 1998–2011, 2011.
- [15] F. Dumortier, J. Llibre, and J. C. Artés, *Qualitative Theory of Planar Differential Systems*, Springer, Berlin, Germany, 2006.
- [16] E. González-Olivares, J. Mena-Lorca, A. Rojas-Palma, and J. D. Flores, "Dynamical complexities in the Leslie-Gower predator-prey model as consequences of the Allee effect on prey," *Applied Mathematical Modelling. Simulation and Computation for Engineering and Environmental Systems*, vol. 35, no. 1, pp. 366–381, 2011.
- [17] J. Wu, *Theory and Applications of Partial Functional-Differential Equations*, vol. 119, Springer, New York, NY, USA, 1996.
- [18] Y. Kuang, *Delay Differential Equations with Applications in Population Dynamics*, vol. 191, Academic Press, New York, NY, USA, 1993.
- [19] B. D. Hassard, N. D. Kazarinoff, and Y. H. Wan, *Theory and Applications of Hopf Bifurcation*, vol. 41, 1981.
- [20] J. K. Hale, *Functional Differential Equations*, Springer, 1971.