Abstract

Ras Genes in Human Breast Cancer

Spiros Miyakis, George Sourvinos and
Demetrios A. Spandidos

*Laboratory of Virology, Medical School,
University of Crete, Heraklion, Crete,
Greece*

The expression of *ras* mRNA levels in 27 human sporadic breast cancer specimens was examined, and compared to the corresponding adjacent normal tissue using the RT-PCR technique. Eighteen out of the 27 specimens (67%) exhibited two- to four-fold increased expression of *ras* mRNA levels, compared to corresponding normal tissue. The rates of augmented mRNA expression were similar among the three *ras* genes. A statistically

significant correlation of overexpression of *ras* genes in specimens classified as Stage I disease was observed, compared to tumors in a more advanced stage (II or III). The incidence of codon 12 point mutations of the K-*ras* gene in fresh tissue samples was also assessed in 61 human sporadic breast cancer cases. Point mutations were detected in four (6.5%) out of the 61 cases examined; no correlation was found with any clinicopathological parameter. This is the first report to our knowledge of the differential expression of the *ras* family genes in breast carcinoma. Our findings indicate that the aberrant expression of *ras* genes may be an initial event in breast cancer oncogenesis and that K-*ras* point mutations are rarely involved in the development of mammary neoplasias.