Review

Natural Health Products, Modulation of Immune Function and Prevention of Chronic Diseases

Pierre S. Haddad^{1,2,5}, Georges A. Azar³, Simon Groom⁴ and Michel Boivin^{1,5}

¹Fondation Lucie et André Chagnon, 2001 McGill College, Suite 1000, ²Department of Pharmacology and ³Department of Microbiology and Immunology, Université de Montréal, 2900 Edouard-Montpetit, Montreal, ⁴Department of Human Nutrition, Macdonald College, McGill University, Ste-Anne-de-Bellevue and ⁵Centre Hospitalier de l'Université de Montréal, Hôpital St-Luc, 1058 St-Denis, Montreal, Quebec, Canada

The immune system is increasingly found to be involved in the development of several chronic illnesses, for which allopathic medicine has provided limited tools for treatment and especially prevention. In that context, it appears worthwhile to target the immune system in order to modulate the risk of certain chronic illnesses. Meanwhile, natural health products (NHPs) are generating renewed interest, particularly in the prevention and treatment of several chronic diseases. Over 20 scientists from fields related to immune function and NHPs were thus convened to establish the state of knowledge on these subjects and to explore future research directions. This review summarizes the result of discussions held during the symposium. It thus seeks to be thought provoking rather than to comprehensively cover such broad areas of research. Notably, a brief overview of the immune system is presented, including potentially useful targets and strategies to keep it in an equilibrated state, in order to prevent certain disorders. The pertinence and limitations of targeting the immune system to prevent chronic diseases is also discussed. The paper then discusses the usefulness and limitations of current experimental tools available to study the immune modulating effects of NHPs. Finally, a concise review of some of the most studied NHPs showing promising immunomodulatory activity is given, and avenues for future research are described.

Keywords: NK cells – Cytokines – Microparticles – Probiotics – Natraprevention – Vitamins – n-3 fatty acids – Green tea – Echinacea – Ginseng

Introduction

This concise review presents an expanded summary of discussions held during a symposium entitled 'Maintenance of an efficient and equilibrated immune system through the novel use of natural health products' and that was the object of a recent meeting report (1). The symposium brought together between 20 and 30 scientists in disciplines ranging from herbalism, ethnobotany and phytochemistry, through to basic and clinical research in fields such as immunology, cancer and nutrition. It was held in Montreal, Canada, on September 23 and 24, 2004 under the auspices of the Lucie and André

Chagnon Foundation, Canada's largest philanthropic organization. Participants were asked to discuss the relationship between the modulation of immune system functions and the prevention of chronic diseases, and to explore natural health products (NHPs) that may find novel and judicious use in this context. Other symposium details, including the list of participants, can be found in the published meeting report (1).

The complexity of the immune system is secondary only to that of the central nervous system although the immune system is older in evolutionary terms (2). Our immune system is often conceptualized as possessing two principal components: innate and adaptive immunity, which work in concert to defend the body against infection.

The innate system is ancestral and present in invertebrates as well as vertebrates. It is neither anticipatory (no memory) nor clonal and does not respond to environmental changes. The

For reprints and all correspondence: Pierre S. Haddad, Research Coordinator, Fondation Lucie et André Chagnon, 2001 McGill College Avenue, Montreal, Quebec, Canada H3A 1G1. Tel: +514-380-2001 ext. 1098; Fax: +514-380-8434; E-mail: pierre.haddad@umontreal.ca

innate system reacts directly to a wide variety of microorganisms through receptors for highly conserved and essential microbial molecules, such as lipopolysaccharides present on the cell wall of Gram-negative bacteria and lipoteichoic acids present in Gram-positive bacteria. Indeed, the cells of the innate immune system represent the first line of defence in the immunosurveillance network. The primary cells involved in the identification and spontaneous lysis of offensive targets (virus-infected cells, tumor cells, bone marrow stem cells and embryonic cells) are natural killer (NK) cells, which have been the object of several ground breaking studies in the later part of the last century (3–8). These cells originate from the bone marrow, from which they travel in a unidirectional and almost exclusive way to the spleen; very few NK cells are found in lymph nodes or other secondary sites such as the gut-associated and bronchi-associated lymphatic tissue (3,9–15). Nonetheless, NK cells are lymphoid in morphology and need cells of the macrophage/monocyte lineage for the full expression of NK cell production or function. NK cells are numerically and functionally low in youth and old age, and maximal during the reproductive periods in mammals. Moreover, an inverse relationship exists between the rise and fall of NK cells and the incidence of tumor growth. This phenomenon is believed to be more than coincidental. The studies of Dussault and Miller (16–18) have elucidated some of the factors underlying this relationship. In elderly mice, bone marrow NK cell production and their capacity to bind onto offensive target cells are significantly reduced relative to that of the young adult mammal, thus attenuating any potential for tumor cell lysis. In contrast, very young mammals produce NK cells at a level comparable to that of young adults, but there exist NK-suppressive factors in the micro-environment of the spleen that prevent NK cells from lysing offensive targets. In humans, this corresponds to a time when pediatric leukemias and lymphomas are prevalent. This demonstrates, in part, the importance of NK cells and of the adaptive system in general for the maintenance of optimal immune function.

In contrast, the adaptive system, also termed specific immunity, is acquired through interactions with the environment. It is subject to induction, anticipation (immune memory) and clonal expansion. Adaptive immune responses are central to the body's ability to eliminate bacterial, viral and parasitic infections. Understanding these responses is the key to understanding the phenomena of allergy, autoimmunity, vaccination, cancer and organ graft rejection/acceptance. Adaptive immunity has been known for over a century and an abundant literature exists, which is fully reviewed in classical works such as those of Roitt and Delves (19) and Paul (20). The cells involved in adaptive responses are thymus (T)-derived cytotoxic T lymphocytes and helper T lymphocytes (the latter existing in two subsets: Th1 and Th2), bone (B) marrow-derived lymphocytes and assorted accessory cells (dendritic cells, macrophages and stromal cells). Lymphoid tissue is distributed all over the body. Lymphocytes (along with several of their accessory cells) circulate to and through the

spleen, lymph nodes, specialized lymphoid tissue in the gut (Peyer's Patches and appendix), bronchi and oropharynx (adenoids and tonsils). These 'secondary' sites serve as clonal generating stations for T and B lymphocytes, once the appropriate antigen (from bacterial, viral or parasitic sources) has been encountered. Thus, an army of identical and antigen-specific lymphocytes is uniquely directed at the offensive antigen, ensuring its elimination in normal animals, including humans. Cytotoxic T lymphocytes destroy virus-infected cells by making direct contact with the latter in order to induce cytolysis. Th1 cells, on the other hand, 'activate' the appropriate accessory cell, and it is the latter that then delivers the cytolytic blow. Th1 cells are the target of the AIDS virus, and when the vital Th1 cells are crippled throughout the body (advanced AIDS), opportunist infections ultimately kill the infected host (human or animal). Th2 lymphocytes activate B lymphocytes whose function is to produce an appropriate antibody against the offensive antigen, leading to its destruction and elimination.

The innate and adaptive systems are highly integrated and interdependent (21). In addition to being a requisite for adaptive immunity, the innate system is responsible for the early killing and clearance of infectious pathogens and the resolution of the inflammatory response. The most important means by which the immune system communicates internally and externally is through the use of cytokines, small soluble signaling molecules. Cytokines play a crucial role in the selection, initiation and modulation of an appropriate immune response. The signaling network of cytokines is characterized by its complexity (synergy and antagonism) and by its redundancy (parallelism) that ensure the speed and flexibility required for an effective immune response.

The immune system can give rise to several disorders when it is weakened or overactivated (Table 1). For instance, a defective adaptive immune response can lead to recurrent infection despite previous encounters with antigens, to lowered tolerance to self-organs and tissues leading to organ-specific autoimmunity, and to faulty recognition and elimination of transformed cells leading to cancer. However, an overactivated innate immune response can cause chronic infection because of inefficient clearing of pathogen or chronic inflammation due to an inefficient regulation or resolution of the inflammatory response. Chronic inflammation can lead to extensive non-specific destruction of neighboring tissue. In fact, inflammation and the immune system are intimately tied. In addition,

Table 1. Health effects of imbalances of the immune system

Immune system status	Attributable pathologies
Weak	Prone to opportunistic infectious diseases
	Prone to cancer establishment and tumor escape
Overactivated	Chronic inflammation and autoimmunity (e.g. Type 1 diabetes)
	Heart disease, cancer (e.g. lymphoid)
	Skin disease, allergies and asthma
	Joint and tissue destruction

inflammation is increasingly found to be involved in the development of several chronic diseases such as arteriosclerosis, diabetes, neurodegenerative diseases and even cancer. Immune dysfunction has also been linked to conditions such as chronic pain, anxiety and depression, albeit sometimes as a consequence rather than a cause, and may be involved in other disease processes in a manner that is currently not fully understood (22). Indeed, its diverse influence upon health may be in part due to its evolution as a defence mechanism and in part due to its origins, possibly as a sensory organ (23).

Is the Immune System a Good Target to Aim at In Order to Prevent Chronic Disease? Yes, But...

A first question to address is whether the immune system is a good target to select in the fight to prevent chronic diseases. Consensus is generally positive with some important reservations being brought forward. In fact, it is accepted that the immune system represents our arsenal to combat infection and disease. As mentioned, an underactivated immune system increases risks of opportunistic infection and uncontrolled neoplastic tissue growth, while an overactivated immune system leads to many afflictions, including inflammation, allergy and autoimmunity. Hence, an unchecked and unbalanced immune system hampers good health and quality of life. This explains the consensus 'Yes' given as an answer by the experts regarding the pertinence of targeting the immune system to prevent chronic disease.

However, it must be noted that the immune system carries out its important defensive function in intimate and coordinated interaction with the nervous and endocrine systems (24,25). The innate complexity of the immune system and of its communication tool, the cytokines, is thus further complicated by these interactions with the nervous and endocrine systems. Some concern also arises when considering that targeting the immune system may be too reductionist an approach, particularly when dealing with complex pathologies like cancer. These considerations form the basis of the 'but...' part of the answer given in the title to this subsection.

Promising Research Avenues and Markers

It is important to consider that invertebrate species lack adaptive immunity but are clearly able to deal effectively with infections. It is also interesting to note that these organisms do not appear to develop cancer or autoimmune disease (at least not in any readily recognizable form), but they do demonstrate senescence (26–28). Therefore, the apparent absence of these age-related diseases in invertebrates could support an evolutionary argument to focus disease prevention upon maintenance of efficient and equilibrated innate immunity.

If this line of reasoning is correct, then the immune cells of greatest interest are ones of the NK cell lineage described

above. These immune cells are present in invertebrate and vertebrate species and serve as a link between innate and adaptive immunity in vertebrates, as mentioned (25). Indeed, NK cell function has been identified as a reliable measure of immunotoxicity for use in pharmaceutical safety testing (29) and could perhaps provide a means to monitor immune system equilibrium.

Another important point to consider relates to the problem of markers for immune system function. It is difficult to reach a consensus on the appropriate markers to use in order to monitor the maintenance of a healthy immune system, primarily because biological heterogeneity precludes readily identifiable thresholds for concern in the prevention of chronic disease. However, there exist potential novel avenues to explore, including the use of NK cell activity, as mentioned above, or the use of a constellation of markers, of which cytokines and cytokine receptors would be a major part. Much has been written about the validity and validation of biomarkers for health monitoring (30). While it is beyond the scope of this paper to discuss these issues at length, it is worth highlighting some general considerations. Firstly, biomarkers must be relevant to the question being asked and practical to use in the desired setting since advances in analytical sensitivity can now provide detection limits at concentrations often below those that have biological importance (31). Secondly, it is generally easier to define thresholds for intervention (i.e. when a disease requires treatment) than for early detection or prevention (i.e. before onset of the disease). Indeed, thresholds for disease prevention require a good definition of what constitutes the 'normal' condition. However, biological heterogeneity and the redundancy present within the immune system makes the establishment of precise 'normal ranges' especially difficult. Therefore, although NK cell activity and cytokine assays can be used as reliable measures for treatment intervention (32,33) they have yet to be shown to be practical biomarkers for defining a threshold for the early detection of chronic illnesses.

Finally, one interesting alternative to conventional immune system biomarkers is the presence and quality of microparticles present in the blood. Such microparticles, including exosomes, originate from several cells, including some prominent immune cells, and may represent a novel type of intercellular signaling mechanism (34,35). Hence, research into the development of novel and pertinent biomarkers will certainly contribute to finding ways to modulate the immune system and prevent certain chronic diseases.

Tools to Study the Immune System and NHPs

In this section we briefly review several experimental models and approaches that have been used to study immunomodulatory NHPs. Table 2 summarizes the advantages and limitations of each modality. Human clinical studies remain the gold standard upon which scientists and health care professionals can rely to determine the usefulness and efficacy of a given NHP. Nonetheless, animal models, *in vitro* bioassays, and

Table 2. Tools to study NHPs as related to the immune system

Experimental models/tools	Pros	Cons
<i>In vitro</i> measurement on cell culture	Can target specific effects of the studied product on specific cell types Very helpful for investigating cellular and molecular mechanisms Large number of very well characterized systems with efficient markers of activation/inhibition	Simplistic model, systemic effects not taken into consideration NHP administration generally not physiological
Genomics/proteomics	New analytical tools Very precise results Can be used in fundamental research as well as clinical trials	Cost of utilization Availability of source material
Animal models	Mouse has a very well characterized genome Complete living organism Genetically modifiable (transgenic and knock out mice) Large number of well characterized models for several human pathologies	Despite high similarities with man, uncertainty whether mice and men react the same way Difficulty in measuring environmental components on a mouse
Clinical trials	Gold standard for safety and efficacy Direct and systemic effects measured on health	Weakness or lack of proper markers to evaluate success of intervention Improper administration/standardization of NHPs
Epidemiological studies	Large-scale studies on the population Can include both the environmental and germ line components of disease	Challenge in scientifically measuring dietary exposure in the population Large cohorts needed for significant conclusions

modern proteomic and genomic technologies can be most helpful in delineating the mode of action of NHPs. Finally, epidemiological studies bring a wealth of information on the influence of genetic (germ line) and environmental factors on health outcomes in the general population and pertinent subgroups.

In addition, it is generally recognized that a great effort should be placed on the standardization of methods used to study immune modulation by NHPs. In terms of herbals, this also includes a crucial component of standardization of the herbal preparations themselves. As mentioned in Table 2 above, the greatest inconsistencies of clinical research related to immunomodulatory NHPs are caused in major part by the lack of quality control of the source materials used (36). The US agency NCCAM came to a similar conclusion after its first 5 years of functioning and is now focusing on more preclinical studies to prepare better clinical studies, notably with herbal products (<http://nccam.nih.gov/about/plans/2005/index.htm>).

NHPs having Immunomodulatory Activity

During the last century, modern medicine has contributed major advancements to sanitation and public health initiatives and discoveries like antibiotics and vaccination. These efficiently hindered the progression of infectious diseases through the population. In fact, modern medicine has succeeded best in treating acute illnesses. It is in the realm of chronic diseases that modern medicine and drugs have been more disappointing and that NHPs have elicited resurgent interest. In this section,

we review the evidence for the immunomodulatory action of several NHPs, including probiotics, products of animal origin (principally milk and thymic extracts), nutritional factors (especially vitamins, minerals and fatty acids) and herbal products. Table 3 summarizes the strongest evidence brought forward for some of the most promising NHPs. Readers are also oriented to recent reviews (37,38) for complementary information.

The gastrointestinal (GI) tract and the skin epidermis are the major barriers humans possess to confront the external environment. Both are responsible for maintaining systemic integrity and use lymph nodes as efficient immune checkpoints. The GI tract distinguishes itself by holding the greatest number of lymph nodes, and also by being the most intricate and complex immune organ. Probiotics are live bacteria that colonize the GI tract. They have received a lot of research attention, particularly in clinical studies in children and the elderly (39–42). Indeed, they positively affect general health going from the development of the immune system in infants right after birth to the enhancement of general immunity in the elderly (critically exposed to immunosenescence). Probiotics offer much promise because they appear not only to help with the maturation of the visceral immune system but also to equilibrate the body's defence mechanisms. In that sense, they act as true immunomodulating agents, enhancing immunity when it is weakened or developing, and buffering exaggerated reactions (e.g. allergies and inflammatory bowel disease). They also represent the only NHP category that research has shown to have well established curing

Table 3. Promising NHPs exhibiting modulatory effects on the immune system

NHP source	Prevention	Treatment
Probiotics	Helps immune system maturation (96) Helps prevent allergies (96), atopic disease and hypersensitivities in infants (98,99) Enhances immunity in the elderly (102,96)	Helps manage atopic eczema/dermatitis (97) Promise in treating inflammatory bowel disease (IBD) through immune modulation (100,101)
Green Tea (EGCG, flavonoids ...)	Daily intake could prevent tumor growth (65,103,104)	Potent antiangiogenic effect (66–68) Potent anti-inflammatory effect (105,106)
Vitamins C and E, n-3 fatty acids	Antioxidant: decrease oxidative stress in several chronic diseases (73) Anti-inflammatory: n-3 fatty acids (74)	Helps manage arthritis, asthma and IBD (76,75)
<i>Echinacea</i>	Enhance antitumor immunity (mouse NK cells) (107) Enhance NK cell immunity in aged mice (107)	May help treat upper respiratory tract infection (108) Putative antifungal properties (109) and antiviral properties (110)
Ginseng (ginsenosides)	Immunosuppressant properties (111) Anti-inflammatory properties (113)	May have benefit as an adjunct to cancer therapy (112)
Ginseng (polysaccharides)	Immune stimulating properties (NK, macrophages and antibody production) (114)	Helps clear infections agents like <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> and <i>Candida albicans</i> in mice (115)

as well as preventive properties. Probiotics exert their actions principally by

1. Improving the intestinal immunologic barrier function through spatial exclusion of other pathogenic microbial that could otherwise colonize the gut (43,44).
2. Restoring normal intestinal permeability (45) in the face of inflammatory or allergic diseases by still ill-defined mechanisms that include cell wall and DNA components playing the role of immunomodulatory products (46) and by lowering the production of pro-inflammatory cytokines (47,48).
3. Synthesizing vitamins (folic acid, biotin and other group B vitamins, vitamin K and so on) (116).

Suggestions have been made that probiotics may absorb heavy metals (49–51) and thus help their elimination through the feces. Probiotics may also promote the production of enzymes that help decrease the severity of certain food intolerances/allergies (52). However, these actions have not been fully documented.

Data on milk peptides, bovine colostrum and thymic extracts are less convincing in terms of immune modulation, most of the research having been done in cellular models *in vitro* and showing contradicting results (53–64). Several peptides released from enzymatic hydrolysis of milk proteins were reported to modulate immune functions such as phagocytosis, lymphocyte proliferation and cytokine production. The type of enzymatic treatment has an influence on the immunomodulatory functions of peptides generated from milk proteins and, without standardized procedures, it is difficult to draw conclusions from those studies. Further research, especially clinical trials, will be required to identify the impact of those products on the immune system in humans.

However, stronger evidence was presented to support the usefulness of a number of micronutrients to prevent certain chronic illnesses. A new term coined ‘nutraprevention’ is

applied to the use of empirical combinations of foods, spices and other food-grade herbals in the fight against cancer (65). So far, this approach is based on epidemiological studies clearly showing relationships between cancer prevalence and food consumption in different populations around the world. Moreover, research is now showing potent antiangiogenic activity for polyphenolic components of green tea (66–68); angiogenesis being the process by which growing tumors promote the development of new blood vessels to ensure their survival. Several plant constituents, including soy isoflavones (69,70) and compounds in cruciferous vegetables (71,72), are also known to modulate enzymes involved in xenobiotic biotransformation, namely, the activation or detoxification of ‘foreign’, pro-carcinogenic chemicals. Vitamin C has been clearly linked to the modulation of immune function, particularly in the context of novel research showing improved cognitive status in patients with neurodegenerative disorders (73), which are increasingly shown to involve an immune dysfunction component. Similarly, polyunsaturated fatty acids, in particular the now famous ω -3 fatty acids, have been used successfully in the management and prevention of several inflammatory and allergic diseases (74–76). However, clinical research on lipids has yielded inconsistent results concerning their potential to modulate immune system function. This state of affairs is related to poor study design and to the great variability in immune cells selected and methods used to assess immune function.

Intake of green tea, *Brassica* vegetables (e.g. broccoli, cabbage and others) and soy foods has been shown to be associated with lower risk of several cancers in epidemiologic studies [reviewed in (77–82)]. Furthermore, data from experimental studies suggest that green tea and its constituents, such as isoflavones, may modulate markers of inflammation and immune function (83–87). A limited number of herbal products also show promise in the realm of immune modulation.

It is notably the case with *Echinacea* that has received significant scientific attention in the past two decades, particularly in studies on upper respiratory tract infections (36,88,89). Among its many immune modulating activities reported, activation of macrophages has been demonstrated most convincingly. However, novel mechanisms of action include effects on B and T lymphocytes as well as NK cells (90,91). In particular, Currier and Miller (90) observed that upon incorporating *Echinacea* in the daily diet of elderly mice for only 14 days NK cell production in the bone marrow, as well as the number of mature functional NK cells reaching the spleen, increased to levels found in young adults (92). Brousseau and Miller (93) have also revealed that daily dietary intake of *Echinacea* throughout life, beginning in youth, increased the number of individuals reaching old age to 74% as opposed to 46% in control animals. Thus, any herbal treatment that would augment NK cells would clearly be worthy of investigation for its therapeutic/prophylactic potential, especially in the realm of immune modulation.

Ginseng is the other herbal that shows much promise as an immune modulating NHP. Indeed, it appears to have both immunostimulant and immunosuppressant activities, which would lend credence to the view that ginseng is an 'adaptogenic' NHP. Immunostimulant activities have been attributed to polysaccharide fractions, notably acting on both macrophages and B lymphocytes (94,95). However, ginsenosides display immunosuppressant activity *in vitro* by inhibiting cytokine release in activated macrophages (I. Lemaire, unpublished results). Hence, further research is warranted on ginseng and other adaptogenic herbals to define their potential use in preventing chronic diseases related to immune dysfunction.

Summary and Conclusion

Dysfunction of the immune system is clearly implicated in the development of several chronic diseases. Hence, attempting to maintain an efficient and equilibrated immune system is a valid approach to prevent certain chronic illnesses. However, this approach is limited by the great complexity of the immune system and its close interactions with the nervous and endocrine functions. This is further hampered by the difficulty in identifying reliable and convenient analytical tools to assess immune function and its modulation. In that context, interesting areas for future research include NK cells, constellation of cytokine markers and receptors and assessment of blood-borne microparticles. Despite these limitations, research on NHPs has shown that certain products offer novel avenues to modulate the immune system and help prevent chronic diseases. The most promising ones are probiotics, nutraceuticals (combinations of biologically-active fruits, vegetables and spices) and so-called adaptogenic plants (e.g. *echinacea* and ginseng). Notwithstanding these encouraging results, future research efforts with NHPs will need to address the important issues of standardization in both product quality and research methodology. Among areas to favor in the research agenda on NHPs and immune modulation, this review highlights the

following needs: (i) studies on the interaction between NHPs and immune cells, particularly the influence of NHPs on the innate immune system; (ii) epidemiological studies and clinical studies with NHPs in chronic illnesses where immune dysfunction may play a role. Finally, social/economic impact of NHPs in health care must also be considered.

Acknowledgements

This review is based on discussions held in September 2004 during a 2 day symposium organized and funded by the Lucie et André Chagnon Foundation in Montreal, Canada. The symposium is the object of a short meeting report in *eCAM* (1). The authors express their gratitude to all participants for valuable comments and suggestions. Special thanks go to Yvan Boutin (Lévis-Lauzon College, Lévis, Canada), Edwin Cooper (UCLA, Los Angeles, CA, USA), Johanna Lampe (Fred Hutchison Cancer Research Center, Seattle, WA, USA), Sandra Miller (McGill University, Montreal, Canada) and Denis Roy (Laval University, Quebec City, Canada). Ms Geneviève Côté and Geneviève Beauchamp are heartfully thanked for devoting their time and innovative ideas to make the symposium organization such a success. We gratefully acknowledge the expert secretarial help of Ms Laure Pinto.

References

1. Haddad PS, Beauchamp G, Côté G, Boivin M. Maintenance of an efficient, equilibrated immune system through the novel use of natural health products: synopsis of a symposium. *eCAM* 2005;2:237-8.
2. Cooper E. Commentary on CAM and NK cells by Kazuyoshi Takeda and Ko Okumura. *eCAM* 2004;1:29-34.
3. Keissling R, Klein E, Pross H, Wigzell H. Natural killer cells in the mouse. II. Cytotoxic cells with specificity for mouse Moloney leukemia cells: characteristics of the killer cell. *Eur J Immunol* 1975;5:117-21.
4. Herberman RB, Nunn ME, Larvin DH. Natural cytotoxic reactivity of mouse lymphoid cells against syngeneic and allogeneic tumors. I. Distribution of reactivity and specificity. *Int J Cancer* 1975;16:216-29.
5. Miller SC. Genetically determined resistance to foreign bone marrow transplantation in mice: characterization of the effector cells. *J Immunol* 1983;131:92-7.
6. Hackett J, Tutt M, Lipscomb M, Bennett M, Koo G, Kumar V. Origin and differentiation of natural killer cells. II. Functional and morphologic studies of purified NK 1.1+ cells. *J Immunol* 1986;136:3124-31.
7. Krishnaraj R. Immunosenescence of human NK cells: effects on tumor target recognition, lethal hit and interferon sensitivity. *Immunol Lett* 1992;34:79-84.
8. Kutza J, Murasko DM. Effects of aging on natural killer cell activity and activation by interleukin-2 and IFN- α . *Cell Immunol* 1994;155:195-204.
9. Haller O, Wigzell H. Suppression of natural killer cell activity with radioactive strontium: effector cells are marrow dependent. *J Immunol* 1977;118:1503-6.
10. Seaman WE, Blackman MA, Gindhart TD, Roubinian JR, Loeb JM, Talal N. Estradiol reduces natural killer cells in mice. *J Immunol* 1978;121:2193-8.
11. Riccardi C, Barlozzari T, Santoni A, Herberman RB, Cesarini C. Transfer to cyclophosphamide-treated mice of natural killer (NK) cells and *in vivo* natural reactivity against tumors. *J Immunol* 1981;126:1284-7.
12. Koo GC, Cayre Y, Mittl LR. Comparative study of splenic natural killer cells and prothymocytes. *Cell Immunol* 1981;62:164-71.
13. Koo GC, Manyak CL. Generation of cytotoxic cells from murine bone marrow by human recombinant IL-2. *J Immunol* 1986;137:1751-6.
14. Kalland T. Interleukin 3 is a major negative regulator of the generation of natural killer cells from mouse bone marrow precursors. *J Immunol* 1986;137:2268-71.

15. Pollack SB, Rosse C. The primary role of murine bone marrow in the production of natural killer cells. *J Immunol* 1987;139:2149–56.
16. Dussault I, Miller SC. Decline in natural killer cell-mediated immunosurveillance in aged mice—a consequence of reduced cell production and tumor binding capacity. *Mech Ageing Dev* 1994;75:115–29.
17. Dussault I, Miller SC. The effectiveness of immunotherapy in aged leukemic mice. *Gerontology* 1995;41:195–204.
18. Dussault I, Miller SC. Effect on leukemia cell numbers of in vivo administration of immunotherapeutic agents is age-dependent. *Oncology* 1996;53:241–6.
19. Roitt IM, Delves PJ (eds). *Encyclopedia of Immunology*, 3rd edition. London/San Diego: Academic Press, 1992.
20. Paul WE (ed). *Fundamental Immunology*, 4th edition. Philadelphia: Lippincott Williams & Wilkins Publishers, 1998.
21. Hoebe K, Janssen E, Beutler B. The interface between innate and adaptive immunity. *Nat Immunol* 2004;5:971–4.
22. Ewald PW. Plague time: how stealth infections cause cancers, heart disease and other deadly ailments. New York: The Free Press, 2002, 282.
23. Blalock JE. The immune system as a sensory organ. *J Immunol* 1984;132:1067–70.
24. Haddad JJ, Saade NE, Safieh-Garabedian B. Cytokines and neuro-immune-endocrine interactions: a role for the hypothalamic-pituitary-adrenal revolving axis. *J Neuroimmunol* 2002;133:1–19.
25. Takeda K, Okumura K. CAM and NK cells. *eCAM* 2004;1:17–27.
26. Flatt T. Assessing natural variation in genes affecting *Drosophila* lifespan. *Mech Ageing Dev* 2004;125:155–9.
27. Luo Y. Long-lived worms and aging. *Redox Rep* 2004;9:65–9.
28. Singh K. Mitochondrial dysfunction is a common phenotype in aging and cancer. *Ann NY Acad Sci* 2004;1019:260–4.
29. USFDA. Immunotoxicology evaluation of investigational new drugs, Guidance for Industry Document issued by the Center for Drug Evaluation and Research.35.
30. World Health Organization. Environmental Health Criteria 222: Biomarkers In Risk Assessment: Validity And Validation, Report published under the joint sponsorship of the United Nations Environment Program, the International Labor Organization, and the World Health Organization, and produced within the framework of the Inter-Organization Program for the Sound Management of Chemicals.
31. Pritzker KPH. Cancer biomarkers: easier said than done. *Clin Chem* 2002;48:1147–50.
32. Kelley DS, Taylor PC, Nelson GJ, Schmidt PC, Mackey BE, Kyle D. Effects of dietary arachidonic acid on human immune response. *Lipids* 1997;32:449–56.
33. Kelley DS, Taylor PC, Nelson GJ, Mackey BE. Dietary docosahexaenoic acid and immunocompetence in young healthy men. *Lipids* 1998;33:559–66.
34. Denzer K, Kleijmeer MJ, Heijnen HF, Stoorvogel W, Geuze HJ. Exosome: from internal vesicle of the multivesicular body to intercellular signalling device. *J Cell Sci* 2000;113:3365–74.
35. Freyssinet JM. Cellular microparticles: what are they bad or good for? *J Thromb Haemost* 2003;1:1655–62.
36. Goel V, Lovlin R, Barton R, Lyon MR, Bauer R, Lee TD, et al. Efficacy of a standardized *Echinacea* preparation (*Echinilin*) for the treatment of the common cold: a randomized, doubled-blind, placebo-controlled trial. *J Clin Pharm Ther* 2004;29:75–83.
37. Plaeger SF. Clinical immunology and traditional herbal medicines. *Clin Diagn Lab Immunol* 2003;10:337–8.
38. Patwardhan B, Gautam M. Botanical immunodrugs: scope and opportunities. *Drug Discov Today* 2005;10:495–502.
39. Shi HN, Walker A. Bacterial colonization and the development of intestinal defence. *Can J Gastroenterol* 2004;18:493–500.
40. Marteau P, Seksik P, Lepage P, Dore J. Cellular and physiological effects of probiotics and prebiotics. *Mini Rev Med Chem* 2004;4:889–96.
41. Brown AC, Valiere A. Probiotics and medical nutrition therapy. *Nutr Clin Care* 2004;7:56–68.
42. Bruzzese E, Canani RB, De Marco G, Guarino A. Microflora in inflammatory bowel diseases: a pediatric perspective. *J Clin Gastroenterol* 2004;38:S91–3.
43. Resta-Lenert S, Barrett KE. Live probiotics protect intestinal epithelial cells from the effects of infection with enteroinvasive *Escherichia coli* (EIEC). *Gut* 2003;52:988–97.
44. Jenkins B, Holsten S, Bengmark S, Martindale R. Probiotics: a practical review of their role in specific clinical scenarios. *Nutr Clin Pract* 2005;20:262–70.
45. Madsen K, Cornish A, Soper P, McKaigney C, Jijon H, Yachimec C, et al. Probiotic bacteria enhance murine and human intestinal epithelial barrier function. *Gastroenterology* 2001;121:580–91.
46. Jijon H, Backer J, Diaz H, Yeung H, Thiel D, McKaigney C, et al. DNA from probiotic bacteria modulates murine and human epithelial and immune function. *Gastroenterology* 2004;126:1358–73.
47. Madsen KL, Doyle JS, Jewell LD, Tavernini MM, Fedorak RN. Lactobacillus species prevents colitis in interleukin 10 gene-deficient mice. *Gastroenterology* 1999;116:1107–14.
48. Lammers KM, Helwig U, Swennen E, Rizzello F, Venturi A, Caramelli E, et al. Effect of probiotic strains on interleukin 8 production by HT29/19A cells. *Am J Gastroenterol* 2002;97:1182–6.
49. Summers AO, Wireman J, Vimy MJ, Lorscheider FL, Marshall B, Levy SB, et al. Mercury released from dental “silver” fillings provokes an increase in mercury- and antibiotic-resistant bacteria in oral and intestinal floras of primates. *Antimicrob Agents Chemother* 1993;37:825–34.
50. Lorscheider FL, Vimy MJ, Summers AO, Zwiers H. The dental amalgam mercury controversy—inorganic mercury and the CNS; genetic linkage of mercury and antibiotic resistances in intestinal bacteria. *Toxicology* 1995;97:19–22.
51. Brudnak MA. Probiotics as an adjuvant to detoxification protocols. *Med Hypotheses* 2002;58:382–5.
52. de Vrese M, Stegelmann A, Richter B, Fenselau S, Laue C, Schrezenmeir J. Probiotics—compensation for lactase insufficiency. *Am J Clin Nutr* 2001;73:421S–9S.
53. Migliore-Samour D, Floc’h F, Jolles P. Biologically active casein peptides implicated in immunomodulation. *J Dairy Res* 1989;56:357–62.
54. Torre PM, Oliver SP. Suppression of mitogenic response of peripheral blood mononuclear cells by bovine mammary secretions. *J Dairy Sci* 1989;72:219–27.
55. Fiat AM, Migliore-Samour D, Jolles P, Drouet L, Bal dit Sollier C, Caen J. Biologically active peptides from milk proteins with emphasis on two examples concerning antithrombotic and immunomodulating activities. *J Dairy Sci* 1993;76:301–10.
56. Schlimme E, Meisel H. Bioactive peptides derived from milk proteins. Structural, physiological and analytical aspects. *Nahrung* 1995;39:1–20.
57. Lahov E, Regelson W. Antibacterial and immunostimulating, casein-derived substances from milk: casecidin, isracidin peptides. *Food Chem Toxicol* 1996;34:131–45.
58. Cross ML, Gill HS. Immunomodulatory properties of milk. *Br J Nutr* 2000;84:81.
59. Shah NP. Effects of milk-derived bioactives: an overview. *Br J Nutr* 2000;84:S3–10.
60. Low PP, Rutherford KJ, Gill HS, Cross ML. Effect of dietary whey protein concentrate on primary and secondary antibody responses in immunized BALB/c mice. *Int Immunopharmacol* 2003;3:393–401.
61. Legrand D, Ellass E, Pierce A, Mazurier J. Lactoferrin and host defence: an overview of its immuno-modulating and anti-inflammatory properties. *Biometals* 2004;17:225–9.
62. Li EW, Mine Y. Immunoenhancing effects of bovine glycomacropeptide and its derivatives on the proliferative response and phagocytic activities of human macrophagelike cells, U937. *J Agric Food Chem* 2004;52:2704–8.
63. Marshall K. Therapeutic applications of whey protein. *Altern Med Rev* 2004;9:136–56.
64. Mercier A, Gauthier S, Fliss I. Immunomodulating effects of whey proteins and their enzymatic digests. *Int Dairy J* 2004;14:175.
65. Beliveau R, Gingras D. Green tea: prevention and treatment of cancer by nutraceuticals. *Lancet* 2004;364:1021–2.
66. Lamy S, Gingras D, Beliveau R. Green tea catechins inhibit vascular endothelial growth factor receptor phosphorylation. *Cancer Res* 2002;62:381–5.
67. Cao Y, Cao R. Angiogenesis inhibited by drinking tea. *Nature* 1999;398:381.
68. Cao Y, Cao R, Brakenhielm E. Antiangiogenic mechanisms of diet-derived polyphenols. *J Nutr Biochem* 2002;13:380–90.
69. Kishida T, Beppu M, Nashiki K, Izumi T, Ebihara K. Effect of dietary soy isoflavone aglycones on the urinary 16 α -to-2-hydroxysterone ratio in C3H/HeJ mice. *Nutr Cancer* 2000;38:209–14.
70. Chan HY, Leung LK. A potential protective mechanism of soya isoflavones against 7,12-dimethylbenz[a]anthracene tumour initiation. *Br J Nutr* 2003;90:457–65.

71. Gill CI, Haldar S, Porter S, Matthews S, Sullivan S, Coulter J, et al. The effect of cruciferous and leguminous sprouts on genotoxicity, in vitro and in vivo. *Cancer Epidemiol Biomarkers Prev* 2004;13:1199–205.
72. Zhang Y. Cancer-preventive isothiocyanates: measurement of human exposure and mechanism of action. *Mutat Res* 2004;555:173–90.
73. Martin A. Antioxidant vitamins E and C and risk of Alzheimer's disease. *Nutr Rev* 2003;61:69–73.
74. Kelley DS. Modulation of human immune and inflammatory responses by dietary fatty acids. *Nutrition* 2001;17:669–73.
75. Simopoulos AP. Omega-3 fatty acids in inflammation and autoimmune diseases. *J Am Coll Nutr* 2002;21:495–505.
76. Endres S, Lorenz R, Loeschke. Lipid treatment of inflammatory bowel disease. *Curr Opin Clin Nutr Metab Care* 1999;2:117–20.
77. Murillo G, Mehta RG. Cruciferous vegetables and cancer prevention. *Nutr Cancer* 2001;41:17–28.
78. Kristal AR, Lampe JW. Brassica vegetables and prostate cancer risk: a review of the epidemiological evidence. *Nutr Cancer* 2002;42:1–9.
79. Le Marchand L. Cancer preventive effects of flavonoids. *Biomed Pharmacother* 2002;56:296–301.
80. Sarkar FH, Li Y. Soy isoflavones and cancer prevention. *Cancer Invest* 2003;21:744–57.
81. Limer JL, Speirs V. Phyto-estrogens and breast cancer chemoprevention. *Breast Cancer Res* 2004;6:119–27.
82. Park OJ, Surh YJ. Chemopreventive potential of epigallocatechin gallate and genistein: evidence from epidemiological and laboratory studies. *Toxicol Lett* 2004;150:43–56.
83. Paradkar PN, Blum PS, Berhow MA, Baumann H, Kuo SM. Dietary isoflavones suppress endotoxin-induced inflammatory reaction in liver and intestine. *Cancer Lett* 2004;215:21–8.
84. Verdrengh M, Jonsson IM, Holmdahl R, Tarkowski A. Genistein as an anti-inflammatory agent. *Inflamm Res* 2003;52:341–6.
85. Surh YJ, Chun KS, Cha HH, Han SS, Keum YS, Park KK, et al. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF-kappa B activation. *Mutat Res* 2001;480:243–68.
86. Katiyar SK, Challa A, McCormick TS, Cooper KD, Mukhtar H. Prevention of UVB-induced immunosuppression in mice by the green tea polyphenol (–)-epigallocatechin-3-gallate may be associated with alterations in IL-10 and IL-12 production. *Carcinogenesis* 1999;20:2117–24.
87. Jenkins DJ, Kendall CW, Connelly PW, Jackson CJ, Parker T, Faulkner D, et al. Effects of high- and low-isoflavone (phytoestrogen) soy foods on inflammatory biomarkers and proinflammatory cytokines in middle-aged men and women. *Metabolism* 2002;51:919–24.
88. Block KI, Mead MN. Immune system effects of Echinacea, ginseng and astragalus: a review. *Integr Cancer Ther* 2003;2:247–67.
89. Barrett B. Medicinal properties of Echinacea: a critical review. *Phytomedicine* 2003;10:66–86.
90. Currier NL, Miller SC. Natural killer cells from aging mice treated with extracts from *Echinacea purpurea* are quantitatively and functionally rejuvenated. *Exp Gerontol* 2000;35:627–39.
91. Currier NL, Miller SC. *Echinacea purpurea* and melatonin augment natural-killer cells in leukemic mice and prolong life span. *J Altern Complement Med* 2001;7:241–51.
92. Sun LZ-Y, Currier NL, Miller SC. The American coneflower: a prophylactic role involving nonspecific immunity. *J Altern Complement Med* 1999;5:437–46.
93. Brousseau M, Miller SC. Enhancement of natural killer cells and increased survival of aging mice fed daily Echinacea root extract from youth. *Biogerontology* 2005; 6: 157–163.
94. Wang M, Guilbert LJ, Ling L, Li J, Wu Y, Xu S, et al. Immunomodulating activity of CVT-E002, a proprietary extract from North American ginseng (*Panax quinquefolium*). *J Pharm Pharmacol* 2001;53:1515–23.
95. Assinewe VA, Amason JT, Aubry A, Mullin J, Lemaire I. Extractable polysaccharides of *Panax quinquefolius* L. (North American ginseng) root stimulate TNF alpha production by alveolar macrophages. *Phytomedicine* 2002;9:398–404.
96. Cross ML. Immunoregulation by probiotic lactobacilli: pro-Th1 signals and their relevance to human health. *Clin Appl Immunol Rev* 2002;3:115–25.
97. Viljanen M, Savilahti E, Haahtela T, Juntunen-Backman K, Korpela R, Poussa T, et al. Probiotics in the treatment of atopic eczema/dermatitis syndrome in infants: a double-blind placebo-controlled trial. *Allergy* 2005;60:494–500.
98. Majamaa H, Isolauri E. Probiotics: a novel approach in the management of food allergy. *J Allergy Clin Immunol* 1997;99:179–85.
99. Ogden NS, Bielory L. Probiotics: a complementary approach in the treatment and prevention of pediatric atopic disease. *Curr Opin Allergy Clin Immunol* 2005;5:179–84.
100. Fedorak RN, Madsen KL. Probiotics and the management of inflammatory bowel disease. *Inflamm Bowel Dis* 2004;10:286–99.
101. Kanauchi O, Matsumoto Y, Matsumura M, Fukuoka M, Bamba T. The beneficial effects of microflora, especially obligate anaerobes, and their products on the colonic environment in inflammatory bowel disease. *Curr Pharm Des* 2005;11:1047–53.
102. Gill HS, Rutherford KJ, Cross ML, Gopal PK. Enhancement of immunity in the elderly by dietary supplementation with the probiotic *Bifidobacterium lactis* HN019. *Am J Clin Nutr* 2001;74:833–9.
103. Greenwald P. Clinical trials in cancer prevention: current results and perspectives for the future. *J Nutr* 2004;134:3507S–12S.
104. Gao YT, McLaughlin JK, Blot WJ, Ji BT, Dai Q, Fraumeni JF Jr. Reduced risk of esophageal cancer associated with green tea consumption. *J Natl Cancer Inst* 1994;86:855–8.
105. Aktas O, Prozorovski T, Smorodchenko A, Savaskan NE, Lauster R, Kloetzel PM, et al. Green tea epigallocatechin-3-gallate mediates T cellular NF-kappa B inhibition and exerts neuroprotection in autoimmune encephalomyelitis. *J Immunol* 2004;173:5794–800.
106. Tedeschi E, Menegazzi M, Yao Y, Suzuki H, Forstermann U, Kleinert H. Green tea inhibits human inducible nitric-oxide synthase expression by down-regulating signal transducer and activator of transcription-1alpha activation. *Mol Pharmacol* 2004;65:111–20.
107. Currier NL, Miller SC. The effect of immunization with killed tumor cells, with/without feeding of *Echinacea purpurea* in an erythroleukemic mouse model. *J Altern Complement Med* 2002;8:49–58.
108. Barrett B. Echinacea for upper respiratory tract infection: an evidence-based clinical review. *J Fam Pract* 1999;48:628–35.
109. Binns SE, Purgina B, Bergeron C, Smith ML, Ball L, Baum BR, et al. Light-mediated antifungal activity of Echinacea extracts. *Planta Med* 2000;66:241–4.
110. Thompson KD. Antiviral activity of Viracea against acyclovir susceptible and acyclovir resistant strains of herpes simplex virus. *Antiviral Res* 1998;39:55–61.
111. Larsen MW, Moser C, Hoiby N, Song Z, Kharazmi A. Ginseng modulates the immune response by induction of interleukin-12 production. *APMIS* 2004;112:369–73.
112. Chang YS, Seo EK, Gyllenhaal C, Block KI. *Panax ginseng*: a role in cancer therapy?. *Integr Cancer Ther* 2003;2:13–33.
113. Park EK, Choo MK, Han MJ, Kim DH. Ginsenoside Rh1 possesses antiallergic and anti-inflammatory activities. *Int Arch Allergy Immunol* 2004;133:113–20.
114. Kim TS, Na K, Choi EM, Chung JY, Hwang JK. Immunomodulating activities of polysaccharides isolated from *Panax ginseng*. *J Med Food* 2004;7:1–6.
115. Song Z, Johansen HK, Faber V, Moser C, Kharazmi A, Rygaard J, et al. Ginseng treatment reduces bacterial load and lung pathology in chronic *Pseudomonas aeruginosa* pneumonia in rats. *Antimicrob Agents Chemother* 1997;41:961–4.
116. Crittenden RG, Martinez NR, Playne MJ. Synthesis and utilisation of folate by yoghurt starter cultures and probiotic bacteria. *Int J Food Microbiol* 2003;15:217–22.

Received March 3, 2005; accepted August 26, 2005