Original Article

Bioactive Constituents of Brazilian Red Propolis

Boryana Trusheva¹, Milena Popova¹, Vassya Bankova¹, Svetlana Simova¹, Maria Cristina Marcucci², Patricia Laguna Miorin², Flavia da Rocha Pasin² and Iva Tsvetkova³

¹Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria, ²Faculdade de Farmacia, Universidade Bandeirante de Sao Paulo, Rua Maria Candida, 1813, Vila Guilherme, Campus MC, 02071-013 Sao Paulo, SP Brazil and ³Institute of Microbiology, Bulgarian Academy of Sciences, Acad. G. Bonchev Street, Building 26, 1113 Sofia, Bulgaria

In a new propolis type, red Brazilian propolis, 14 compounds were identified (six of them new for propolis), among them simple phenolics, triterepenoids, isoflavonoids, prenylated benzophenones and a naphthoquinone epoxide (isolated for the first time from a natural source). Three of the major components demonstrated significant antimicrobial activity, and two (obtained as inseparable mixture) possessed radical scavenging activity against 1,1-diphenyl-2-picrylhydrazyl (DPPH).

Keywords: antibacterial activity - chemical constituents - propolis - radical scavenging activity

Introduction

Propolis (bee glue) has a long history of being used as a remedy, dating back to the times of ancient Greece and Rome. Nowadays, it is still used for the treatment of various diseases, and in products like 'health foods', 'biocosmetics', etc., because of its versatile biological activities (1). Tropical propolis samples, and especially Brazilian ones, have shown significant differences in their chemical composition to propolis from temperate zone (2,3). For this reason, Brazilian bee glue has recently become a subject of increasing interest for scientists (4-7). It was found that propolis from different regions of Brazil display different chemical composition, depending on the local flora at the site of collection. Park et al. (8) have specified 12 types of Brazilian propolis according to its geographical origin, chemical composition and source plant. The most popular and well studied Brazilian propolis is the so-called green or Alecrim propolis, which originates from *Baccharis* dracunculifolia (Asteraceae) (9-12). Till now, no chemical data have been published on red propolis from Brazil. In Brazil,

red propolis is collected in the North regions. Red colored propolis is reported to be typical for Cuba, where its plant source was identified as *Clusia nemorosa* (Clusiaceae) (13), and for Venezuela, where bees collect it from *Clusia scrobiculata* (14). In this study, we report our results on antibacterial and antioxidant activity of chemical constituents of red Brazilian propolis.

Materials and Methods

Nuclear magnetic resonance (NMR) spectra were measured on a Bruker AVANCE 250 MNR spectrometer; mass-spectra were measured on a Hewlett Packard 5972 mass spectrometer system.

Propolis. Propolis was collected near Maceio city, Alagoas State, Brazil.

Extraction of propolis. Propolis (61 g) was cut into small pieces and extracted with 70% ethanol (1 : 10, w/v) at room temperature for 24 h. The ethanol extract was concentrated *in vacuo* and extracted successively with petrol ether (40–60°C) three times. The petrol ether extract was evaporated to give 5 g dry residue after evaporation.

Isolation of compounds. The petrol ether extract was subjected to column chromatography on silica gel (300 g) with an n-hexane/acetone gradient (1 : 0.05/1 : 0.4) to produce 20 fractions (**I**–**XX**).

© The Author (2006). Published by Oxford University Press. All rights reserved.

For reprints and all correspondence: Vassya Bankova, Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, 1113 Sofia, Bulgaria. Tel: +359-2-9606149; Fax: +359-2-8700225; E-mail: bankova@orgchm.bas.bg

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that: the original authorship is properly and fully attributed; the Journal and Oxford University Press are attributed as the original place of publication with the correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oxfordjournals.org

Fraction I (300 mg) was rechromatographed on a silica gel column eluted with *n*-hexane/diethyl ether gradient (1: 0.01/1: 1). The first fraction of this column, **I.1** (12 mg), was subjected to gas chromatography-mass spectrometry (GC-MS) analysis with a Hewlett Packard Gas Chromatograph 5890 Series II Plus linked to Hewlett Packard 5972 mass spectrometer system equipped with a 23 m long, 0.25 mm id, 0.5 um film thickness HP5-MS capillary column. The temperature was programmed from 100 to 310°C at a rate of 5°C.min⁻¹. Helium was used as a carrier gas, flow rate 0.7 ml min⁻¹, split ratio 1:80, injector temperature 280°C, ionization voltage 70 eV. Using computer searches on a NIST98 MS data library, the following compounds were identified in the mixture: transanethol 1 (13%), methyl eugenol 2 (14%), trans-methyl isoeugenol 3 (18%), elemicin 4 (26%) and trans-isoelemicin 5 (11%).

The second fraction **I.2** (37 mg), after additional separation by preparative thin layer chromatography (TLC) (silica gel, *n*-hexane/ethyl methyl ketone 1 : 0.06) yielded 20(29)-lupen-3-one **6** (6 mg) and 2,3-epoxy-2-(3-methyl-2-butenyl)-1,4-naphthalenedione **7** (8.6 mg).

Fraction II (300 mg) was rechromatographed on a silica gel column eluted with *n*-hexane/diethyl ether gradient (1 : 0.01/1 : 1). After further purification by preparative TLC (silica gel, *n*-hexane/ethyl methyl ketone 1 : 0.06), a mixture of triterpenic alcohols (36 mg) (¹H-NMR) was obtained. This mixture was analyzed after silylation, using the above mentioned GC-MS apparatus and the same analysis conditions as with fraction I.1. Using computer searches on a NIST98 MS data library, α -amyrin 8, β -amyrin 9 (identity also confirmed by comparison with an authentic sample), cycloartenol 10 and lupeol 11 were identified.

Fraction **VIII** (241 mg) was rechromatographed on a silica gel column, mobile phase *n*-hexane/acetone gradient (1 : 0.05/1 : 0.8). Additional purification by preparative TLC (silica gel, toluene/acetone 1 : 0.2) yielded isosativan **12** (40.8 mg).

Fraction **X** (454 mg) was rechromatographed on a silica gel column with mobile phase *n*-hexane/acetone gradient (1 : 0.05/1 : 1). Further purification by preparative TLC (*n*-hexane/ethyl methyl ketone 1 : 0.1) led to the isolation of 11.8 mg medicarpin **13**.

Fraction **XIII** (620 mg) was rechromatographed on a silica gel column with mobile phase *n*-hexane/acetone gradient (*n*-hexane/acetone 1 : 0.01/1 : 1) to yield 20.5 mg of an inseparable mixture of guttiferone E **14** and xanthochymol **15**.

20(29)-Lupen-3-one **6** was identified based on comparison of its EIMS, ¹H- and ¹³C-NMR spectra and optical rotation with literature data (15).

2,3-Epoxy-2-(3-methyl-2-butenyl)-1,4-naphthalenedione 7, colorless oil, $[\alpha]_D$ 0 (c 0.2, acetone). MS (EI, 70 eV), *m/z* (rel. int. %): 242, M⁺. (28), 227 (M-15)⁺ (85), 213 (M-29)⁺(100), 196 (64), 171 (81), 105 (35%), 89 (36), 69 (30). HRMS (EI) *m/z*: 242.09553 (Calc. for C₁₅H₁₄O₃ : 242.09430). For ¹H-and ¹³C-NMR, see Table 1.

Table 1. NMR data of 7 in CDCl ₃ , δ in ppm (J in Hz)	Hz)
---	-----

С	$\delta^{\mathrm{a}}_{\mathrm{H}}$	$\delta^{\rm b}_{ m C}$	HMBC (H \rightarrow C)
1	_	192.0	
2	_	63.5	
3	3.85 s	59.0	C-2, C-4, C-4a
4	_	191.8	
4a	_	131.9	
5	8.03 m	127.4	C-4, C-4a, C-7
6	7.73–7.76 m	134.5	
7	7.73–7.76 m	134.3	
8	7.96 m	126.8	C-1, C-6, C-8a
8a	_	132.4	
1'	2.69 dd (15.4; 8) 3.05 dd (15.4; 7)	26.1	C-2, C-3, C-2', C-3'
2′	5.09 m	115.4	C-1
3′	_	137.3	
4′	1.73 s	25.8	C-2'. C-3', C-5'
5′	1.69 s	18.0	C-2'. C-3', C-4'

^{a1}H-NMR, 250 MHz.

^{b13}C-NMR, 62.9 MHz.

HMBC, heteronuclear multiple bond correlation.

Isosativan **12**, colorless crystals. UV, EIMS, ¹H- and ¹³C-NMR spectra identical with literature data (16), $[\alpha]_D 0$ (c 0.27, chloroform).

(6aS,11aS)-Medicarpin **13**. UV, EIMS, ¹H- and ¹³C-NMR spectra identical with literature data (17), $\lceil \alpha \rceil_D + 184$ (c 0.51, acetone).

Guttiferone E **14** and xanthochymol **15**. The components of this inseparable mixture were identified by comparison of the spectral data for the same mixture published by Gustafson *et al.* (18) ¹H- and ¹³C-NMR, MS (Fig. 1).

Antimicrobial tests. For the investigation of the antibacterial and antifungal activity, the agar cup method (19) was used with test strains *Staphylococcus aureus* 209 (obtained from the Bulgarian Type Culture Collection, Institute for State Control of Drugs, Sofia), *Escherichia coli* WF+ (obtained from the Collection of ZIMET, Central Institute of Microbiology and Experimental Therapy, Jena, Germany) and *Candida albicans* 562 (obtained from the Bulgarian Type Culture Collection, Institute for State Control of Drugs, Sofia). An inhibitory zone with a diameter <10 mm corresponds to lack of activity (10 mm is the diameter of the cup). The test solution (0.1 ml) containing 0.4 mg of each substance in ethanol was applied to every cup (concentration of the test solution 4 mg ml⁻¹). Control experiments with solvents showed that solvents do not have any activity.

DPPH free radical scavenging activity. DPPH free radical scavenging activity was measured according to the procedure described in the literature (20). The decrease of the absorption at 516 nm of the DPPH solution after addition of the tested solution was measured. An aliquot (2960 μ l) of 0.1 mM ethanolic DPPH solution was mixed with 40 μ l of a 3.6 mM solution of the tested substance. The radical scavenging activity was expressed as percentage decrease with respect to control values. Caffeic acid was used as positive control.

=CH₂

Results and Discussion

The petrol ether fraction of the ethanol extract of the investigated propolis sample was subjected to column chromatography on silica gel and several fractions were produced. After further purification by repeated column chromatography and preparative TLC, two complex mixtures, one inseparable mixture of two isomers and four pure compounds were obtained.

The most unpolar fraction, isolated by repeated column chromatography, was of complex composition and was analyzed by GC-MS. It turned out to be composed of following phenylpropene derivatives: *trans*-anethol **1**, methyl eugenol

2. R = H

4. $R = OCH_3$









ő

ÓН

R

15. R = 3-methyl-3-butenyl

Figure 1. Continued.

2, *trans*-methyl isoeugenol **3**, elemicin **4** and *trans*-isoelemicin **5**. Elemicin was the most abundant. Of these compounds, methyl eugenol, methyl isoeugenol, elemicin and isoelemicin were found for the first time in propolis. The composition of this fraction also explains the very unusual anis-like odor of this Brazilian red propolis sample.

The second complex mixture (see Materials and Methods) was comprised of triterpenic alcohols, which were identified by means of GC-MS as α -amyrin **8**, β -amyrin **9**, cycloartenol **10** and lupeol **11**. The most abundant among them was β -amyrin. Triterpenic alcohols are typical for Brazilian propolis (2).

Table 2. Antimicrobial and antiradical activity of isolated compounds

Sample	Antimicro zo	microbial activity inhibitory zone ± SD, mm ^a		DPPH radical scavenging activity
	S. aureus	E. coli	C. albicans	% inhibition
12	14 ± 0	0	15 ± 1	4.5
13	23 ± 1	14 ± 0	26 ± 0	0.7
14/15	19 ± 1	12 ± 0	0	49
Caffeic acid	-	-	-	85.6

^aMean of three measurements.

One of the pure compounds isolated was also of triterpenic nature: the ketone 20(29)-lupen-3-one **6** [identified by comparison of spectral information with literature data (13)], found for the first time in propolis. This compound has recently been found to possess antibiotic activity against bacteria and fungi, and antioxidant activity similar to that of tocopherol (15).

Compound 7 deserves special attention. Its structure was determined as 2,3-epoxy-2-(3-methyl-2-butenyl)-1,4-naphthalenedione on the basis of its MS, infrared, ¹H- and ¹³C-NMR spectra. This is the first isolation of 7 from a natural source. Till now, it was known only as a synthetic product (21,22). The mass- and ¹H-NMR spectra of our compound were identical with the literature data (no ¹³C-NMR data have been published). Compound 7, obtained synthetically from a natural product, demonstrated antibacterial, antifungal and cytotoxic properties (22).

Two isoflavonoids were isolated and identified: the isoflavan isosativan 12 and the pterocarpan medicarpin 13, based on comparison of their spectral properties with literature data (including absolute stereochemistry of 13, confirmed by optical rotation measurements; 12 was racemic). This is the first report of isoflavonoids in propolis other than Cuban. Compounds 12 and 13 were till now found only in Cuban propolis (16,17). This fact suggests that Cuban red propolis and Brazilian red propolis might have a common plant source, but a plant that produces isoflavonoids as components of its exudates is not yet known. The presence of isoflavonoids suggests some plant of the Leguminosae family but further studies are needed for confirmation. Especially 13 is of particular interest: it is an important plant phytoalexin well known for its antimicrobial and especially antifungal activity (23).

Compounds 14 and 15 are double bond isomers which occur as an inseparable mixture, but the structures were deduced by comparison of the spectral data of the mixture with the values for 14 and 15 from the literature (22–25). ¹H- and ¹³C-NMR spectra of the mixture were virtually identical to those of 14 and 15 and all HMBC correlations were fully consistent with these structures. Moreover, Gustafson *et al.* (18) reported the isolation of the same inseparable mixture from *Clusia rosea* leaves as the active anti-HIV principle. These compounds have been detected as traces in red Cuban propolis (13) originating from *C. rosea* floral resins. In Cuban propolis nemorosone, another polyisoprenylated benzophenone, is the most important constituent (13). In our case, however, the mixture of **14** and **15** is among the major components of the extract. So the main plant source is most probably some other *Clusia* species. Moreover, the presence of isoflavonoids in our sample is an indication that another plant source could be involved, as isoflavonoids have never been found in the resins of Clusiaceae plants.

Three of the isolated compounds were tested for their antibacterial and radical scavenging activity against DPPH radicals. The results are represented in Table 2. The results indicated that the isoflavonoids **12** and **13** are important antimicrobial components of red propolis, especially concerning the activity against *C. albicans*. This is not surprising, taking into consideration that pterocarpans are known for their antifungal activity and play a defensive role in many plants due to this activity (26). The mixture of prenylated benzophenones **14/15** demonstrated good activity against *S. aureus*. The mixture showed also significant radical scavenging activity against DPPH, obviously it is one of the most important antioxidant components of the extract.

The identification of new propolis constituents in red Brazilian propolis, most of them having antibacterial, antimycotic and antiradical activities, is a further confirmation of the fact that propolis, independently of its plant source and chemical composition, always possesses antimicrobial and antioxidant activity. This is due to the role that propolis plays in the hive: it is the 'chemical weapon' of bees against pathogen microorganisms and the elements of weather. However, in different propolis types, different chemical constituents are responsible for the valuable activities (27). The results obtained demonstrate once again that propolis remains a fascinating subject for further studies and application to CAM.

Acknowledgments

The authors wish to thank Mrs Diana Nikolova for running GC-MS analyses.

References

- 1. Burdock GA. Review of the biological properties and toxicity of bee propolis (propolis). *Food Chem Toxicol* 1998;36:347–63.
- Marcucci MC, Bankova VS. Chemical composition, plant origin and biological activity of Brazilian propolis. *Curr Top Phytochem* 1999;2: 115–23.
- Bankova VB, De Castro SL, Marcucci MC. Propolis: recent advances in chemistry and plant origin. *Apidologie* 2000;31:3–15.
- Banskota A, Tezuka Y, Midorikawa K, Matsushige K, Kadota S. Two novel cytotoxic benzofuran derivatives from Brazilian propolis. J Nat Prod 2000;63:1277–9.
- Claus R, Kinscherf Ch, Bonaterra G, Basnet P, Metz J, Deigner H-P. Antiapoptotic effects of propolis extract and propol on human macrophages exposed to minimally modified low density lipoprotein. *Arzneim-Forsch/ Drug Res* 2000;5:373–9.
- Hayacibara MF, Koo H, Rosalen PL, Duarte S, Franco EM, Bowen WH, et al. *In vitro* and *in vivo* effects of isolated fractions of Brazilian propolis on caries development. *J Ethnopharmacol* 2005;101:110–5.
- Shimazawa M, Chikamatsu S, Morimoto N, Mishima S, Nagai H, Hara H. Neuroprotection by Brazilian green propolis against *in vitro* and *in vivo* ischemic neuronal damage. *Evid Based Complement Alternat Med* 2005;2:201–7.

- Park KY, Alencar SM, Aguiar CL. Botanical origin and chemical composition of Brazilian propolis. J Agric Food Chem 2002;50:2502–6.
- Bankova V, Boudourova-Krasteva G, Sforcin JM, Frete X, Kujumgiev A, Maimoni-Rodella R, et al. Phytochemical evidence for the plant origin of Brazilian propolis from Sao Paulo state. Z Naturforsch 1999;54c:401–5.
- 10. Kumazawa S, Yoneda M, Shibata I, Kanaeda J, Hamasaka T, Nakayama T. Direct evidence for the plant origin of Brazilian propolis by the observation of honeybee behavior and phytochemical analysis. *Chem Pharm Bull* 2003;51:740–2.
- Slatino A, Weinstein , Teixeira E, Negri G, Message D. Origin and chemical variation of Brazilian propolis. *Evid Based Complement Alternat Med* 2005;2:33–8.
- Teixeira EW, Negri G, Meira RMSA, Message D, Salatino A. Plant origin of green propolis: bee behavior, plant anatomy and chemistry. *Evid Based Complement Alternat Med* 2005;2:85–92.
- Cuesta-Rubio O, Fontana-Uriba BA, Ramirez-Apan T, Cardenas J. Polyisoprenylated benzophenones in Cuban propolis: biological activity of nemorosone. Z Naturforsch 2002;57c:372–8.
- Trusheva B, Popova M, Naydenski H, Tsvetkova I, Rodriguez JG, Bankova V. New polyisoprenylated benzophenones from Venezuelan propolis. *Fitoterapia* 2004;75:683–9.
- Kim E-M, Jung H-R, Min T-J. Purification, structure determination and biological activities of 20(29)-lupen-3-one from *Daedaleopsis tricolor* (Bull. ex Fr.) Bond. et Sing. *Bull Korean Chem Soc* 2001;22:59–62.
- Questa-Rubio O, Frontana-Uriba B, Perez JC. Isoflavonoides en propoleos Cubanos. *Rev Cubana Farm* 2001;35:(Suppl):58–61.
- Picinelli AL, Campo Fernandez M, Questa Rubio O, Marques Hernandez I, De Simone F, Rastrelli L. Isoflavonoids isolated form Cuban propolis. J Agric Food Chem 2005;53:9010–6.

- Gustafson KR, Blunt JW, Munro MHG, Fuller RW, McKee TC, Cardellina JH, et al. The guttiferones, HIV-inhibitory benzophenones from *Symphomia globuliferea, Garcinia livingstonei, Garcinia ovalifolia* and *Clusia rosea. Tetrahedron* 1992;48:10093–102.
- Spooner FD, Sykes G. Laboratory assessment of antibacterial activity. In: Norris JR, Ribbons DW (eds). *Methods in Microbiology*, vol. 7B. London, New York: Academic Press, 1972, 216–7.
- Nenadis N, Tsimidou M. Observations on the estimation of scavenging activity of phenolic compounds using rapid 1,1-diphenyl-2picrylhydrazyl (DPPH) tests. J Am Oil Chem Soc 2002;79:1191–5.
- Pluim H, Wynberg H. Catalytic asymmetric induction in oxidation reactions. Synthesis of optically active epoxynaphthoquinones. J Org Chem 1980;45:2498.
- Perry NB, Blunt JW, Munro MHG. A cytotoxic and antifungal 1,4naphtoquinone and related compounds from a New Zealand brown alga, *Landsburgia quercifolia. J Nat Prod* 1991;54:978–85.
- Dixon RA, Paiva NL. Stress-induced phenylpropanoid metabolism. *Plant Cell* 1995;7:1085–97.
- Blount JF, Williams TH. Revised structure by xanthochymol. *Tetrahedron* Lett 1976;17:2921–4.
- Rama Rao AV, Venkatswamy G, Yemul SS. Xanthochymol and isoxanthochymol, two novel polyisoprenylated benzophenones from *Garcinia xanthochymus. Indian J Chem* 1980;19b:627–33.
- Reynaud J, Guilet D, Terreux R, Lussignol M, Walchshofer N. Isoflavonoids in non-leguminous families: an update. *Nat Prod Rep* 2005;22: 504–15.
- 27. Bankova V. Recent trends and important developments in propolis research. *Evid Based Complement Alternat Med* 2005;2:29–32.

Received November 10, 2005; accepted February 10, 2006



The Scientific **World Journal**



Gastroenterology Research and Practice





Journal of Diabetes Research



Disease Markers



Immunology Research





Submit your manuscripts at http://www.hindawi.com





BioMed **Research International**



Journal of Ophthalmology

Computational and Mathematical Methods in Medicine





Behavioural Neurology









Research and Treatment





Oxidative Medicine and Cellular Longevity



Stem Cells International

