Original Article

Anti-Inflammatory Potential of Ethanolic Leaf Extract of *Eupatorium adenophorum* Spreng. Through Alteration in Production of TNF- α , ROS and Expression of Certain Genes

Ashim K. Chakravarty,¹ Tamal Mazumder,¹ and Shankar N. Chatterjee²

¹ Immunology and Cell Biology Laboratory, Department of Zoology, School of life Sciences, University of North Bengal, Siliguri 734013, India

² Former Joint Director, SeriBiotech, Ministry of Textiles, Government of India, Kolkata, India

Correspondence should be addressed to Ashim K. Chakravarty, prof.ashim.chakravarty@rediffmail.com

Received 28 September 2009; Accepted 22 March 2010

Copyright © 2011 Ashim K. Chakravarty et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Search for a novel anti-inflammatory agent from a herbal source, such as *Eupatorium adenophorum* Spreng., a plant from the Eastern Himalayas, is of prime interest in the present investigation. Inflammation causes tissue destruction and development of diseases such as asthma, rheumatoid arthritis, and so forth. The ethanolic leaf extract of *E. adenophorum* (EEA) was administered intravenously and in other cases topically at the site of delayed type hypersensitivity (DTH) reaction in mouse foot paw induced with dinitrofluorobenzene. EEA can effectively inhibit DTH reaction and bring back normalcy to the paw much earlier than the controls. Efficacy of EEA on regulatory mechanisms for inflammation has also been considered. Intravenous administration of EEA increased the number of CD4⁺ T cells in spleen and tumor necrosis factor (TNF)- α in serum of DTH mice. Initially it was difficult to reconcile with the anti-inflammatory role of EEA and simultaneous induction of TNF- α , an established pro-inflammatory cytokine. EEA induces higher expression of TNF- α gene and amount of the cytokine in serum. We discussed the other role of TNF- α , its involvement in repairing tissue damage incurred in course of inflammatory reaction. EEA also induces TGF- β encoding a cytokine involved in tissue repair mechanism. EEA inhibits expression of another pro-inflammatory cytokine gene *IL-1 β* and downregulates cyclooxygenase 2 (COX2) gene responsible for metabolism of inflammatory mediators like prostaglandins. Furthermore, anti-inflammatory role of EEA is also revealed through its inhibition of hydroxyl radical generation. Notably EEA does not necessarily affect the expression of other inflammation-related genes such as *IL-6*, *IL-10* and *IKK*. The present study reports and analyzes for the first time the anti-inflammatory property of the leaf extract of *E. adenophorum*.

1. Introduction

Inflammation is the body's way of dealing with infections, maintaining a subtle balance between the beneficial effects of inflammation cascades to restrict the infection and potential for long-term tissue destruction [1–3]. If not controlled, inflammation can lead to development of diseases such as chronic asthma, rheumatoid arthritis, multiple sclerosis, inflammatory bowel disease, and so forth, [4–9]. Till date a very few anti-inflammatory drugs from herbal origin have been found, and a number of plants from ethno-medicinal databases are under laboratory investigation across the world [10].

The plant *Eupatorium adenophorum* Spreng. (Figure 1) belongs to the family Asteraceae (Compositae) [11]. A num-

ber of plants of this family are commonly used in folklore medicine in different parts of the world [12–19]. In Kurseong and Darjeeling hill region of the Eastern Himalayas, local people use leaves of *E. adenophorum* Spreng., growing at an altitude of 800–2050 m, for remedial purposes against oral and skin sores. These observations suggested a probable anti-inflammatory and immunomodulatory activity of the plant's leaf extract. Earlier Mandal et al. [20] reported analgesic property of methanolic extract of the leaves. The present investigation intends to explore the anti-inflammatory property of ethanolic leaf extract of *E. adenophorum* Spreng. (EEA) in delayed type hypersensitivity (DTH) induced by 2,4-dinitrofluorobenzene (DNFB) in mouse model.

DTH reaction is initiated by pre-sensitized CD4⁺ T_{DTH} cells [21] and then other inflammatory cells and cytokines

are involved at the site of reaction. The number of CD4⁺ T cells in course of DTH reaction and treatment with EEA has been enumerated to understand the effect of EEA on these cells. TNF- α is the most important cytokine that plays a major role in all the inflammation reactions. Serum TNF- α of DTH mice has also been investigated in the present study in course of inflammation.

Many reports reveal that reactive oxygen species (ROS) play an important role in developing various pathophysiological conditions including inflammation [22–25] and potent anti-inflammatory agents can scavenge the free radicals to quench the biochemical fire [26–28]. The hydroxyl radical (OH[•]) especially plays a crucial role in developing inflammation. Scavenging of hydroxyl radical (OH[•]) by EEA has been studied.

Besides TNF- α , many other cytokines play a key role in orchestrating immune responses in inflammation [29–36]. Here, expression of some cytokine genes such as TNF- α , IL-1 β , IL-6, IL-10 and TGF- β in splenic T cells of DTH mice has been studied at transcription level with and without intravenous (i.v.) application of the plant extract. Expression of inhibitory kappa kinase (IKK) gene has also been judged. The enzyme degrades I κ B subunit to release active NF- κ B [37, 38], that is, involved in activating inflammatory responses as shown by Tak and Firestein [39] and Yamamoto and Gaynor [40].

The expression of COX1 and COX2 genes encoding two isozymes of cyclooxygenase has been taken into account here. Cyclooxygenase is known to play a significant role in induction of inflammation by producing inflammatory mediators like prostaglandins and leukotrienes from arachidonic acid [41–47].

2. Methods

2.1. Animal. Inbred adult Swiss Albino mice of both sexes, 12–16 weeks old have been used. They are maintained in our animal house with food and water *ad libitum*. Animals of approximately equal age and weight were used for experimental and control groups in the experiment. The experimental protocols used in the study have been approved by the Animal Ethical Committee (Regn No. 840/ac/04/CPCSEA).

2.2. Extract Preparation. Extract was prepared from fresh leaves of the plant *E. adenophorum* using the protocol outlined earlier [48–51]. Leaves were collected from their natural habitat at about 1400 m high slope of the Eastern Himalayas, mainly around Kurseong hill. The scientific identification of the plant has been checked by Professor A. P. Das, Plant Taxonomy Lab., Department of Botany, University of North Bengal. The leaves were cleaned thoroughly with water and allowed to air dry. In total, 10 g of leaves were crushed to a paste with a mortar and pestle. An amount of 10 mL of absolute alcohol (ethanol) was added to the paste and kept in a refrigerator overnight for extraction. The alcoholic extract was then filtered first through Whatmann filter paper and the filtrate was refiltered again through cellulose acetate filter paper (0.2 μ m porosity, Sartorius) for

FIGURE 1: Photograph of *E. adenophorum* Spreng. in its natural habitat.

sterilization and finally stored in airtight sterilized vial at 4°C. The extract was used as such for different experiments.

2.3. Experimental and Control Sets and Statistical Analysis. In all the experiments, the effect of EEA was compared with two sets of control, one with equivalent amount of ethanol present in EEA and the other without any treatment. Each experiment was done on the basis of triplicate readings and such an experiment was repeated thrice or more. Results are expressed as Mean $\pm$ SD of at least nine observations. Statistical significance was analyzed using one-way ANOVA software package.

2.4. Induction of Delayed Type Hypersensitivity Reaction with 2,4-DNFB and Application of EEA. Delayed type hypersensitive (DTH) reaction was induced in mouse foot paw by subcutaneous application of 2,4-DNFB [48, 52]. Primary sensitization was carried out by applying 0.0001% DNFB subcutaneously in the right foot pad. After 8 days, mice were resensitized with 0.000001% DNFB on the left foot pad. Two different volumes of percentage solutions of DNFB, 25 or 50 μ L for both sensitization and resensitization, were used in separate experimental setups. The day of resensitization was considered as “0” day for enumeration of DTH reaction. The size of the left paw before resensitization was considered as normal size for the paw. The degree of inflammatory swelling set in the resensitized left paw was measured by a slide caliper. The effect of EEA on DTH reaction set in by two different doses of DNFB was judged after topical or i.v. application of the extract. For topical application, 5 μ L of EEA was applied on the resensitized paw per day from first day of resensitization. For i.v. administration, 25 μ L of EEA was used 1 h prior to resensitization. The percentage of inhibition of inflammation by EEA in reference to the

ethanol control has been calculated by using the following formula:

Inhibition of DTH (%)

$$= \frac{\text{Untreated DTH paw size} - \text{Experimental paw size}}{\text{Untreated DTH paw size}} \times 100. \quad (1)$$

2.5. Isolation of CD4⁺ T Cells Through Magnetic Assorted Cell Sorter (MACS). The splenic lymphocytes were obtained from untreated DTH mice and mice intravenously injected with EEA and ethanol after 24, 48 and 72 h of resensitization following the protocol of Chakravarty and Maitra [53]. Erythrocytes in the spleen cell suspension were lysed by exposure to Tris-buffered ammonium chloride (0.83%, pH 7.2). For depletion of adherent cells, the suspension was incubated in a plastic petri dish at 37°C in humidified atmosphere for 30 min. Non-adherent lymphocyte population was collected and centrifuged and finally resuspended at a concentration of 10⁷ cells in 80 µL. To the aliquot of 80 µL cell suspension, 20 µL of CD4⁺ (L3TH) microbeads (130-049-201, Miltenyi Biotech, Germany) [54–56] with a magnetic probe was added in the test tube. The tubes were refrigerated at 4–6°C for attachment of the bead to the CD4⁺ cells for 15 min. The mixture of cells and magnetic beads is then poured into the magnetic separation (MS) column fitted in the slot of the magnet of MACS. The unlabeled cells passed through the column and were collected in a tube. The MS column was removed from the separator and placed in a fresh collection tube. Of PBS, 1 ml was pipetted onto the MS column and labeled CD4⁺ cells were flushed out from the column by

firmly pushing the designated plunger into the column. The magnetic labeled CD4⁺ cells were then counted in a hemocytometer.

2.6. Estimation of Serum TNF-α. The quantitation of serum TNF-α of DTH bearing mice treated (i.v.) with EEA and control mice was performed by solid phase sandwich enzyme-linked immunosorbent assay (ELISA) kit (Pharmingen, USA) following the protocol outlined by Paul et al. [57–59].

2.7. Hydroxyl Radical Scavenging Assay. The hydroxyl radical (OH[•]) was generated from Fe²⁺-ascorbate-EDTA-H₂O₂ system (Fentons' reaction). Assay reaction mixture was prepared by mixing a 20 mM phosphate buffer, 2 mM FeCl₃, 1 mM EDTA, 2.8 mM 2-deoxy D-ribose, 1 mM H₂O₂ and 1 mM L-ascorbic acid. OH[•] reacts with the deoxy D-ribose and a series of reaction follows to form malonaldehyde (MDA) [60]. An aliquot of 1 ml of the reaction mixture was added in each tube of experimental, alcohol control and normal control sets and incubated at 37°C for 1 h. Two different doses of EEA, 10 and 25 µL, were tested for scavenging hydroxyl radical. In alcohol and normal control, same volume of ethanol and water was added respectively. After incubation, 2 mL of TBA–TCA reagent was added in each tube and boiled for 15 min for generation of MDA. MDA generated was measured at 552 nm in a spectrophotometer. The effect of both EEA and alcohol on generation of hydroxyl radical has been expressed as percentage of inhibition in MDA generation over normal control sets. The formula used is given below:

$$\text{Percentage of inhibition} = \left[100 - \frac{\text{MDA generated in experimental/alcohol control tubes}}{\text{MDA generated in normal control tubes}} \right] \%. \quad (2)$$

2.8. Gene Expression Analysis. Expression of the genes of interest in this study has been carried out using single-cell RNA phenotyping procedure as outlined by Rappolee et al. [47, 61, 62].

2.9. RNA Isolation. RNA was isolated from splenic T cells of nine mice in each group using RNeasy Mini kit (74104, Qiagen, Valencia, CA, USA), as per manufacturer's protocol. Briefly, 6 × 10⁶ T cells were homogenized with a 300 µL RLT buffer and passing them through a 2 mL syringe fitted with a 27-gauge needle. Of 70% ethanol, 300 µL was added to the homogenate and collected in a spin column fitted upon a collection tube. The spin columns and collection tubes were supplied by the manufacturer. After a brief centrifugation for 15 s at 10 000 rpm, the fluid was passed into the collection tube that was then decanted and reattached to the spin column. With addition of 500 µL of buffer RW1 into the spin column centrifugation was made again for 15 s at 10 000 rpm. Following decantation of collection tube, 500 µL of buffer RPE was added to the spin column and centrifuged similarly,

and the step was repeated one more time. Finally, the spin column was fitted upon a fresh collection tube and washed twice with 15 µL of DEPC-treated water to come up with a total of 30 µL volume containing the RNA sample.

The concentration of RNA was measured spectrophotometrically at 400× dilution with Shimadzu UV-160, Japan. The extracted RNA was used for cDNA synthesis.

2.10. cDNA Synthesis. The isolated RNA was used for First strand cDNA synthesis utilizing the RevertAid First strand cDNA synthesis kit # K1621 from Fermentas and the manufacturer's protocol was followed. For synthesis of first strand cDNA, the primer used for PCR amplification was oligo(dT¹³) synthesized by GMBH. cDNA constructed was stored at –20°C for further use.

2.11. Primer Utilized and Amplification Schedule. Primers were designed from various geneBank accession retrieved from PUBMED Data Bank as listed in Table 1, using the primer program available on the internet. The designed

TABLE 1: Primers.

Primers	Accession number	Sense (5'–3')	Antisense (3'–5')	T_m (°C)
IKK	NM_010546	CCAGACTCCAAGGTGGTGT	TGCAGATCACAGGCAGAAAC	60.0
TNF- α	NM_013693	TGGCACAGCCAAG	GGGACCCCTGCTC	52.36
TGF- β	NM_011577	TTGCTTCAGTCCACAGAGA	TGGTTGTAGAGGGCAAGGAC	59.99
IL-1 β	NM_008361	GTGGCAGCTACCTGTGTCTT	GGAGCCTGTAGTGCAGTTGT	57.96
IL-6	NM_031168	GGGAAATCGTGGA	AGGTTTGCCGAGT	43.9
IL-10	NM_010548	CCAAGCCTTATCGGAAATGA	TTTTCACAGGGGAGAAATCG	60.035
COX1	BC023322	AGAAACTGGTCTGCCTCA	AACCCACATCAAGGACTG	54.02
COX2	NM_011198	AGCACCATTCTCCTTGAA	GTAGGCTGTGGATCTTGC	54.0
PKC- θ	NM_008859	AAGTGAGAAACCCCGCTAT	AGGCAAATCCCTTCCAGTCT	60.01
Perforin	NM_011073	ACCCTGAATGGGCTCACA	GCAGCAGTCTGGTTGGT	57.0

primers were synthesized by GMBH, Germany. Details of the primers are given in Table 1.

PCR was performed using a thermocycler (PeqLab, Germany) for 35 cycles in a 30 μ L reaction mixture containing Taq DNA polymerase buffer, all four dNTPs, oligonucleotide primers, Taq DNA polymerase and cDNA products. After amplification, PCR products were analyzed on a 0.8% (w/v) agarose gel. The band density was quantified on the basis of the known concentration of lambda DNA (30 ng) through ImageAide, Spectronics Corporation, NY.

3. Results

3.1. Effect of EEA on Delayed Type Hypersensitive Reaction

3.1.1. With Topical Application. Figure 2 represents the data of delayed type hypersensitivity reaction induced with 25 μ L DNFB and treatment with ethanolic leaf extract of *E. adenophorum* (EEA) (experimental) and ethanol in control. Swelling of resensitized paw in both control and experimental mice was maximum on third day; with topical application of EEA, the maximum swelling was in the range of 0.4733 ± 0.0227 cm, and in ethanol-treated control mice the maximum value was 0.5667 ± 0.173 cm (Figure 2). The paw size in EEA-treated mice gets back to normal range by the ninth day of resensitization. Inhibition with EEA is a continuous process throughout the experiment. Figure 2 also represents the percentages of inhibition on different days.

A higher dose (50 μ L) of DNFB causes the swelling to be more and the peak on the second day; however, recovery with EEA treatment is on the ninth day (Figure 3). In fact EEA inhibited the swelling by 60% on the ninth day. Figure 4 is the photographic representation of DTH reaction induced with 50 μ L DNFB and inhibition with EEA treatment.

3.1.2. With i.v. Application. When EEA was administered intravenously in DTH bearing mice, inflammatory swelling of the resensitized paw persisted longer than in cases of topical application of EEA. The dose of 25 μ L DNFB could induce maximum swelling of 0.963 ± 0.012 cm on the second day in untreated mice. Intravenous application of EEA

FIGURE 2: Inhibitory effect of topical application of EEA on delayed type hypersensitive (DTH) reaction induced with 25 μ L 2,4-DNFB (significance at $*P < .01$).

caused the DTH reaction to slow down. In alcohol group the swelling was maximum, 0.7367 ± 0.0045 cm, on fifth day (also on sixth). EEA treatment restricted the swelling to 0.5123 ± 0.0112 cm on fifth day and brought back normalcy (0.2967 ± 0.0103 cm) by 11th day; the inhibition over the ethanol control was found to be 45.39% (Figure 5).

The degree of swelling was more at DTH site with higher dose (50 μ L) of DNFB (Figure 6.) inhibited the reaction more effectively than alcohol alone (control) and allowed regaining normalcy by 13 days (Figure 6).

3.2. Estimation of the Number of CD4⁺ T Cells in DTH Mice. CD4⁺ T cells from splenic lymphocyte population were isolated by labeling with microbeads containing a paramagnetic probe and passing through MACS. The number of CD4⁺ T lymphocytes in the spleen of DTH mice treated with EEA

FIGURE 3: Indurations of DTH swelling, induced with 50 μ L DNFB, and inhibition after topical application of EEA and alcohol (significance at $*P < .01$).

FIGURE 4: Photographs showing effect of topical application of EEA and alcohol on resensitized paw of DTH mice, induced by 50 μ L DNFB, during 9-day period of the study.

was significantly more than the controls, about two and a half times at 24 and 48 h and two times at 72 h (Figure 7). A slight increase in the cell number in the alcohol control group was noted (Figure 7).

FIGURE 5: Changes in paw size of DTH mice, induced with 25 μ L DNFB, and after i.v. application of EEA and alcohol (significance of inhibition by EEA at $*P < .01$ and $**P < .05$).

FIGURE 6: Changes in paw size of DTH mice, induced with 50 μ L DNFB, and after i.v. application of EEA and alcohol (significance of inhibition by EEA at $*P < .01$).

3.3. Serum TNF- α of DTH-Bearing Mice and Upon EEA Treatment. Serum TNF- α in three groups of mice bearing DTH reaction—untreated, alcohol and EEA treated (i.v.)—was measured by a solid phase sandwich enzyme-linked immunosorbent assay (ELISA).

There was no appreciable difference in the level of serum TNF- α in the three groups of mice at 24 h (Figure 8). The TNF- α level in DTH mice treated with EEA (i.v.) was more than that in other groups by 48 h (93.23 pg mL^{-1}) and the level was maintained upto 72 h (Figure 8).

3.4. Generation of Hydroxyl Radical and Effect of EEA. The hydroxyl radical ($\text{OH}^{\bullet}$) is potentially harmful for the cellular

FIGURE 7: Effect of intravenously administered EEA and alcohol on count of CD4⁺ T cells from spleen of mice after 24, 48 and 72 h of induction of DTH with 25 µL DNFB (significance of results with EEA over control at * $P < .01$).

FIGURE 8: Level of serum TNF- α from DTH mice (induced with 25 µL DNFB) at three different hours in presence and absence of EEA (significance compared to controls at * $P < .01$).

macromolecules and is implicated in pathophysiology of inflammation. EEA inhibits hydroxyl radical generation up to 57.98% at a dose of 25 µL. The same dose of ethanol (control) showed 23.62% inhibition. At a lower dose of 10 µL, the degree of inhibition was about 49.25% in experimental ones and 14.88% in case of alcohol control (Figure 9).

3.5. Expression of Certain Genes in Splenic T Cells of DTH Mice with and without EEA Treatment. The T cells actively

FIGURE 9: The effect of two different doses of EEA on inhibition of generation of hydroxyl radical (OH[•]).

participate in the progression of DTH reaction. Up- or downregulation of certain genes in the cells must be related to the production of pro- and anti-inflammatory cytokines, transcription factors and mediators. The level of expression of some of these genes at transcription level in splenic T cells of DTH mice untreated and treated with EEA was judged, by quantitating the cDNA PCR product amplified with specific primers. The quantitation was done against 30 ng of lambda DNA as a standard using ImageAide, Spectronics Corporation, NY. The data are presented in Figures 10 and 11.

EEA caused an increment in the expression of TNF- α , a pro-inflammatory cytokine (Figures 10 and 11). At the same time, EEA inhibited expression of pro-inflammatory cytokine IL-1 β and showed no effect on IL-6, another pro-inflammatory cytokine.

EEA did not influence the expression of IL-10, an anti-inflammatory cytokine beyond alcohol control. The expression of TGF- β encoding a cytokine involved in regeneration was induced by EEA beyond the level in the controls.

EEA apparently did not influence expression of IKK and COX1 genes but downregulated the expression of COX2 gene.

4. Discussion

The ethanolic leaf extract of *E. adenophorum* Spreng. (EEA) could effectively suppress the inflammatory reaction set in foot paw by injecting two different doses of 2,4-DNFB (Figure 4). The topical application of EEA was more effective in inhibition of the swelling of foot paw and gaining normalcy faster than its i.v. application (Figures 2, 3, 5, and 6).

EEA treatment caused significant increase in the number of the CD4⁺ T cell population in DTH mice (Figure 7). These cells are known to play a central role in inflammatory reactions by secreting all different kinds of cytokines that regulate participation of other kinds of cells [21, 63, 64].

FIGURE 10: Expression of certain inflammation-related genes in T cells obtained from DTH mice, and with and without EEA treatment, containing nine mice in each group. Agarose gel electrophoresis of cDNA PCR products amplified with different gene-specific primers.

FIGURE 11: Graphical presentation of quantification of the DNA bands presented in Figure 10 showing expression of the inflammation-related genes in T cells of DTH mice with and without EEA treatment, containing nine mice in each group. The DNA bands were quantified against 30 ng of lambda DNA through ImageAide, Spectronics Corporation, NY.

TNF-α is a major cytokine involved in DTH reaction. EEA induced higher level of serum TNF-α, surpassing the level in mice undergoing DTH reaction or in mice treated with alcohol (Figure 8). It is worthwhile to note EEA capable

FIGURE 12: Effect of EEA on inhibition of DTH reaction and repair mechanism presented schematically.

of inhibiting DTH reaction is inducing a pro-inflammatory cytokine like TNF-α. Banno et al. [65] showed that TNF-α promotes tissue repair of damage skin by inducing basement membrane components and collagen degrading proteases to participate actively in reconstruction of extracellular matrix. Kuwano and his coworkers [66] found that TNF-α can also induce growth-promoting event like angiogenesis by increasing mRNA level of IL-8, vascular endothelial growth factor and fibroblast growth factors in endothelial cells. We think of similar participation of TNF-α in tissue repair and regeneration to bring back normalcy aftermath of DTH reaction. Thus, TNF-α plays a double role in DTH reaction—pro-inflammatory cum restoring agent.

The ability of EEA for induction of TNF-α gene was studied next. Indeed, it induced higher level of expression of the gene (Figures 10 and 11). EEA does not necessarily affect the expression of gene for other pro-inflammatory cytokines such as IL-1β and IL-6 in a similar fashion. EEA inhibited IL-1β expression and did not influence the expression of IL-6 gene (Figures 10 and 11). It seems that IL-1β without any known function in repair mechanism manifests more inhibitory effect of EEA on inflammation. Stimmeder and his co-workers [67] observed that lornoxicam and other non-steroidal anti-inflammatory drugs inhibit IL-1β expression as well as inflammation. Kohli et al. [68] reported that curcumin, the active component in the rhizome of *Curcuma longa* Linn., demonstrates its anti-inflammatory activity by inhibiting production of IL-1β in lung inflammatory cells.

An anti-inflammatory agent does not necessarily always regulate all the anti-inflammatory cytokine genes as we find that in the present study EEA does not influence the anti-inflammatory cytokine gene IL-10 (Figures 10 and 11).

EEA upregulates expression of *TGF- β* (Figures 10 and 11). *TGF- β* performs as a growth factor in all different kinds of events of collagen production and extracellular matrix reorganization as shown by Barcellos-Hoff [69]. This cytokine might function here to restore normalcy along with *TNF- α* in repair mode as discussed earlier. Simultaneous upregulation of these two genes have also been observed by Chao et al. [70] in microglial cell culture. Sullivan et al. [71] also reported similar trend in expression of these two cytokines in interstitial pulmonary fibrosis-affected lung fibroblasts.

Tak and Firestein [39] and Yamamoto and Gaynor [40] elucidated involvement of NF- κ B pathway for induction of inflammation. Activation of NF- κ B is mediated by the action of Inhibitory kappa kinase (IKK) degrading inhibitory I κ B subunit. Thus, measuring the expression of *IKK*, one can derive the involvement of NF- κ B pathway in a reaction. EEA could not induce the expression of *IKK* gene beyond the controls (Figures 10 and 11) indicating non-involvement of NF- κ B activation pathway for DTH reaction induced with DNFB.

COX1 and COX2 gene products are two isoforms of the cyclooxygenase enzyme that metabolizes arachidonic acid into the inflammatory mediators like prostaglandins and leukotrienes [41–47]. Notably EEA only influences expression of *COX2* gene by way of inhibition (Figures 10 and 11). This may be another way of execution of anti-inflammatory activity by EEA. Salvioli et al. [72] reviewed curcumin, a potent anti-inflammatory agent, also inhibits *COX2* in abetting inflammation.

It is known that reactive oxygen species play an important role in development of inflammation [22–25]. In the present study, the ability of EEA in quenching the generation of hydroxyl radical has been tested and found to be effective (Figure 9).

In summary, ethanolic leaf extract of *E. adenophorum* (EEA) exerts anti-inflammatory activity (Figure 12), likely through inhibition of *IL-1 β* , *COX2* genes and quenching ROS like hydroxyl radical. Simultaneously EEA induces production of *TNF- α* , a pro-inflammatory cytokine. This paradox can only be resolved in the light of participation of *TNF- α* in tissue repair in the aftermath of inflammation. Interestingly, the expression of *TGF- β* gene encoding the cytokine responsible for growth and repair mechanism is also inducible by EEA. *E. adenophorum* as a source of anti-inflammatory substance seems worthy to report. The active compound from the extract is yet to be identified. Isolation of that compound will allow understanding molecular mechanism of the activity of the substance. So far, Zhang et al. [73] reported presence of a few flavonones and sesquiterpene lactones in *E. adenophorum*.

The present investigation also intends that any herbal agent having anti-inflammatory property might be screened faster by estimating its ability to induce or inhibit the genes encoding substances participating in inflammation.

Funding

University Grants Commission (UGC), India.

Acknowledgments

Authors are thankful to the University Grants Commission (UGC), India, for providing financial assistance. They also express their heartfelt gratitude to Dr Arnab Sen, Department of Botany, NBU, and Dr Ranadhir Chakraborty, Department of Biotechnology, NBU, to let us avail instrumental facilities in their laboratories. They are also thankful to Dr Bholanath Paul, IITR, Lucknow, India for providing valuable chemicals to carry out ELISA in his laboratory. They would like to extend their sincere thanks to Professor A. P. Das, Department of Botany, NBU, for identification of the plant.

References

- [1] D. L. Simmons, "What makes a good anti-inflammatory drug target?" *Drug Discovery Today*, vol. 11, no. 5-6, pp. 210–219, 2006.
- [2] K. Saukkonen, S. Sande, C. Cioffe et al., "The role of cytokines in the generation of inflammation and tissue damage in experimental gram-positive meningitis," *Journal of Experimental Medicine*, vol. 171, no. 2, pp. 439–448, 1990.
- [3] N. L. Parenteau and J. Hardin-Young, "The biological mechanisms behind injury and inflammation: how they can affect treatment strategy, product performance, and healing," *Wounds*, vol. 19, no. 4, pp. 87–96, 2007.
- [4] L. M. Fabbri, M. Saetta, G. Picotti, and C. E. Mapp, "Late asthmatic reactions, airway inflammation and chronic asthma in toluene-diisocyanate-sensitized subjects," *Respiration*, vol. 58, no. 1, pp. 18–21, 1991.
- [5] R. J. Stevens, K. M. J. Douglas, A. N. Saratzis, and G. D. Kitas, "Inflammation and atherosclerosis in rheumatoid arthritis," *Expert Reviews in Molecular Medicine*, vol. 7, no. 7, pp. 1–24, 2005.
- [6] L. Klareskog, L. Padyukov, J. Rönnelid, and L. Alfredsson, "Genes, environment and immunity in the development of rheumatoid arthritis," *Current Opinion in Immunology*, vol. 18, no. 6, pp. 650–655, 2006.
- [7] M. Simka, "Blood brain barrier compromise with endothelial inflammation may lead to autoimmune loss of myelin during multiple sclerosis," *Current Neurovascular Research*, vol. 6, no. 2, pp. 132–139, 2009.
- [8] R. D. Zipser, "Mediators of inflammation in inflammatory bowel disease," *Digestive Diseases and Sciences*, vol. 33, no. 3, pp. 4S–5S, 1988.
- [9] J. Drews and S. Ryser, "Classic drug targets," *Nature Biotechnology*, vol. 15, pp. 1318–1319, 1997.
- [10] D.-X. Kong, X.-J. Li, and H.-Y. Zhang, "Where is the hope for drug discovery? Let history tell the future," *Drug Discovery Today*, vol. 14, no. 3-4, pp. 115–119, 2009.
- [11] B. A. AuLD and P. M. Martins, "The autecology of *Eupatorium adenophorum* Spreng. In Austraha," *Weed Research*, vol. 15, pp. 27–31, 1975.
- [12] L. Muschietti, M. Derita, V. Sülsen et al., "In vitro antifungal assay of traditional Argentine medicinal plants," *Journal of Ethnopharmacology*, vol. 102, no. 2, pp. 233–238, 2005.
- [13] E. Ahmed, M. Arshad, M. Ahmad, M. Saeed, and M. Ishaque, "Ethnopharmacological survey of some medicinally important planks of Galliyat areas of NWFP, Pakistan," *Asian Journal of Plant Sciences*, vol. 3, pp. 410–415, 2004.

- [14] P. Castillo-España, A. Cisneros-Estrada, M. L. Garduño-Ramírez, O. Hernández-Abreu, R. Ramírez, and S. Estrada-Soto, "Preliminary ethnopharmacological survey of plants used in Mexico for the treatment of hypertension," *Pharmacognosy Reviews*, vol. 3, pp. 41–65, 2009.
- [15] A. Pieroni, M. E. Giusti, C. de Pasquale et al., "Circum-Mediterranean cultural heritage and medicinal plant uses in traditional animal healthcare: a field survey in eight selected areas within the RUBIA project," *Journal of Ethnobiology and Ethnomedicine*, vol. 2, pp. 16–27, 2006.
- [16] M. Heinrich, M. Robles, J. E. West, B. R. Ortiz De Montellano, and E. Rodríguez, "Ethnopharmacology of Mexican asteraceae (compositae)," *Annual Review of Pharmacology and Toxicology*, vol. 38, pp. 539–565, 1998.
- [17] N. R. Monks, A. Ferraz, S. A. Bradgnon, K. R. Machado, M. F. Richter, and A. B. DaRocha, "In vitro cytotoxicity of extracts from Brazilian Asteraceae," *Pharmaceutical Biology*, vol. 40, pp. 494–500, 2002.
- [18] E. K. Akkol, U. Koca, I. Pesin, and D. Yilmazer, "Evaluation of the wound healing potential of *Achillea biebersteinii* Afan. (Asteraceae) by in vivo excision and incision models," *Evidence-Based Complementary and Alternative Medicine*. In press.
- [19] S. J. Uddin, I. D. Grice, and E. Tiralongo, "Cytotoxic effects of Bangladeshi plant extracts," *Evidence-Based Complementary and Alternative Medicine*. In press.
- [20] S. K. Mandal, R. Boominathan, B. Parimaladevi, S. Dewanjee, and S. C. Mandal, "Analgesic activity of methanol extract of *Eupatorium adenophorum* Spreng. leaves," *Indian Journal of Experimental Biology*, vol. 43, no. 7, pp. 662–663, 2005.
- [21] G. K. Matsushima and S. A. Stohlman, "Maturation of the delayed-type hypersensitivity response in SJL mice: absence of effector cell induction," *European Journal of Immunology*, vol. 18, no. 9, pp. 1411–1416, 1988.
- [22] U. Bandyopadhyay, D. Das, and R. K. Banerjee, "Reactive oxygen species: oxidative damage and pathogenesis," *Current Science*, vol. 77, no. 5, pp. 658–666, 1999.
- [23] P. C. Dedon and S. R. Tannenbaum, "Reactive nitrogen species in the chemical biology of inflammation," *Archives of Biochemistry and Biophysics*, vol. 423, no. 1, pp. 12–22, 2004.
- [24] B. Halliwell, J. R. Houlst, and D. R. Blake, "Oxidants, inflammation, and anti-inflammatory drugs," *FASEB Journal*, vol. 2, no. 13, pp. 2867–2873, 1988.
- [25] P. J. Barnes, "Reactive oxygen species and airway inflammation," *Free Radical Biology and Medicine*, vol. 9, no. 3, pp. 235–243, 1990.
- [26] E. L. Nguemfol, T. Dimol, A. B. Dongmo et al., "Anti-oxidative and anti-inflammatory activities of some isolated constituents from the stem bark of *Allanblackia monticola* Staner L.C (Guttiferae)," *Inflammopharmacology*, vol. 17, pp. 37–41, 2009.
- [27] P. Abreu, S. Matthew, T. González, D. Costa, M. A. Segundo, and E. Fernandes, "Anti-inflammatory and antioxidant activity of a medicinal tincture from *Pedilanthus tithymaloides*," *Life Sciences*, vol. 78, no. 14, pp. 1578–1585, 2006.
- [28] S. Joseph, B. Sabulal, V. George, T. P. Smina, and K. K. Janardhanan, "Antioxidative and antiinflammatory activities of the chloroform extract of *Ganoderma lucidum* found in South India," *Scientia Pharmaceutica*, vol. 77, no. 1, pp. 111–121, 2009.
- [29] C. A. Dinarello, "Proinflammatory cytokines," *Chest*, vol. 118, pp. 503–508, 2000.
- [30] R. Strieter, S. Kunkel, and R. Bone, "Role of tumor necrosis factor-alpha in disease states and inflammation," *Critical Care Medicine*, vol. 21, pp. S447–S463, 1993.
- [31] N. W. Lukacs, R. M. Strieter, S. W. Chensue, M. Widmer, and S. L. Kunkel, "TNF- α mediates recruitment of neutrophils and eosinophils during airway inflammation," *Journal of Immunology*, vol. 154, no. 10, pp. 5411–5417, 1995.
- [32] K. J. Tracey and A. Cerami, "Tumor necrosis factor: a pleiotropic cytokine and therapeutic target," *Annual Review of Medicine*, vol. 45, pp. 491–503, 1994.
- [33] A. Schweizer, U. Feige, A. Fontana, K. Muller, and C. A. Dinarello, "Interleukin-1 enhances pain reflexes: mediation through increased prostaglandin E2 levels," *Agents Actions*, vol. 25, pp. 246–251, 1998.
- [34] C. Gabay, "Interleukin-6 and chronic inflammation," *Arthritis Research & Therapy*, vol. 8, supplement 2, pp. S3–S8, 2006.
- [35] S. Omoigui, "The interleukin-6 inflammation pathway from cholesterol to aging—role of statins, bisphosphonates and plant polyphenols in aging and age-related diseases," *Immunity and Ageing*, vol. 4, pp. 1–22, 2007.
- [36] M. Jutel, M. Akdis, F. Budak et al., "IL-10 and TGF- β cooperate in the regulatory T cell response to mucosal allergens in normal immunity and specific immunotherapy," *European Journal of Immunology*, vol. 33, no. 5, pp. 1205–1214, 2003.
- [37] I. Alkalay, A. Yaron, A. Hatzubai et al., "Stimulation-dependent I kappa B alpha phosphorylation marks the NF-kappa B inhibitor for degradation via the ubiquitin-proteasome pathway," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 92, pp. 10599–10603, 1995.
- [38] V. Pande and M. J. Ramos, "NF- κ B in human disease: current inhibitors and prospects for *de novo* structure based design of inhibitors," *Current Medicinal Chemistry*, vol. 12, no. 3, pp. 357–374, 2005.
- [39] P. P. Tak and G. S. Firestein, "NF-kappaB: a key role in inflammatory diseases," *Journal of Clinical Investigation*, vol. 107, pp. 7–11, 2001.
- [40] Y. Yamamoto and R. B. Gaynor, "Therapeutic potential of inhibition of the NF-kappaB pathway in the treatment of inflammation and cancer," *Journal of Clinical Investigation*, vol. 107, pp. 135–142, 2001.
- [41] M. E. Turini and R. N. DuBois, "Cyclooxygenase-2: a therapeutic target," *Annual Review of Medicine*, vol. 53, pp. 35–57, 2002.
- [42] C. J. Smith, Y. Zhang, C. M. Koboldt et al., "Pharmacological analysis of cyclooxygenase 1 in inflammation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, pp. 13313–13318, 1998.
- [43] W. L. Smith, R. M. Garavito, and D. L. DeWitt, "Prostaglandin endoperoxide H synthase (cyclooxygenases)-1 and -2," *The Journal of Biological Chemistry*, vol. 271, pp. 33157–33160, 1996.
- [44] J. A. Mitchell, S. Larkin, and T. J. Williams, "Cyclooxygenase 2: regulation and relevance in inflammation," *Biochemical Pharmacology*, vol. 50, pp. 1535–1542, 1995.
- [45] J. R. Vane, Y. S. Bakhle, and R. M. Botting, "Cyclooxygenases 1 and 2," *Annual Review of Pharmacology and Toxicology*, vol. 38, pp. 97–120, 1998.
- [46] F. Chen, S. Sun, D. C. Kuhn et al., "Involvement of NF- κ B in silica-induced cyclooxygenase II gene expression in rat alveolar macrophages," *American Journal of Physiology*, vol. 272, no. 4, pp. L779–L786, 1997.
- [47] Y. Devaux, C. Seguin, S. Grosjean et al., "Lipopolysaccharide-induced increase of prostaglandin E(2) is mediated by

- inducible nitric oxide synthase activation of the constitutive cyclooxygenase and induction of membrane-associated prostaglandin E synthase," *Journal of Immunology*, vol. 167, pp. 3962–3971, 2001.
- [48] A. K. Chakravarty, S. N. Chatterjee, H. Yasmin, and T. Mazumder, "Comparison of efficacy of turmeric and commercial curcumin in immunological functions and gene regulation," *International Journal of Pharmacology*, vol. 5, pp. 333–345, 2009.
- [49] A. K. Chakravarty and H. Yasmin, "Activation of cell mediated immune response and apoptosis towards malignant cells with turmeric treatment in murine model," *Indian Journal of Biochemistry and Biophysics*, vol. 45, no. 1, pp. 23–29, 2008.
- [50] A. K. Chakravarty and H. Yasmin, "Alcoholic turmeric extract simultaneously activating murine lymphocytes and inducing apoptosis of Ehrlich ascitic carcinoma cells," *International Immunopharmacology*, vol. 5, no. 10, pp. 1574–1581, 2005.
- [51] A. K. Chakravarty, H. Yasmin, and S. K. Das, "Two way efficacy of alcoholic turmeric extract: stimulatory for murine lymphocytes and inhibitory for ascetic fibrosarcoma," *Pharmaceutical Biology*, vol. 42, pp. 217–224, 2004.
- [52] M. Iijima and S. I. Katz, "Specific immunologic tolerance to dinitrofluorobenzene following topical application of dinitrothiocyanobenzene: modulation by suppressor T cells," *Journal of Investigative Dermatology*, vol. 81, no. 4, pp. 325–330, 1983.
- [53] A. K. Chakravarty and U. K. Maitra, "Polyclonally stimulated murine T cells in control fibrosarcoma *in situ*," *Journal of the Indian Institute of Science*, vol. 70, pp. 503–516, 1990.
- [54] R. Busch, D. Cesar, D. Higuera-Alhino, T. Gee, M. K. Hellerstein, and J. M. McCune, "Isolation of peripheral blood CD4⁺ T cells using RosetteSep^{TM} and MACS^{TM} for studies of DNA turnover by deuterium labeling," *Journal of Immunological Methods*, vol. 286, pp. 97–109, 2004.
- [55] M. P. Matheu and M. D. Cahalan, "Isolation of CD4⁺ T cells from mouse lymph nodes using Miltenyi MACS purification," *Journal of Visualized Experiments*, no. 9, p. 409, 2007.
- [56] L. A. Stanciu and R. Djukanović, "Isolation of T-cell subsets by magnetic cell sorting (MACS)," *Methods in Molecular Biology*, vol. 134, pp. 133–141, 2000.
- [57] V. Sureshkumar, B. Paul, M. Uthirappan et al., "Pro-inflammatory and anti-inflammatory cytokine balance in gasoline exhaust induced pulmonary injury in mice," *Inhalation Toxicology*, vol. 17, pp. 161–168, 2005.
- [58] I. V. M. L. R. Sirish Kumar, B. N. Paul, R. Asthama, A. K. Saxena, S. Mehrotra, and G. Rajan, "Swertia Chirayita mediated modulation of interleukin-1 β , interleukin-6, interleukin-10, interferon- γ and tumor necrosis factor- α in arthritic mice," *Immunopharmacology and Immunotoxicology*, vol. 25, pp. 573–583, 2003.
- [59] P. D. Drew and J. A. Chavis, "Female sex steroids: effects upon microglial cell activation," *Journal of Neuroimmunology*, vol. 111, no. 1-2, pp. 77–85, 2000.
- [60] B. Halliwell, J. M. C. Gutteridge, and O. I. Aruoma, "The deoxyribose method: a simple 'test-tube' assay for determination of rate constants for reactions of hydroxyl radicals," *Analytical Biochemistry*, vol. 165, no. 1, pp. 215–219, 1987.
- [61] D. A. Rappolee, D. Mark, M. J. Banda, and Z. Werb, "Wound macrophages express TGF- α and other growth factors *in vivo*: analysis by mRNA phenotyping," *Science*, vol. 241, pp. 708–712, 1988.
- [62] D. A. Rappolee, C. A. Brenner, R. Schultz, D. Mark, and Z. Werb, "Developmental expression of PDGF, TGF- α , and TGF- β genes in preimplantation mouse embryos," *Science*, vol. 241, no. 4874, pp. 1823–1825, 1988.
- [63] S. Ehlers, M. E. A. Mielke, and H. Hahn, "CD4⁺ T cell associated cytokine gene expression during experimental infection with *Listeria monocytogenes*: the mRNA phenotype of granuloma formation," *International Immunology*, vol. 6, no. 11, pp. 1727–1737, 1994.
- [64] L. W. Poulter, G. J. Seymour, O. Duke, G. Janossy, and G. Panayi, "Immunohistological analysis of delayed-type hypersensitivity in man," *Cell Immunology*, vol. 74, pp. 358–369, 1982.
- [65] T. Banno, A. Gazel, and M. Blumenberg, "Effects of tumor necrosis factor- α (TNF α) in epidermal keratinocytes revealed using global transcriptional profiling," *Journal of Biological Chemistry*, vol. 279, no. 31, pp. 32633–32642, 2004.
- [66] S. Yoshida, M. Ono, T. Shono et al., "Involvement of interleukin-8, vascular endothelial growth factor, and basic fibroblast growth factor in tumor necrosis factor alpha-dependent angiogenesis," *Molecular and Cellular Biology*, vol. 17, no. 7, pp. 4015–4023, 1997.
- [67] J. Berg, H. Fellier, T. Christoph, J. Grarup, and D. Stimmeder, "The analgesic NSAID lornoxicam inhibits cyclooxygenase (COX)-1/-2, inducible nitric oxide synthase (iNOS), and the formation of interleukin (IL)-6 *in vitro*," *Inflammation Research*, vol. 48, no. 7, pp. 369–385, 1999.
- [68] K. Kohli, J. Ali, M. J. Ansari, and Z. Raheman, "Curcumin: a natural antiinflammatory agent," *Indian Journal of Pharmacology*, vol. 37, no. 3, pp. 141–147, 2005.
- [69] M. H. Barcellos-Hoff, "Radiation-induced transforming growth factor β and subsequent extracellular matrix reorganization in murine mammary gland," *Cancer Research*, vol. 53, no. 17, pp. 3880–3886, 1993.
- [70] C. C. Chao, S. Hu, W. S. Sheng, M. Tsang, and P. K. Peterson, "Tumor necrosis factor- α mediates the release of bioactive transforming growth factor- β in murine microglial cell cultures," *Clinical Immunology and Immunopathology*, vol. 77, pp. 358–365, 1995.
- [71] D. E. Sullivan, M. Ferris, D. Pociask, and A. R. Brody, "Tumor necrosis factor- α induces transforming growth factor- β expression in lung fibroblasts through the extracellular signal-regulated kinase pathway," *American Journal of Respiratory Cell and Molecular Biology*, vol. 32, pp. 342–349, 2005.
- [72] S. Salvioli, E. Sikora, E. L. Cooper, and C. Franceschi, "Curcumin in cell death processes: a challenge for CAM of age-related pathologies," *Evidence-Based Complementary and Alternative Medicine*, vol. 4, no. 2, pp. 181–190, 2007.
- [73] M.-L. Zhang, M. Wu, J.-J. Zhang, D. Irwin, Y.-C. Gu, and Q.-W. Shi, "Chemical constituents of plants from the genus *Eupatorium*," *Chemistry and Biodiversity*, vol. 5, no. 1, pp. 40–55, 2008.