Original Article

Diaphragmatic Breathing Reduces Exercise-Induced Oxidative Stress

Daniele Martarelli, Mario Cocchioni, Stefania Scuri, and Pierluigi Pompei

Department of Experimental Medicine and Public Health, University of Camerino, Via Madonna delle Carceri, 62032 Camerino, Macerata, Italy

Correspondence should be addressed to Daniele Martarelli, daniele.martarelli@unicam.it

Received 31 March 2009; Accepted 2 October 2009

Copyright © 2011 Daniele Martarelli et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Diaphragmatic breathing is relaxing and therapeutic, reduces stress, and is a fundamental procedure of Pranayama Yoga, Zen, transcendental meditation and other meditation practices. Analysis of oxidative stress levels in people who meditate indicated that meditation correlates with lower oxidative stress levels, lower cortisol levels and higher melatonin levels. It is known that cortisol inhibits enzymes responsible for the antioxidant activity of cells and that melatonin is a strong antioxidant; therefore, in this study, we investigated the effects of diaphragmatic breathing on exercise-induced oxidative stress and the putative role of cortisol and melatonin hormones in this stress pathway. We monitored 16 athletes during an exhaustive training session. After the exercise, athletes were divided in two equivalent groups of eight subjects. Subjects of the studied group spent 1 h relaxing performing diaphragmatic breathing and concentrating on their breath in a quiet place. The other eight subjects, representing the control group, spent the same time sitting in an equivalent quiet place. Results demonstrate that relaxation induced by diaphragmatic breathing increases the antioxidant defense status in athletes after exhaustive exercise. These effects correlate with the concomitant decrease in cortisol and the increase in melatonin. The consequence is a lower level of oxidative stress, which suggests that an appropriate diaphragmatic breathing could protect athletes from long-term adverse effects of free radicals.

1. Introduction

Stress is defined as a physiological reaction to undesired emotional or physical situations. Initially, stress induces an acute response (fight or flight) that is mediated by catecholamines. When stress becomes chronic and lasts for a long time, the stressed organism reacts with physiological alterations to adapt to the unfavorable conditions. This ACTH-mediated reaction affects the immune and neuroendocrine systems, and it is responsible for several diseases [1]. Numerous data support the hypothesis that the pathophysiology of chronic stress can be due, at least partially, to an increase in oxidative stress [2–4], which may also contribute to heart disease [5, 6], rheumatoid arthritis [7, 8], hypertension [9, 10], Alzheimer's disease [11, 12], Parkinson's disease [13], atherosclerosis [14] and, finally, aging [15].

Some authors have attributed stress-induced oxidative stress to an increase in glucocorticoids. In fact, there is evidence to suggest that glucocorticoids induce oxidative stress mainly by altering the expression and activity of

antioxidant enzymes, thus impairing the antioxidant defense of the body [16–19]. High levels of glucocorticoids are known to decrease blood reduced glutathione (GSH) and erythrocyte superoxide dismutase (SOD) activity in rats [20]. Other enzymes are also involved, and NADPH oxidase, xanthine oxidase and uncoupled endothelial nitric oxide synthase are important sources of reactive oxygen species (ROS) in glucocorticoid-induced oxidative stress (see [9] for a review on this argument).

A number of studies support the fact that meditation, through the modulation of the neuroendocrine response, combats stress and its related diseases. In fact, beyond its psychological and social effects, clinical studies have documented that meditation improves the immune system [21] and decreases cardiovascular risk factors such as hyperlipidemia, hypertension and atherosclerosis [22–30]. A reduction in both glucocorticoids and oxidative stress has been documented in people who practice meditation regularly. Hormonal reactions to stressors, in particular plasma cortisol levels, are lower in people who meditate than

in people who do not [31–36], suggesting that it is possible to modulate the neuroendocrine system through neurological pathways. Analysis of oxidative stress levels in people who meditate indicated that transcendental meditation, Zen meditation and Yoga correlate with lower oxidative stress levels [37–43].

Melatonin could also be involved in the reduction of oxidative stress because increased levels of this hormone have been reported after meditation [44–46]. This neurohormone is considered a strong antioxidant and is used as a treatment for aging. Melatonin in fact, increases several intracellular enzymatic antioxidant enzymes, such as SOD and glutathione peroxidase (GSH-Px) [47, 48], and induces the activity of γ -glutamylcysteine synthetase, thereby stimulating the production of the intracellular antioxidant GSH [49]. A number of studies have shown that melatonin is significantly better than the classic antioxidants in resisting free-radical-based molecular destruction. In these *in vivo* studies, melatonin was more effective than vitamin E, β -carotene [50–52] and vitamin C [53–55]. In addition to mental stress, physical stress also increases the production of ROS. In exhaustive and prolonged exercise, ROS production is elevated, and changes in exercise intensity (aerobic-anaerobic) have been associated with a higher degree of oxidative stress [56–59]. Although it has been established that a continuous and moderate physical activity reduces stress, intense and prolonged exercise is deleterious and needs a proper recovery procedure. The link between physical and psychological stress is apparent because they have equivalent hormonal responses. Actually, both types of stress are characterized by activation of the neuroendocrine axis, which leads to the production of ACTH and cortisol. The beneficial or detrimental role of cortisol in athletes has been debated, as some believe that its catabolic actions are detrimental to muscle recovery, whereas others believe that its anti-inflammatory actions are beneficial to muscle recovery. Plasma cortisol levels increase in response to intense and prolonged exercise [60, 61]. Ponjee et al. [62] demonstrated that cortisol increased significantly in male athletes after they ran a marathon. In another study, plasma ACTH and cortisol were found elevated in highly trained runners and in sedentary subjects after intense treadmill exercise [63].

Additionally, melatonin levels are affected by physical activity. There are some conflicting reports regarding the effects of exercise on melatonin levels, with some studies reporting an increase, some a decrease and some reporting no change in melatonin concentrations after exercise [64–70]. However, these contradictory results could be due to light conditions and the timing or intensity of exercise. Moreover, sex, age and training of the monitored athletes may contribute to the different results reported in these studies. It has been speculated that intense sport increases melatonin secretion due to the necessity of combating the free radical production that occurs during exercise, and melatonin could be responsible for amenorrhea in female athletes as an effect of overtraining [71].

Most, if not all, meditation procedures involve diaphragmatic breathing (DB), which is the act of breathing deeply into the lungs by flexing the diaphragm rather than the rib

cage. DB is relaxing and therapeutic, reduces stress and is a fundamental procedure of Pranayama Yoga, Zen, transcendental meditation and other meditation practices.

Although exercise-induced ROS production can be produced via different pathways [56], we speculated that by combating the exercise-induced increase in cortisol levels and by stimulating melatonin levels, DB could improve antioxidant defenses and, therefore, decrease oxidative stress. We have recently demonstrated that in master athletes, oxidative stress induced by intense exercise reaches dangerous levels [72]. Therefore, in this study, we investigated the effects of DB on exercise-induced oxidative stress and the putative role of cortisol and melatonin hormones in this stress pathway.

2. Methods

2.1. Subjects and Exercise. Athletes were monitored during a training session for a 24-h long contest. This type of race lasts for 24 h, generally starting at 10:00 am and ending at 10:00 am the following day. Bikers ride as many kilometers as possible on a specific circuit trail in the 24-h period. Athletes are allowed to stop, to sleep, to rest and to eat as much food as they want to eat.

The session analyzed in this study was a reproduction of the first 8 h of the race, which is generally the most intense. Athletes started to ride at 10:00 am and stopped at 6:00 pm. They consumed the same food and rested the same time.

Since the parameters measured can differ for each individual, we performed preliminary analyses to select subjects with comparable cortisol, melatonin, antioxidant and oxidative stress values.

We selected 16 amateur male cyclists, aged 44.4 ± 2 years ($\pm$ SD). Their mean height and weight were 175.4 ± 7.5 cm and 68.8 ± 5.7 kg, respectively, (Table 1). Subjects were informed of the purpose of the study, and all of them gave their informed consent prior to their inclusion. This study has been performed in accordance with the ethical standards laid down in the 1964 declaration of Helsinki.

None of the subjects had taken medication or supplements within the past 30 days that might alter the study outcome, and none of them had a history of medical or surgical events that could affect the study outcome, including cardiovascular disease or metabolic, renal, hepatic or musculoskeletal disorders.

2.2. Experimental Procedure. After exercising, athletes took a shower and drank water to rehydrate. They were then divided in two equivalent groups of eight subjects (Table 1). Subjects of the studied group were previously trained to relax by performing DB and concentrating on their breath. These athletes spent 1 h (6:30–7:30 pm) relaxing performing DB in a quiet place. The other eight subjects, representing the control group, spent the same time sitting in an equivalent quiet place. The only activity allowed was reading magazines. Lighting levels were monitored throughout the experiment and did not exceed 15 lux, a level well below that known to influence melatonin secretion [73, 74].

TABLE 1: Characteristics of the sample studied.

Athletes	DB group				Control group				All athletes			
	Age (years)	Height (cm)	Weight (kg)	Kilometers covered	Age (years)	Height (cm)	Weight (kg)	Kilometers covered	Age (years)	Height (cm)	Weight (kg)	Kilometers covered
Mean	44.50	175.25	68.63	130.13	44.38	175.50	69.13	128.63	44.44	175.38	68.88	129.38
SD	2.32	6.94	4.52	5.01	2.12	8.34	6.31	5.60	2.16	7.52	5.66	5.19

After the resting and DB periods, athletes consumed the same food and retired for sleeping at 10:00 pm. At 11:00 pm, all of them were sleeping.

We referred to the DB applied here as a relaxation technique. Instead of training the athletes with some form of meditation, we preferred DB because it is easy to learn and to perform and because it does not require any moral conviction that could generate psychologically adverse reactions. However, DB associated with a focused mind (in this case, awareness on the breath as specified in the methods section) can be considered a form of meditation such as focused meditation or others [75].

2.3. Oxidative Stress Determination. Oxidative stress was measured by performing the d reactive oxygen metabolites (d-ROMs) test [76, 77], which determines the plasma ROMs produced by ROS. The d-ROMs test is based on the concept that plasmahydroperoxides react with the transition metal ions liberated from the proteins in the acidic medium and are converted to alkoxy and peroxy radicals. These newly formed radicals are able to oxidize *N,N*-diethyl-*p*-phenyldiamine to the corresponding radical cation, and its concentration can be determined through spectrophotometric procedures (absorption at 505 nm). The d-ROMs test is expressed in U CARR (Carratelli units), where 1 U CARR = 0.08 mg H₂O₂ dl⁻¹. Values higher than 300 U CARR indicate oxidative stress. ROMs were determined before starting the exercise (9:30 am), at the end of the exercise (6:00 pm), immediately after the DB periods (7:30 pm), at 2:00 am, and 24 h after the exercise (10:00 am of the following day).

2.4. Biological Antioxidant Potential Determination. The antioxidant defense status was assessed by determining the biological antioxidant potential (BAP test), which depends on the plasma levels of antioxidants. The BAP test is based on the ability of a coloured solution, containing a source of ferric (Fe³⁺) ions adequately bound to a special chromogenic substrate, to lose colour when Fe³⁺ ions are reduced to ferrous ions (Fe²⁺), which occurs when a reducing/antioxidant system is added. The ferric chloride reagent (50 µL) is transferred into a cuvette containing the thiocyanate derivative reagent. The resulting colored solution is gently mixed by inversion and its absorbance is measured at 550 nm. Then, 10 µL of plasma is added to the same cuvette, the solution is gently mixed, incubated in a thermostatic block for 5 min at 37°C, and its absorbance at 550 nm is remeasured [78, 79]. The BAP test results are expressed in µmol Fe²⁺/liter of sample. Values higher

than 2200 µmolFe²⁺/liter are considered a normal BAP. d-ROMs and BAP tests were performed using apposite kits and dedicated instrumentation Free Radical Analytical System 4 (FRAS4, Health & Diagnostics Limited Co., Parma, Italy). Since the BAP test must be performed at least 3 h after food was last consumed, the BAP was determined before breakfast at 8:00 am, during the night at 2:00 am, and 24 h post-exercise (8:00 am).

2.5. Saliva Collection. The subjects abstained from alcoholic and caffeinated beverages from the beginning of the training session and were only allowed to drink water. Subjects washed their mouths with distilled water before salivary samples were obtained using the Bühlmann saliva collection device (Bühlmann Laboratories AG, Switzerland). Immediately after collection, the saliva samples were frozen and stored at -80°C until they were assayed for cortisol and melatonin concentrations.

2.6. Cortisol Assay. Salivary cortisol was determined before the exercise began (10:00 am), at the end of the exercise (6:00 pm), immediately after the DB period (7:30 pm), at 2:00 am, and 24 h after the exercise (10:00 am of the next day) using a commercially available EIA kit (Cortisol Express, Cayman Chemical Ann Arbor, MI, USA). Absorbance values were determined at 415 nm using a plate reader. Samples were assayed in triplicate.

2.7. Melatonin Assay. Salivary nocturnal melatonin was determined at 2:00 am using the Bühlmann Direct Saliva Melatonin Elisa (Bühlmann Laboratories AG, Switzerland). This assay is based on a melatonin biotin conjugate antibody, streptavidin conjugated to horseradish peroxidase and a tetramethyl benzidine (TMB) substrate. The product of the substrate was measured spectrophotometrically at 450 nm. The assay sensitivity range was 1–60.6 pg ml⁻¹.

2.8. Statistical Analysis. The characteristics of the studied sample and the effects of DB were analyzed by two-way ANOVA with repeated measurements. A two-sided *t*-test (*post-hoc* comparisons) and the non parametric Wilcoxon-Mann-Whitney test were used for the comparison of numerical data across groups for each time point. A *P*-value < .05 was considered statistically significant. Statistics were compiled using Microsoft Excel and Winstat software. Changes in melatonin levels were analyzed by the two-sided *t*-test and the non-parametric Wilcoxon-Mann-Whitney test.

FIGURE 1: ROMs levels were determined at different times, before and after exercise. Athletes were divided in two equivalent groups of eight subjects. Subjects of the studied group spent 1 h (6:30–7:30 pm) relaxing performing DB and concentrating on their breath in a quiet place. The other eight subjects, representing the control group, spent the same time sitting in an equivalent quite place. Values shown are mean $\pm$ SD. ** $P < .01$ DB versus control group.

3. Results

3.1. Characteristics of the Studied Sample. Subjects were divided into two similar groups, as shown in Table 1. There were no statistical differences for age, height, weight, or km covered between the groups [$F(1,62) = 0.023$; $P > .5$].

3.2. Oxidative Stress Changes. As expected, the exercise induced a strong oxidative stress in athletes (Figure 1).

The ROMs levels were significantly increased after exercise compared to pre-exercise levels. All athletes had an elevation in ROMs in response to the training exercise, reaching particularly high levels of oxidative stress. The overall ANOVA revealed a significant DB effect [$F(1,78) = 11.184$; $P < .01$] and time effect [$F(4,75) = 130.481$; $P < .01$]. After completing the training exercise, there was a significant amount of variability between the ROMs levels of individual athletes, suggesting that each athlete has an individual response to oxidative stress. However, *post-hoc* comparisons confirmed that the mean level of ROMs in athletes of the DB group was significantly lower than the control-group athletes both at 2:00 am ($P < .01$ DB versus control group) and 24 h post-exercise ($P < .01$ DB versus control group). For the DB group, the increase in ROMs levels post-exercise compared to pre-exercise levels was 161.7% at 6:00 pm, 150.9% at 7:30 pm, 141.6% at 2:00 am and 126.8% 24 h post-exercise. For the control group, the increase in ROMs levels post-exercise compared to pre-exercise levels was 160.9% at 6:00 pm, 157.1% at 7:30 pm, 159.9% at 2:00 am and 154% 24 h post-exercise.

3.3. Biological Antioxidant Potential Changes. Figure 2 shows the BAP which significantly increased in both groups.

FIGURE 2: BAP levels were determined at different times, before and after exercise. Athletes were divided in two equivalent groups of eight subjects. Subjects of the studied group spent 1 h relaxing performing DB and concentrating on their breath in a quiet place. The other eight subjects, representing the control group, spent the same time sitting in an equivalent quite place. Since this test must be performed several hours after food ingestion, BAP levels were determined pre-exercise at 8:00 am before breakfast, at 2:00 am, and at 8:00 am 24 h post-exercise. Values shown are mean $\pm$ SD. * $P < .05$ DB versus control group. ** $P < .01$ DB versus control group.

Again, a significant variation among the subjects was observed, but athletes of the DB group presented BAP levels significantly higher than the control group [$F(1,46) = 21.001$; $P < .01$]. This difference was more evident at 2:00 am ($P < .01$ DB versus control group, *post-hoc* comparisons) than 24 h post-exercise ($P < .05$ DB versus control group, *post-hoc* comparisons), where BAP began to return to basal levels. With respect to the pre-exercise values, for the DB group, the increase in BAP levels was 129.1% at 2:00 am and 111.1% at 24 h post-exercise.

For the control group, the increase was 114.2% at 2:00 am and 106.2% at 24 h post-exercise with respect to the pre-exercise values. ANOVA also revealed a significant time effect [$F(2,45) = 91.587$; $P < .01$].

3.4. Changes in Cortisol Levels. ANOVA revealed a significant DB effect [$F(1,78) = 4.028$; $P < .05$]. As shown in Figure 3, significant differences between the groups were observed only at 7:30 pm, after the DB ($P < .05$ DB versus control group, *post-hoc* comparisons). At 2:00 am and 24 h post-exercise, cortisol levels were lower in athletes of the DB group, but differences were not statistically significant. In athletes of the DB group, the decrease in cortisol levels (07:30 p.m.) temporarily precedes the decrease in ROMs levels (2:00 am). It was not possible to determine the effects of exercise on cortisol levels, as hormone concentrations were determined at different times during its circadian rhythm. With respect to the pre-exercise values, for the DB group, cortisol values were 82.2% by 06:00 pm, 61.1% by 7:30 pm,

FIGURE 3: Salivary cortisol levels were determined at different times, before and after exercise. Athletes were divided in two equivalent groups of eight subjects. Subjects of the studied group spent 1 h (6:30–7:30 pm) relaxing performing DB and concentrating on their breath in a quiet place. The other eight subjects, representing the control group, spent the same time sitting in an equivalent quiet place. Values shown are mean $\pm$ SD. * $P < .05$ DB versus control group.

47.7% by 02:00 am and 74.7% 24 h post-exercise. For the control group, with respect to the pre-exercise values, values were 83.4% by 06:00 pm, 79.1% by 7:30 pm, 54.6% by 02:00 am and 86.9% 24 h post-exercise respect to the pre-exercise values. ANOVA also revealed a significant time effect [$F(4,75) = 17.459$; $P < .01$].

3.5. Changes in Melatonin Levels. Figure 4 shows the differences in nocturnal melatonin levels between the two groups of athletes. Melatonin levels were significantly higher in athletes of the DB group ($P < .05$ DB versus control group). These data are congruent with the lower ROMs levels, with the higher BAP levels and with the lower cortisol levels at 7:30 pm.

4. Discussion

This study demonstrates that DB reduces the oxidative stress induced by exhaustive exercise. To our knowledge, this is the first study which explores the effect of DB on the stress caused by exhaustive physical activity.

It is known that cortisol inhibits enzymes responsible for the antioxidant activity of cells and that melatonin is a strong antioxidant. After the training exercise, athletes who underwent DB presented higher levels of BAP, which are congruent with the reduced levels of cortisol and ROMs and with the increased levels of nocturnal melatonin. As in our previous study [72], after exercise, we found an increase in BAP levels in both of the groups analyzed. However, the elevated levels of plasma antioxidant markers after exercise can be explained considering three processes: (i) the suspension of exercise decreases oxidant production, so antioxidant

FIGURE 4: Salivary nocturnal melatonin levels variation after exercise. Values shown are mean $\pm$ SD. * $P < .05$ DB versus control group.

defense can return to normal levels; (ii) up-regulation of antioxidants and (iii) the mobilization of antioxidants from tissues to blood [80]. Beyond these mechanisms, these results also suggest that cortisol and melatonin levels could affect the modulation of antioxidant defenses and are relevant in determining the final level of oxidative stress. The decrease of ROS concentrations in subjects performing DB could be attributed to the reduced neuroendocrine response induced by relaxation.

The rationale is as follows (Figure 5):

- (i) intense exercise increases cortisol production;
- (ii) a high plasmatic level of cortisol decreases body antioxidant defenses;
- (iii) a high plasmatic level of cortisol correlates with a high level of oxidative stress;
- (iv) DB reduces the production of cortisol;
- (v) DB increases melatonin levels;
- (vi) melatonin is a strong antioxidant;
- (vii) DB increases the BAP and
- (viii) DB reduces oxidative stress.

If these results are confirmed in other intense physical activity programs, relaxation could be considered an effective practice to significantly contrast the free radical-mediated oxidative damage induced by intense exercise. Therefore, similar to the way that antioxidant supplementation has been integrated into athletic training programs, DB or other meditation techniques should be integrated into many sports as a method to improve performance and to accelerate recovery. However, wider health implications can be accounted for the use of DB, as it can find applications in several pathologies. For example, the oxidative stress that occurs in the hyperventilation syndrome can be cured by learning

FIGURE 5: Modulation of oxidative stress by exercise and DB.

DB. Hyperventilation, in fact, induces hyperoxia which is known to be related with oxidative stress [81, 82]. The hyperventilation syndrome affects 15% of the population and occurs when breathing rates elevate to 21–23 bpm as a result of constricted non-DB. DB can treat hyperoxia and its consequences acting by two synergic ways: restoring the normal breath rhythm and reducing oxidative stress mainly through the increase in melatonin production which is known for its ability to reduce oxidative stress induced by exposure to hyperbaric hyperoxia [83]. Moreover, Orme-Johnson observed greatly reduced pathology levels in regular meditation practitioners [84, 85]. A 5 years statistic of approximately 2000 regular participants demonstrated that Transcendental Meditation reduced benign and malignant tumors, heart disease, infectious diseases, mental disorders and diseases of the nervous system. Mourya et al. evidenced that slow-breathing exercises may influence autonomic functions reducing blood pressure in patients with essential hypertension [86]. Finally, there are also evidences that procedures which involve the control of the breathing can positively affect type 2 Diabetes [87], depression, pain [88], high glucose level and high cholesterol [89].

Our results contribute to explain these effects as oxidative stress may also play a role in the development of these pathologies [2–15]. The role of melatonin must also be emphasized. Beyond its antioxidant properties, melatonin is involved in the regulation of the circadian sleep-wake rhythm and in the modulation of hormones and the immune system. Due to its wide medical implications, the increase in melatonin levels induced by DB suggests that this breath procedure deserves to be included in public health improvement programs.

In this work, we explored the acute effects of DB, but these outcomes should also be investigated for longer periods, for which we would expect a more intense and beneficial response. For example, it is likely that expert practitioners who frequently utilize of DB could obtain a more significant reduction in oxidative stress and, perhaps, an improvement

in exercise performance. Moreover, relaxation could also be improved by adding another relaxation method to the formula, for example music. In fact, Khalfa et al. [90] demonstrated that relaxing music facilitates recovery from a psychologically stressful task, decreasing the salivary cortisol.

Our results must also be discussed in light of the fact that cortisol has an ACTH-dependent circadian rhythm with peak levels in the early morning and a nadir at night. Athletes start to ride at 10:00 am and stop at 6:00 pm. The DB session started at 6:30 pm and stopped at 7:30 pm. It is probable that these results would be different if the time of physical activity and the DB session were changed. The same is true for melatonin. In fact, significant differences have been reported in melatonin secretion when exercises were performed at different times and under different light conditions [64–70]. We collected the saliva at 2:00 am, when a peak in melatonin must be expected. DB increased the levels of melatonin in athletes, and this correlates with lower oxidative stress (ROMs), with lower cortisol levels and with the higher antioxidant status (BAP) in these athletes.

The mechanism by which relaxation might induce an increase in melatonin levels is uncertain, and whether the melatonin increase is simply due to the cortisol decrease remains to be elucidated. Different mechanisms could be involved. Tooley et al. [46] speculated that meditation-reduced hepatic blood flow [91] could raise the plasma levels of melatonin. Alternatively, since meditation increases plasma levels of noradrenaline [92] and urine levels of the metabolite 5HIAA [93], a possible direct action on the pineal gland could be hypothesized, as melatonin is synthesized in the pineal by serotonin under a noradrenaline stimulus [94]. More likely, we suspect that the increase in melatonin levels determined in our experiment can be mainly attributed to the reduced cortisol levels. Actually, a relationship between cortisol and melatonin rhythms has been observed [95], indicating that melatonin onset typically occurs during low cortisol secretion. In addition, Monteleone et al. [96] found that exercise-induced increases in plasma cortisol preceded

the lower night-time melatonin secretion, thus suggesting a connection between the metabolisms of these two hormones.

More studies are needed to clarify the link between cortisol and melatonin; however, due to the complexity of the pathways involved in maintaining homeostasis and in initiating the stress response, it is plausible that the relationship between the two hormones could be mediated by several mechanisms.

Overall, these data demonstrate that relaxation induced by DB increases the antioxidant defense status in athletes after exhaustive exercise. These effects correlate with the concomitant decrease in cortisol, which is known to negatively affect antioxidant defenses, and the increase in melatonin, a strong antioxidant. The consequence is a lower level of oxidative stress, which suggests that an appropriate recovery could protect athletes from long-term adverse effects of free radicals.

Funding

Department of Experimental Medicine and Public Health, University of Camerino, Macerata, Italy.

References

- [1] S. Cohen, D. Janicki-Deverts, and G. E. Miller, "Psychological stress and disease," *Journal of the American Medical Association*, vol. 298, no. 14, pp. 1685–1687, 2007.
- [2] T. Itoh, T. Saito, M. Fujimura, S. Watanabe, and K. Saito, "Restraint stress-induced changes in endogenous zinc release from the rat hippocampus," *Brain Research*, vol. 618, no. 2, pp. 318–322, 1993.
- [3] F. Scarpellini, M. Sbracia, and L. Scarpellini, "Psychological stress and lipoperoxidation in miscarriage," *Annals of the New York Academy of Sciences*, vol. 709, pp. 210–213, 1994.
- [4] S. Adachi, K. Kawamura, and K. Takemoto, "Oxidative damage of nuclear DNA in liver of rats exposed to psychological stress," *Cancer Research*, vol. 53, no. 18, pp. 4153–4155, 1993.
- [5] B. Molavi and J. L. Mehta, "Oxidative stress in cardiovascular disease: molecular basis of its deleterious effects, its detection, and therapeutic considerations," *Current Opinion in Cardiology*, vol. 19, no. 5, pp. 488–493, 2004.
- [6] N. S. Dhalla, R. M. Temsah, and T. Netticadan, "Role of oxidative stress in cardiovascular diseases," *Journal of Hypertension*, vol. 18, no. 6, pp. 655–673, 2000.
- [7] C. A. Hitchon and H. S. El-Gabalawy, "Oxidation in rheumatoid arthritis," *Arthritis Research and Therapy*, vol. 6, no. 6, pp. 265–278, 2004.
- [8] L. I. Filippin, R. Vercelino, N. P. Marroni, and R. M. Xavier, "Redox signalling and the inflammatory response in rheumatoid arthritis," *Clinical and Experimental Immunology*, vol. 152, no. 3, pp. 415–422, 2008.
- [9] S. L. H. Ong, Y. Zhang, and J. A. Whitworth, "Reactive oxygen species and glucocorticoid-induced hypertension," *Clinical and Experimental Pharmacology and Physiology*, vol. 35, no. 4, pp. 477–482, 2008.
- [10] K. V. Kumar and U. N. Das, "Are free radicals involved in the pathobiology of human essential hypertension?" *Free Radical Research Communications*, vol. 19, no. 1, pp. 59–66, 1993.
- [11] Y. Christen, "Oxidative stress and Alzheimer disease," *American Journal of Clinical Nutrition*, vol. 71, no. 2, pp. 621S–629S, 2000.
- [12] W. R. Markesbery, "Oxidative stress hypothesis in Alzheimer's disease," *Free Radical Biology and Medicine*, vol. 23, no. 1, pp. 134–147, 1997.
- [13] P. Jenner and C. W. Olanow, "Oxidative stress and the pathogenesis of Parkinson's disease," *Neurology*, vol. 47, no. 6, pp. S161–S170, 1996.
- [14] F. Bonomini, S. Tengattini, A. Fabiano, R. Bianchi, and R. Rezzani, "Atherosclerosis and oxidative stress," *Histology and Histopathology*, vol. 23, no. 1–3, pp. 381–390, 2008.
- [15] K. C. Kregel and H. J. Zhang, "An integrated view of oxidative stress in aging: basic mechanisms, functional effects, and pathological considerations," *American Journal of Physiology*, vol. 292, no. 1, pp. R18–R36, 2007.
- [16] L. J. McIntosh and R. M. Sapolsky, "Glucocorticoids increase the accumulation of reactive oxygen species and enhance adriamycin-induced toxicity in neuronal culture," *Experimental Neurology*, vol. 141, no. 2, pp. 201–206, 1996.
- [17] Y. Z. Eid, A. Ohtsuka, and K. Hayashi, "Tea polyphenols reduce glucocorticoid-induced growth inhibition and oxidative stress in broiler chickens," *British Poultry Science*, vol. 44, no. 1, pp. 127–132, 2003.
- [18] A. Ohtsuka, H. Kojima, T. Ohtani, and K. Hayashi, "Vitamin E reduces glucocorticoid-induced oxidative stress in rat skeletal muscle," *Journal of Nutritional Science and Vitaminology*, vol. 44, pp. 779–786, 1998.
- [19] E. S. Epel, E. H. Blackburn, J. Lin et al., "Accelerated telomere shortening in response to life stress," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 49, pp. 17312–17315, 2004.
- [20] A. Orzechowski, P. Ostaszewski, A. Brodnicka et al., "Excess of glucocorticoids impairs whole-body antioxidant status in young rats. Relation to the effect of dexamethasone in soleus muscle and spleen," *Hormone and Metabolic Research*, vol. 32, no. 5, pp. 174–180, 2000.
- [21] R. J. Davidson, J. Kabat-Zinn, J. Schumacher et al., "Alterations in brain and immune function produced by mindfulness meditation," *Psychosomatic Medicine*, vol. 65, no. 4, pp. 564–570, 2003.
- [22] V. A. Barnes, F. A. Treiber, J. R. Turner, H. Davis, and W. B. Strong, "Acute effects of transcendental meditation on hemodynamic functioning in middle-aged adults," *Psychosomatic Medicine*, vol. 61, no. 4, pp. 525–531, 1999.
- [23] V. A. Barnes, F. A. Treiber, and H. Davis, "Impact of Transcendental Meditation on cardiovascular function at rest and during acute stress in adolescents with high normal blood pressure," *Journal of Psychosomatic Research*, vol. 51, pp. 597–605, 2001.
- [24] S. R. Wenneberg, R. H. Schneider, K. G. Walton, C. R. Maclean, D. K. Levitsky, and J. W. Salerno, "A controlled study of the effects of the Transcendental Meditation program on cardiovascular reactivity and ambulatory blood pressure," *International Journal of Neuroscience*, vol. 89, pp. 15–28, 1997.
- [25] R. Calderon Jr., R. H. Schneider, C. N. Alexander, H. F. Myers, S. I. Nidich, and C. Haney, "Stress, stress reduction and hypercholesterolemia in African Americans: a review," *Ethnicity and Disease*, vol. 9, no. 3, pp. 451–462, 1999.
- [26] M. J. Cooper and M. M. Aygen, "A relaxation technique in the management of hypercholesterolemia," *Journal of Human Stress*, vol. 5, no. 4, pp. 24–27, 1979.
- [27] A. Castillo-Richmond, R. H. Schneider, C. N. Alexander et al., "Effects of stress reduction on carotid atherosclerosis

- in hypertensive African Americans," *Stroke*, vol. 31, no. 3, pp. 568–573, 2000.
- [28] M. S. King, T. Carr, and C. D'Cruz, "Transcendental meditation, hypertension and heart disease," *Australian Family Physician*, vol. 31, no. 2, pp. 164–168, 2002.
 - [29] P. Lehrer, Y. Sasaki, and Y. Saito, "Zazen and cardiac variability," *Psychosomatic Medicine*, vol. 61, no. 6, pp. 812–821, 1999.
 - [30] K. Takeo, H. Minamisawa, K. Kanda, and S. Hasegawa, "Heart rates during the daily activity of bZenQ priests," *Journal of Human Ergology*, vol. 13, pp. 83–87, 1984.
 - [31] C. N. Alexander, P. Robinson, D. W. Orme-Johnson, R. H. Schneider, and K. G. Walton, "Effects of Transcendental Meditation compared to other methods of relaxation and meditation in reducing risk factors, morbidity and mortality," *Homeostasis*, vol. 35, pp. 243–264, 1994.
 - [32] R. Sudsuang, V. Chentanez, and K. Veluvan, "Effect of Buddhist meditation on serum cortisol and total protein levels, blood pressure, pulse rate, lung volume and reaction time," *Physiology and Behavior*, vol. 50, no. 3, pp. 543–548, 1991.
 - [33] R. Jevning, A. F. Wilson, and W. R. Smith, "The transcendental meditation technique, adrenocortical activity, and implications for stress," *Experientia*, vol. 34, no. 5, pp. 618–619, 1978.
 - [34] R. Jevning, A. F. Wilson, and J. M. Davidson, "Adrenocortical activity during meditation," *Hormones and Behavior*, vol. 10, no. 1, pp. 54–60, 1978.
 - [35] C. R. MacLean, K. G. Walton, S. R. Wenneberg, D. K. Levitsky, J. P. Mandarino, and R. Waziri, "Effects of the Transcendental Meditation program on adaptive mechanisms: changes in hormone levels and responses to stress after 4 months of practice," *Psychoneuroendocrinology*, vol. 22, pp. 277–295, 1997.
 - [36] R. R. Michaels, J. Parra, D. S. McCann, and A. J. Vander, "Renin, cortisol, and aldosterone during transcendental meditation," *Psychosomatic Medicine*, vol. 41, no. 1, pp. 50–54, 1979.
 - [37] D. H. Kim, Y. S. Moon, H. S. Kim, J. S. Jung, H. M. Park, and H. W. Suh, "Effect of Zen Meditation on serum nitric oxide activity and lipid peroxidation," *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, vol. 29, pp. 327–331, 2005.
 - [38] R. H. Schneider, S. I. Nidich, J. W. Salerno et al., "Lower lipid peroxide levels in practitioners of the transcendental meditation program," *Psychosomatic Medicine*, vol. 60, no. 1, pp. 38–41, 1998.
 - [39] S. Sinha, S. N. Singh, Y. P. Monga, and U. S. Ray, "Improvement of glutathione and total antioxidant status with yoga," *Journal of Alternative and Complementary Medicine*, vol. 13, no. 10, pp. 1085–1090, 2007.
 - [40] R. K. Yadav, R. B. Ray, R. Vempati, and R. L. Bijlani, "Effect of a comprehensive yoga-based lifestyle modification program on lipid peroxidation," *Indian Journal of Physiology and Pharmacology*, vol. 49, no. 3, pp. 358–362, 2005.
 - [41] E. P. Van Wijk, H. Koch, S. Bosman, and R. Van Wijk, "Anatomical characterization of human ultraweak photon emission in practitioners of transcendental meditation and control subjects," *Journal of Alternative and Complementary Medicine*, vol. 12, pp. 31–38, 1998.
 - [42] E. P. A. Van Wijk, R. Lüdtkke, and R. Van Wijk, "Differential effects of relaxation techniques on ultraweak photon emission," *Journal of Alternative and Complementary Medicine*, vol. 14, no. 3, pp. 241–250, 2008.
 - [43] T. Kita, M. Yokode, N. Kume et al., "The concentration of serum lipids in Zen monks and control males in Japan," *Japanese Circulation Journal*, vol. 52, no. 2, pp. 99–104, 1988.
 - [44] K. Harinath, A. S. Malhotra, K. Pal, R. Prasad, R. Kumar, and T. C. Kain, "Effects of hatha yoga and omkar meditation on cardiorespiratory performance, psychologic profile, and melatonin secretion," *The Journal of Alternative and Complementary Medicine*, vol. 10, pp. 261–268, 2004.
 - [45] E. E. Solberg, A. Holen, Ø. Ekeberg, B. Østerud, R. Halvorsen, and L. Sandvik, "The effects of long meditation on plasma melatonin and blood serotonin," *Medical Science Monitor*, vol. 10, no. 3, pp. CR96–CR101, 2004.
 - [46] G. A. Tooley, S. M. Armstrong, T. R. Norman, and A. Sali, "Acute increases in night-time plasma melatonin levels following a period of meditation," *Biological Psychology*, vol. 53, no. 1, pp. 69–78, 2000.
 - [47] C. Rodriguez, J. C. Mayo, R. M. Sainz et al., "Regulation of antioxidant enzymes: a significant role for melatonin," *Journal of Pineal Research*, vol. 36, no. 1, pp. 1–9, 2004.
 - [48] R. J. Reiter, D.-X. Tan, and M. D. Maldonado, "Melatonin as an antioxidant: physiology versus pharmacology," *Journal of Pineal Research*, vol. 39, no. 2, pp. 215–216, 2005.
 - [49] K. Winiarska, T. Fraczyk, D. Malinska, J. Drozak, and J. Bryla, "Melatonin attenuates diabetes-induced oxidative stress in rabbits," *Journal of Pineal Research*, vol. 40, no. 2, pp. 168–176, 2006.
 - [50] G. Baydas, H. Canatan, and A. Turkoglu, "Comparative analysis of the protective effects of melatonin and vitamin E on streptozocin-induced diabetes mellitus," *Journal of Pineal Research*, vol. 32, no. 4, pp. 225–230, 2002.
 - [51] M. H. Abdel Wahab, E.-S. E. M. S. Akoul, and A.-A. H. Abdel-Aziz, "Modulatory effects of melatonin and vitamin E on doxorubicin-induced cardiotoxicity in Ehrlich ascites carcinoma-bearing mice," *Tumori*, vol. 86, no. 2, pp. 157–162, 2000.
 - [52] P. Montilla, A. Cruz, F. J. Padillo, I. Túnez, F. Gascon, and M. C. Muñoz, "Melatonin versus vitamin E as protective treatment against oxidative stress after extra-hepatic bile duct ligation in rats," *Journal of Pineal Research*, vol. 31, pp. 138–144, 2001.
 - [53] C.-H. Hsu, B.-C. Han, M.-Y. Liu, C.-Y. Yeh, and J. E. Casida, "Phosphine-induced oxidative damage in rats: attenuation by melatonin," *Free Radical Biology and Medicine*, vol. 28, no. 4, pp. 636–642, 2000.
 - [54] F. Gultekin, N. Delibas, S. Yasar, and I. Kilinc, "In vivo changes in antioxidant systems and protective role of melatonin and a combination of vitamin C and vitamin E on oxidative damage in erythrocytes induced by chlorpyrifos-ethyl in rats," *Archives of Toxicology*, vol. 75, pp. 88–96, 2001.
 - [55] S. Rosales-Corral, D. X. Tan, R. J. Reiter, M. Valdivia-Velázquez, G. Martínez-Barboza, and J. P. Acosta-Martínez, "Orally administered melatonin reduces oxidative stress and proinflammatory cytokines induced by amyloid-beta peptide in rat brain: a comparative, in vivo study versus vitamin C and E," *Journal of Pineal Research*, vol. 35, pp. 80–84, 2003.
 - [56] C. M. Deaton and D. J. Marlin, "Exercise-associated oxidative stress," *Clinical Techniques in Equine Practice*, vol. 2, no. 3, pp. 278–291, 2003.
 - [57] N. Ilhan, A. Kamanli, R. Ozmerdivenli, and N. Ilhan, "Variable effects of exercise intensity on reduced glutathione, thiobarbituric acid reactive substance levels, and glucose concentration," *Archives of Medical Research*, vol. 35, pp. 294–300, 2004.
 - [58] K. P. Skenderi, M. Tsironi, C. Lazaropoulou et al., "Changes in free radical generation and antioxidant capacity during ultra-marathon foot race," *European Journal of Clinical Investigation*, vol. 38, no. 3, pp. 159–165, 2008.

- [59] C. Leeuwenburgh and J. W. Heinecke, "Oxidative stress and antioxidants in exercise," *Current Medicinal Chemistry*, vol. 8, no. 7, pp. 829–838, 2001.
- [60] W. J. Kraemer, C. C. Loebel, J. S. Volek et al., "The effect of heavy resistance exercise on the circadian rhythm of salivary testosterone in men," *European Journal of Applied Physiology*, vol. 84, no. 1-2, pp. 13–18, 2001.
- [61] S. P. Bird and K. M. Tarpenning, "Influence of circadian time structure on acute hormonal responses to a single bout of heavy-resistance exercise in weight-trained men," *Chronobiology International*, vol. 21, pp. 131–146, 2004.
- [62] G. A. E. Ponjee, H. A. M. De Rooy, and H. L. Vader, "Androgen turnover during marathon running," *Medicine and Science in Sports and Exercise*, vol. 26, no. 10, pp. 1274–1277, 1994.
- [63] A. Luger, P. A. Deuster, S. B. Kyle, W. T. Gallucci, L. C. Montgomery, and P. W. Gold, "Acute hypothalamic-pituitary-adrenal responses to the stress of treadmill exercise. Physiologic adaptations to physical training," *The New England Journal of Medicine*, vol. 316, pp. 1309–1315, 1987.
- [64] D. B. Carr, S. M. Reppert, B. Bullen, G. Skrinar, I. Beitins, and M. Arnold, "Plasma melatonin increases during exercise in women," *Journal of Clinical Endocrinology & Metabolism*, vol. 53, pp. 224–225, 1981.
- [65] J. J. Theron, J. M. C. Oosthuizen, and M. M. Rautenbach, "Effect of physical exercise on plasma melatonin levels in normal volunteers," *South African Medical Journal*, vol. 66, no. 22, pp. 838–841, 1984.
- [66] P. Monteleone, M. Maj, M. Fusco, C. Orasso, and D. Kemali, "Physical exercise at night blunts the nocturnal increase of plasma melatonin levels in healthy humans," *Life Sciences*, vol. 47, no. 22, pp. 1989–1995, 1990.
- [67] P. Monteleone, M. Maj, A. Fuschino, and D. Kemali, "Physical stress in the middle of the dark phase does not affect light-depressed plasma melatonin levels in humans," *Neuroendocrinology*, vol. 55, no. 4, pp. 367–371, 1992.
- [68] K. Yaga, D.-X. Tan, R. J. Reiter, L. C. Manchester, and A. Hattori, "Unusual responses of nocturnal pineal melatonin synthesis and secretion to swimming: attempts to define mechanisms," *Journal of Pineal Research*, vol. 14, no. 2, pp. 98–103, 1993.
- [69] T. Miyazaki, S. Hashimoto, S. Masubuchi, S. Honma, and K.-I. Honma, "Phase-advance shifts of human circadian pacemaker are accelerated by daytime physical exercise," *American Journal of Physiology*, vol. 281, no. 1, pp. R197–R205, 2001.
- [70] A. N. Elias, A. F. Wilson, M. R. Pandian, F. J. Rojas, R. Kayaleh, and S. C. Stone, "Melatonin and gonadotropin secretion after acute exercise in physically active males," *European Journal of Applied Physiology and Occupational Physiology*, vol. 66, pp. 357–361, 1993.
- [71] G. A. Laughlin, A. B. Loucks, and S. S. C. Yen, "Marked augmentation of nocturnal melatonin secretion in amenorrheic athletes, but not in cycling athletes: unaltered by opioidergic or dopaminergic blockade," *Journal of Clinical Endocrinology and Metabolism*, vol. 73, no. 6, pp. 1321–1326, 1991.
- [72] D. Martarelli and P. Pompei, "Oxidative stress and antioxidant changes during a 24-hours mountain bike endurance exercise in master athletes," *Journal of Sports Medicine and Physical Fitness*, vol. 49, no. 1, pp. 122–127, 2009.
- [73] I. McIntyre, T. R. Norman, G. D. Burrows, and S. M. Armstrong, "Human melatonin suppression by light is intensity dependent," *Journal of Pineal Research*, vol. 6, pp. 149–156, 1988.
- [74] J. Trinder, S. M. Armstrong, C. O'Brien, D. Luke, and M. J. Martin, "Inhibition of melatonin secretion onset by low levels of illumination," *Journal of Sleep Research*, vol. 5, no. 2, pp. 77–82, 1996.
- [75] E. Baron Short, S. Kose, Q. Mu, J. Borckardt, A. Newberg, and M. S. George, "Regional brain activation during meditation shows time and practice effects: an exploratory fMRI study," *Evidence-Based Complementary and Alternative Medicine*. In press.
- [76] N. Ilhan, A. Kamanli, R. Ozmerdivenli, and N. Ilhan, "Variable effects of exercise intensity on reduced glutathione, thiobarbituric acid reactive substance levels, and glucose concentration," *Archives of Medical Research*, vol. 35, pp. 294–300, 2004.
- [77] M. R. Cesarone, G. Belcaro, M. Carratelli et al., "A simple test to monitor oxidative stress," *International Angiology*, vol. 18, no. 2, pp. 127–130, 1999.
- [78] A. Alberti, L. Bolognini, D. Macciantelli, and M. Caratelli, "The radical cation of N,N-diethyl-para-phenyldiamine: a possible indicator of oxidative stress in biological samples," *Research on Chemical Intermediates*, vol. 26, no. 3, pp. 253–267, 2000.
- [79] K. Dohi, K. Satoh, H. Ohtaki et al., "Elevated plasma levels of bilirubin in patients with neurotrauma reflect its pathophysiological role in free radical scavenging," *In Vivo*, vol. 19, no. 5, pp. 855–860, 2005.
- [80] T. A. Watson, R. Callister, R. D. Taylor, D. W. Sibbritt, L. K. Macdonald-Wicks, and M. L. Garg, "Antioxidant restriction and oxidative stress in short-duration exhaustive exercise," *Medicine and Science in Sports and Exercise*, vol. 37, no. 1, pp. 63–71, 2005.
- [81] M. J. Alcaraz-García, M. D. Albaladejo, C. Acevedo et al., "Effects of hyperoxia on biomarkers of oxidative stress in closed-circuit oxygen military divers," *Journal of Physiology and Biochemistry*, vol. 64, no. 2, pp. 135–142, 2008.
- [82] M. Phillips, R. N. Cataneo, J. Greenberg, R. Grodman, R. Gunawardena, and A. Naidu, "Effect of oxygen on breath markers of oxidative stress," *European Respiratory Journal*, vol. 21, no. 1, pp. 48–51, 2003.
- [83] R. J. Reiter, D. X. Tan, L. C. Manchester, M. Pilar Terron, L. J. Flores, and S. Koppisepi, "Medical implications of melatonin: receptor-mediated and receptor-independent actions," *Advances in Medical Sciences*, vol. 52, pp. 11–28, 2007.
- [84] D. Orme-Johnson, "Medical care utilization and the transcendental meditation program," *Psychosomatic Medicine*, vol. 49, no. 5, pp. 493–507, 1987.
- [85] D. Orme-Johnson, "Evidence that the Transcendental Meditation program prevents or decreases diseases of the nervous system and is specifically beneficial for epilepsy," *Medical Hypotheses*, vol. 67, no. 2, pp. 240–246, 2006.
- [86] M. Mourya, A. S. Mahajan, N. P. Singh, and A. K. Jain, "Effect of slow- and fast-breathing exercises on autonomic functions in patients with essential hypertension," *Journal of Alternative and Complementary Medicine*, vol. 15, no. 7, pp. 711–717, 2009.
- [87] K. Yang, L. M. Bernardo, S. M. Sereika, M. B. Conroy, J. Balk, and L. E. Burke, "Utilization of 3-month Yoga program for adults at high risk for type 2 diabetes: a pilot study," *Evidence-Based Complementary and Alternative Medicine*. In press.
- [88] J. D. Adams Jr. and C. Garcia, "Palliative care among Chumash people," *Evidence-Based Complementary and Alternative Medicine*, vol. 2, no. 2, pp. 143–147, 2005.
- [89] K. Yang, "A review of yoga programs for four leading risk factors of chronic diseases," *Evidence-Based Complementary and Alternative Medicine*, vol. 4, no. 4, pp. 487–491, 2007.

- [90] S. Khalfa, S. D. Bella, M. Roy, I. Peretz, and S. J. Lupien, "Effects of relaxing music on salivary cortisol level after psychological stress," *Annals of the New York Academy of Sciences*, vol. 999, pp. 374–376, 2003.
- [91] R. Jevning, A. F. Wilson, W. R. Smith, and M. E. Morton, "Redistribution of blood flow in acute hypometabolic behavior," *American Journal of Physiology*, vol. 4, no. 1, pp. R89–R92, 1978.
- [92] R. Lang, K. Dehof, K. A. Meurer, and W. Kaufmann, "Sympathetic activity and transcendental meditation," *Journal of Neural Transmission*, vol. 44, no. 1-2, pp. 117–135, 1979.
- [93] M. Bujatti and P. Riederer, "Serotonin, noradrenaline, dopamine metabolites in transcendental meditation technique," *Journal of Neural Transmission*, vol. 39, no. 3, pp. 257–267, 1976.
- [94] R. Y. Moore, "The innervation of the mammalian pineal gland," *Progress in Reproductive Biology*, vol. 4, pp. 1–29, 1978.
- [95] N. Zisapel, R. Tarrasch, and M. Laudon, "The relationship between melatonin and cortisol rhythms: clinical implications of melatonin therapy," *Drug Development Research*, vol. 65, no. 3, pp. 119–125, 2005.
- [96] P. Monteleone, A. Fuschino, G. Nolfi, and M. Maj, "Temporal relationship between melatonin and cortisol responses to nighttime physical stress in humans," *Psychoneuroendocrinology*, vol. 17, no. 1, pp. 81–86, 1992.